



Fungicide

(The Original 76% Water Dispersible Granules)

NOTIFICATION

ACTIVE INGREDIENT:

Ziram (zinc dimethyl dithiocarbamate)*.....76.0%

OTHER INGREDIENTS:.....24.0%

TOTAL 100.0%

NOV 30 2004

*Zinc expressed as metallic.....16.25%

EPA Reg. No. 45728-12

EPA Est. No. 32557-BEL-1

EPA Est. No. 72178-SC-001

**KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO**

**Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)**

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergencies involving a Spill, Leak, Fire, Exposure, or Accident, Contact: CHEMTREC at (800) 424-9300. For product usage information, phone Taminco, Inc., toll free at (800) 426-3820 from 9:00 AM to 5:00 PM Eastern time.	
NOTE TO PHYSICIAN	
Possible mucosal damage may contraindicate gastric lavage.	

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks
- Protective eye wear
- Chemical-resistant headgear for overhead exposure
- Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protection Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering control statements:
When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:
User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS
This product is toxic to fish and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS
Keep away from fire and sparks.
GENERAL USE INFORMATION

Ziram Granuflo is a broad spectrum fungicide for the control of certain diseases in fruit and nut crops, and ornamentals. Taminco, Inc.'s recommendations for the use of this product are based upon tests believed to be reliable. Ziram Granuflo is not for residential use.
FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.
Read entire label carefully and use only as directed.

See Other Panels for Additional Precautionary Statements.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER: Corrosive, causes eye damage. **May be fatal if inhaled.** Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed or absorbed through skin.
PERSONAL PROTECTIVE EQUIPMENT (PPE)
Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistant category selection chart.
Applicators and other handlers must wear:

DIRECTIONS FOR USE

3/10

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eye wear
- Chemical-resistant headgear for overhead exposure

CHEMIGATION: Do not apply this product through any type of irrigation system.

DIRECTIONS AND RECOMMENDATIONS:

Use recommended rates for concentrate, dilute and aerial applications. For dilute application, use Ziram Granuflo® at the recommended rates per 100 gallons. Ground application is preferred, especially when full canopy of foliage exists. ZIRAM GRANUFLO® may be applied as a concentrate in a minimum of 20 gallons or more of spray solution per acre by ground spray equipment or in a minimum of 10 gallons or more per acre by aircraft equipment. Increase spray volume and fungicide rate (according to labeling specifications) as crop growth increases to ensure thorough coverage and adequate protection of the foliage and fruit. Under low disease conditions, minimum label rates per application should be used, however, maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions.

Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use. Attention should be given to sprayer speed and calibration, wind speed, and tree and vine canopy to ensure adequate spray coverage with ZIRAM GRANUFLO®.

For dilute application, use ZIRAM GRANUFLO® at the recommended rates per 100 gallons. For concentrate ground application, use ZIRAM GRANUFLO® at the recommended rate per acre. The following should be used to determine the equivalent amount of product per acre in ½X, 1X, 2X, 3X, 4X (where X = per 100 gal), etc. spray solutions:

POUNDS ZIRAM GRANUFLO® FUNGICIDE PER ACRE FOR DILUTE AND CONCENTRATE SPRAYS			
POME FRUIT (APPLE)		STONE FRUIT (PEACH & NECTARINE)	
½X or 50 gal/acre	6 - 8 lbs	½X or 50 gal/acre	3.75 - 8 lbs
1X or 100 gal/acre	6 - 8 lbs	1X or 100 gal/acre	3.75 - 8 lbs
2X or 200 gal/acre	6 - 8 lbs	2X or 200 gal/acre	3.75 - 8 lbs
3X or 300 gal/acre	6 - 8 lbs	3X or 300 gal/acre	3.75 - 8 lbs
4X or 400 gal/acre	6 - 8 lbs	4X or 400 gal/acre	3.75 - 8 lbs

Where ZIRAM GRANUFLO® rate per acre is not available, the following can be used to determine dilute and concentrate sprays per acre use rates based on per 100 gallon volume use rates.

Examples:

- 1) A vineyard consisting of mature grapevines requires 3 to 4 pounds of ZIRAM GRANUFLO® for adequate black rot, downy mildew, phomopsis cane and leaf spot, ripe rot and botrytis bunch rot control. The following table reflects how the rates are equivalent comparing both dilute and concentrate applications:

POUNDS ZIRAM GRANUFLO FUNGICIDE DILUTE AND CONCENTRATE SPRAYS BASED ON PER 100 GALLON RECOMMENDATION		
CROP	DILUTE (PER 100 GALLONS)	CONCENTRATE (PER 20-100 GALLONS)
Grape (Eastern U.S.)	3 - 4 lbs	3 - 4 lbs

- 2) A blueberry field consisting of mature blueberries requires 3 pounds of ZIRAM GRANUFLO® for adequate mummyberry control. The following table reflects how the rates are equivalent comparing both dilute and concentrate applications:

4/10

POUNDS ZIRAM GRANUFLO FUNGICIDE DILUTE AND CONCENTRATE SPRAYS BASED ON PER 100 GALLON RECOMMENDATION		
CROP	DILUTE (PER 100 GALLONS)	CONCENTRATE (PER 20-100 GALLONS)
Blueberries	3 lbs	3 lbs

These rate recommendations can be used in concentrate, dilute and aerial applications. Ground application is preferred, especially when full canopy or foliage exists.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Make new dilutions for each spraying. If a spreader/sticker is added to the finished spray mixture, do not use for pre-harvest sprays. Also, it is not recommended that a spreader/sticker be used on fruit that will not be washed following harvest and prior to commercial pack-out. The use of a spreader/sticker for applications to apricots is not recommended for risk of adverse effects to fruit finish.

MIXING INSTRUCTIONS

To mix, fill the spray tank about 2/3 full of water, and with agitator running, slowly sift in the required amount of Ziram Granuflo®. Finish filling the tank with water and let mixing continue in the tank until thoroughly mixed. Continue agitation and apply immediately.

Do not graze or feed cover crops from treated orchards.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Keep away from fire and sparks.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. Do not put concentrate or dilute material into food or drink containers.

In case of spill: Avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

For Emergencies Involving a Spill, Leak, Fire, Exposure, or Accident, contact: CHEMTREC at (800) 424-9300. For product usage information, phone Taminco, Inc., toll free at (800) 426-3820 from 9:00 AM to 5:00 PM Eastern time.

Confine spill: If liquid, dike surrounding area or absorb with sand, cat litter, or commercial clay. Sweep up material and dispose according to Pesticide Disposal. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TERMS OF SALE OR USE

On purchase of this product, buyer and user agree to the following conditions:

Warranty: Taminco, Inc. makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Ziram Granuflo® is a Registered Trademark of Taminco N.V., a Belgian corporation.

NET WEIGHT:



Taminco, Inc.
1950 Lake Park Drive
Smyrna, GA 30080

ZIRAM GRANUFLO APPLICATION INFORMATION SPECIFIC TO CROP

5/10

FRUIT AND NUT TREE CROPS	DISEASES	POUNDS ZIRAM GRANUFLO®		DIRECTIONS FOR USE
		per acre	per 100 gal	
ALMONDS	Brown Rot Blossom Blight and Twig Blight (<i>Monilinia</i> spp.)	8 lbs	2 2/3 lbs ⁽³⁾	Apply pre-bloom through petal fall periods. Do not apply later than 5 weeks after petal fall. Do not apply more than 32 lbs/A per crop cycle. ⁽⁴⁾
	Leaf Blight (<i>Seimatosporium lichenicola</i>)			
	Scab (<i>Cladosporium carpophilum</i>)			
	Shot Hole (<i>Coryneum</i> Blight) (<i>Stigmina carpophilum</i>)			
APPLES	Apple Scab (<i>Venturia inaequalis</i>)	6 to 8 lbs	1 1/2 to 2 lbs ⁽²⁾	Apply from pre-bloom through early cover sprays as needed. Do not apply within 14 days of harvest. Eastern U.S.: Do not apply more than 56 lbs/A per crop cycle. ⁽⁴⁾ Western U.S.: Do not apply more than 32 lbs/A per crop cycle. ⁽⁴⁾
	Cedar Apple Rust⁽¹⁾, Quince Rust⁽¹⁾ (<i>Gymnosporangium</i> spp.)			
	Sooty Blotch⁽¹⁾ (<i>Gloeodes pomigena</i>)			
	Fly Speck⁽¹⁾ (<i>Schizothyrium perexiguum</i>)			
	Bitter Rot⁽¹⁾ (<i>Glomerella cingulata</i>)			
	Bull's-Eye Rot (Pacific Northwest only) (<i>Pezizula malicorticis</i>)	6 to 8 lbs	1 1/2 to 2 lbs ⁽²⁾	Apply 12 to 14 days after calyx stage and repeat as necessary in summer sprays. Do not apply within 14 days of harvest. Do not apply more than 32 lbs/A per crop cycle. ⁽⁴⁾
	Perennial Canker (Oregon only) (<i>Neofabraea perennans</i>)	6 to 8 lbs	1 1/2 to 2 lbs ⁽²⁾	Apply pre-bloom and calyx sprays. Do not apply more than 32 lbs/A per crop cycle. ⁽⁴⁾
APRICOTS	Brown Rot Blossom Blight, Twig Blight and Fruit Rot (<i>Monilinia</i> spp.)	6 to 8 lbs (except CA) 8 lbs (CA only)	1 1/2 to 2 2/3 lbs ⁽³⁾ 2 2/3 lbs	Apply in pre-bloom, bloom and petal fall through early cover sprays. Do not apply within 30 days of harvest. Do not apply more than 40 lbs/A per crop cycle. ⁽⁴⁾ The use of spreader/stickers for applications to apricots is not recommended based on risk of adverse effects to fruit finish.
CHERRIES Eastern U.S.	Brown Rot Blossom Blight, Twig Blight and Fruit Rot (<i>Monilinia</i> spp.)	5 to 8 lbs	1 2/3 to 2 2/3 lbs ⁽³⁾	Apply in pre-bloom, bloom, petal fall and shuck stages, and approximately 2 weeks after shuck fall. Do not apply within 14 days of harvest. Do not apply more than 40 lbs/A per crop cycle. ⁽⁴⁾
	Leaf Spot (<i>Blumeriella jaapii</i>)			
CHERRIES Western U.S.	Brown Rot Blossom Blight, Twig Blight and Fruit Rot (<i>Monilinia</i> spp.)	4 to 6 lbs (except CA) 4 to 5 lbs (CA only)	1 1/3 to 2 lbs ⁽³⁾ 1 1/3 to 1 2/3 lbs ⁽³⁾	Apply at pre-bloom, bloom, petal fall and shuck stages, and approximately 2 weeks after shuck fall. Do not apply more than 30 lbs/A per crop cycle. ⁽⁴⁾ Do not apply within 30 days of harvest. (except California) Do not apply within 7 days of harvest. (California only)
PEACHES AND NECTARINES Eastern U.S.	Brown Rot Blossom Blight, Twig Blight and Fruit Rot (<i>Monilinia</i> spp.)	4 1/2 to 8 lbs	1 1/2 to 2 2/3 lbs ⁽³⁾	Apply at pre-bloom through cover sprays as needed. Do not apply within 14 days of harvest. Do not apply more than 72 lbs/A per crop cycle. ⁽⁴⁾
	Peach Blight (i.e. Shot Hole) (<i>Stigmina carpophila</i>)	4 1/2 to 8 lbs	1 1/2 to 2 2/3 lbs ⁽³⁾	Apply at pre-bloom through cover sprays as needed. In addition to spray program, apply 8 lbs/A in the fall before the heavy rains. Do not apply within 14 days of harvest. Do not apply more than 72 lbs/A per crop cycle. ⁽⁴⁾
	Peach Leaf Curl (<i>Taphrina deformans</i>)	3 3/4 to 8 lbs	1 1/4 to 2 2/3 lbs ⁽³⁾	Make dormant applications after leaf drop and/or prior to bud swell. Do not apply more than 72 lbs/A per crop cycle. ⁽⁴⁾
PEACHES AND NECTARINES Western U.S.	Brown Rot Blossom Blight, Twig Blight and Fruit Rot (<i>Monilinia</i> spp.)	6 to 8 lbs (except CA) 8 lbs (CA only)	1 1/2 to 2 2/3 lbs ⁽³⁾ 2 2/3 lbs ⁽³⁾	Apply at pre-bloom through cover sprays as needed. Do not apply within 30 days of harvest. Do not apply more than 56 lbs/A per crop cycle. ⁽⁴⁾
	Peach Blight (i.e. Shot Hole) (<i>Stigmina carpophila</i>)	6 to 8 lbs (except CA) 8 lbs (CA only)	1 1/2 to 2 2/3 lbs ⁽³⁾ 2 2/3 lbs ⁽³⁾	Apply at pre-bloom through cover sprays as needed. In addition to preharvest spray program, apply 8 lbs/A in the fall (after harvest) before the heavy rains. Do not apply within 30 days of harvest. Do not apply more than 56 lbs/A per crop cycle. ⁽⁴⁾
	Peach Leaf Curl (<i>Taphrina deformans</i>)	8 to 10 lbs (except CA) 8 lbs (CA only)	2 2/3 to 3 1/3 lbs ⁽³⁾ 2 2/3 lbs ⁽³⁾	Make dormant applications in late fall after leaf drop and/or in late winter prior to bud swell. Do not apply more than 56 lbs/A per crop cycle. ⁽⁴⁾

(1) Not for use in California.
 Based on dilute sprays of 400 gallons per acre.
 (2) Based on dilute sprays of 300 gallons per acre.
 (4) Crop cycle is defined as pre-bloom through post-harvest.

ZIRAM GRANUFLO APPLICATION INFORMATION SPECIFIC TO CROP

6/10

FRUIT AND NUT

POUNDS ZIRAM GRANUFLO

TREE CROPS	DISEASES	per acre	per 100 gal	DIRECTIONS FOR USE
PEARS	Pear Scab (<i>Venturia pyrina</i>) Quince Rust ⁽¹⁾ (<i>Gymnosporangium</i> spp.) Sooty Blotch ⁽¹⁾ (<i>Gloeodes pomigena</i>) Fly Speck ⁽¹⁾ (<i>Schizothyrium pomi</i>) Blitter Rot ⁽¹⁾ (<i>Glomerella cingulata</i>) Fabraea Leaf Spot ⁽¹⁾ (<i>Diplocarpon mespili</i>)	6 to 8 lbs	1 1/2 to 2 lbs ⁽²⁾	Apply from pre-bloom through cover sprays as needed. Eastern U.S.: Do not apply within 14 days of harvest. Do not apply more than 56 lbs/A per crop cycle. ⁽⁴⁾ Western U.S.: Do not apply within 5 days of harvest. Do not apply more than 32 lbs/A per crop cycle. ⁽⁴⁾
	Bull's-Eye Rot (Pacific Northwest) (<i>Pezizicia malicorticis</i>)	6 to 8 lbs	1 1/2 to 2 lbs ⁽²⁾	Apply in the first cover spray and in pre-harvest spray before fall rains begin. Repeat as necessary, especially if excessive rainfall occurs. Do not apply within 5 days of harvest. Do not apply more than 32 lbs/A per crop cycle. ⁽⁴⁾
	Storage Rot (Oregon only) (<i>Phialophora malorum</i>)	6 to 8 lbs	1 1/2 to 2 lbs ⁽²⁾	Apply as a pre-harvest spray. Do not apply within 5 days of harvest. Do not apply more than 32 lbs/A per crop cycle. ⁽⁴⁾
PECANS	Pecan Scab (<i>Cladosporium effusum</i>) Anthracnose (<i>Glomerella cingulata</i>)	6 to 8 lbs	1 1/2 to 2 2/3 lbs ⁽²⁾	Apply up to 3 times beginning when leaves are 1/4 to 1/2 in. long through pollination. Apply up to 5 more cover sprays at 3 to 4 week intervals beginning after pollination. Do not apply within 55 days of harvest. Do not apply more than 64 lbs/A per crop cycle. ⁽⁴⁾
ORNAMENTAL CROPS(1)				
Asters	Rust (<i>Puccinia</i> spp.)		1 1/2 to 2 lbs	Begin application early in the growing season and continue every 7 days as needed until blossoms form.
Azaleas Camellias	Petal Blight (<i>Pestalotia</i> spp.)		1 1/2 to 2 lbs	Begin application with first bloom. Apply at 3 to 5 day intervals as a drench to mulch and litter beneath and around bushes through bloom period.
Carnations Chrysanthemums Marigolds Morning Glory Snapdragons	Rust (<i>Puccinia</i> spp.)		1 2/3 to 2 lbs	Apply at 7 to 10 day intervals as needed throughout the growing season.
Carnations Chrysanthemums Dahlias Gladiolus Snapdragons	Leaf Spot (<i>Alternaria</i> spp.)		1 2/3 to 2 lbs	Apply at 7 day intervals as needed throughout the growing season.
Cyclamens Gardenias	Botrytis Blight (<i>Botrytis</i> spp.) Leaf Spot (<i>Phyllostica</i> spp.)		1 1/2 to 2 lbs	Apply at 7 to 10 day intervals as needed throughout the growing season.
Dahlias Delphinium Golden Glow Hollyhocks Phlox	Powdery Mildew (<i>Erysiphe cichoracearum</i>)		1 2/3 to 2 lbs	Apply at 7 day intervals as needed throughout the growing season.
Lilies Peonies Poinsettias Tulips	Botrytis Blight (<i>Botrytis</i> spp.)		1 1/2 to 2 lbs	Apply at 7 to 10 day intervals as needed throughout the growing season.
Roses	Black Spot (<i>Diplocarpon rosae</i>) Powdery Mildew (<i>Sphaerotheca</i> spp.)		1 2/3 to 2 lbs	Begin application when leaves unfold. Apply at 7 to 10 day intervals as needed throughout the growing season.

(1) Not for use in California.

(2) Based on dilute sprays of 400 gallons per acre.

Based on dilute sprays of 300 gallons per acre.

(4) Crop cycle is defined as pre-bloom through post-harvest.

ZIRAM GRANUFLO APPLICATION INFORMATION SPECIFIC TO CROP

7/10

ORNAMENTAL TREES AND FRUIT(1)	DISEASES	POUNDS ZIRAM GRANUFLO		DIRECTIONS FOR USE
		per acre	per 100 gal	
Alfalfa Pine Seedlings	Fusiform Rust		2 lbs	Apply at 3 to 5 day intervals from plant emergence to mid-June.
Douglas Fir Christmas Trees	Melampsora Needle Rust		2 lbs	Make one application in the spring when new shoot growth is 1/2 to 2 inches in length.
Shasta Fir Christmas Trees	Puccineastrum Needle Rust		2 lbs	Make a single application in the spring when new shoot growth is 1/2 to 2 inches in length.
Apple Pear	Anthracnose (<i>Pezicula malicorticis</i>) European (Nectria) Canker Fusiform Rust		2 to 3 lbs	Apply as a thorough coverage spray in the fall and dormant periods. DO NOT USE FRUIT FOR HUMAN CONSUMPTION.
Flowering Crab Apple Oregon Crab Apple Quince	Anthracnose (<i>Pezicula malicorticis</i>)		2 to 3 lbs	Apply as a thorough coverage spray in the fall and dormant periods. DO NOT USE FRUIT FOR HUMAN CONSUMPTION.
Sweet Cherry Wild Cherry Choke Cherry Cherry laurel Flowering Plum (non-edible only)	Witches' Broom, Leaf Curl, Plum Pockets (<i>Taphrina</i> spp.)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell. DO NOT USE FRUIT FOR HUMAN CONSUMPTION.
Peach Apricot	Twig Blight (<i>Monilinia</i> spp.) Leaf Curl (<i>Taphrina deformans</i>)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell. DO NOT USE FRUIT FOR HUMAN CONSUMPTION.
Nectarine	Leaf Curl (<i>Taphrina deformans</i>) Shot Hole (<i>Stigmata carpophila</i>)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell. DO NOT USE FRUIT FOR HUMAN CONSUMPTION.
American Hornbeam	Leaf Blister (<i>Taphrina australis</i>)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell.
Birch Maple	Leaf Blister (<i>Taphrina</i> spp.)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell.
California Buckeye	Yellow Leaf Blister (<i>Taphrina aesculi</i>)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell.
Oak	Leaf Blister (<i>Taphrina caerulescens</i>)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell.
Alder	Leaf Curl, Catkin Hypertrophy (<i>Taphrina</i> spp.)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell.
Poplar	Yellow Leaf Blister (<i>Taphrina populina</i>) Shoot Blight (<i>Venturia populina</i>)		2 to 3 lbs	Apply as a thorough coverage spray in the dormant period prior to bud swell.

(1) Not for use in California.

ZIRAM GRANUFLO APPLICATION INFORMATION SPECIFIC TO CROP

8/16

SMALL FRUIT AND VEGETABLE CROPS	DISEASES	POUNDS ZIRAM GRANUFLO®		DIRECTIONS FOR USE
		per acre	per 100 gal	
Blackberries ⁽¹⁾	Evergreen blackberry canker	3 lbs	3 lbs	Apply in sufficient water for thorough coverage. Make one application between mid-June and early July.
Blueberries ⁽¹⁾	Mummyberry (Monilinia vaccinii-corymbosi)	3 lbs	3 lbs	Apply in sufficient water for thorough coverage. Apply at loose bud scale stage and 7 days later. Do not apply later than 3 weeks after full bloom.
Grape Eastern U.S. (East of the Rockies)	Black Rot (Guignardia bidwellii) Downy Mildew (Plasmopara viticola) Phomopsis Cane and Leaf Spot (Phomopsis viticola) Ripe Rot (Glomerella cingulata) Botrytis Bunch Rot (Botrytis cinerea) (aides in control)	3 to 4 lbs	3 to 4 lbs	Begin applications when shoots are at least 1 inch long and continue at 7-14 day intervals or as necessary according to the integrated pest management guidelines for the particular growing area. Use shorter intervals under heavy disease pressure. Do not apply more than 28 lbs per acre per crop cycle. Do not apply later than 21 days to harvest.
Grape Western U.S. (West of the Rockies)	Black Rot (Guignardia bidwellii) Downy Mildew (Plasmopara viticola) Phomopsis Cane and Leaf Spot (Phomopsis viticola) Ripe Rot (Glomerella cingulata) Botrytis Bunch Rot (Botrytis cinerea) (aides in control)	3 to 4 lbs	3 to 4 lbs	Apply when shoots are ½ to 1½ inches long and repeat at 7-10 day intervals as needed. Do not apply after bloom.
Tomato ⁽¹⁾ (Do not use on Cherry tomatoes)	Anthracnose (Colletotrichum coccodes) Early Blight (Alternaria solani) Septoria Leaf Spot (Septoria lycopersici)	3 to 4 lbs	3 to 4 lbs	Begin applications at first sign of infection and continue at 7-14 day intervals. Use the shorter spray interval under high disease pressure. Do not apply more than 24 lbs per acre per crop cycle. Do not apply within 7 days of harvest. Ziram can be mixed with copper fungicides to enhance bacterial disease control.

(1) Not for use in California.

ChemReg®

INTERNATIONAL, LLC

9/10

1990 OLD BRIDGE ROAD, SUITE 201
LAKE RIDGE, VIRGINIA 22192-3026

DIRECT: 703-492-0753
MAIN: 703-492-0445
Fax: 703-492-0668

E-MAIL: spanogle@chemreg.com
WEB SITES: www.chemreg.com
www.pesticide.net

TERRI L. SPANOGLA

Hand Delivered

November 16, 2004

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
Room 266A, Crystal Mall 2
1801 Bell Street
Arlington, VA 22202-4501

Attn.: Cynthia Giles-Parker
Product Manager (22)

Subject: Notification to Correct the Current Label Use Pattern for Blueberries for Ziram Granuflo Fungicide (EPA Reg. No. 45728-12) Per PR Notice 98-10

Dear Cynthia:

On behalf of Taminco, Inc., please find enclosed a notification correcting the current use profile for Ziram Granuflo Fungicide (EPA Reg. No. 45728-12) for blueberries because of information that was inadvertently added to the label.

Taminco, Inc. submitted an amendment on May 14, 2003, to address minor changes to the "Precautionary Statements" and "Environmental Hazards" sections of the label. EPA accepted this label on September 29, 2003. However, we have since realized that the September 29, 2003, EPA stamp-accepted label inadvertently contains additional diseases, a footnote indicating an application rate of 4 lbs per acre for the disease, *Botrytis*, and directions for use that were not approved previously for blueberries.

CONSULTANTS TO SUCCESS®

Expertise in Toxicology, Residue and Product Chemistry, Environmental Fate, Efficacy, Registration Strategies and Implementation, Regulatory Compliance, Label Development, Exposure and Risk Assessments, Data Citation and Compensation, Study Design and Monitoring.

10/10

Cynthia Giles-Parker
November 16, 2004
Page 2


We are submitting this notification to correct the erroneous label by deleting all but one of the diseases (Mummyberry) and revising the directions for use language for blueberries to reflect the correct use profile.

Please find enclosed:

1. EPA Form 8570-1: Application for Registration
2. One copy of the new label and one copy of highlighted changes for Ziram Granuflo® Fungicide.

If you have any questions, please call me at (703) 492-0753.

Sincerely,



Terri Spanogle
Scientific and Regulatory Consultant
ChemReg International, LLC
Authorized Representative
Taminco, Inc.

Enclosures

cc: Jean-Michel Denis, Taminco, Inc.
Mia Laget, Taminco N.V.

Label code: Ziram Granuflo 11/16/04.TLS(EPA09/29/03, CDPR12/05/01)