

MAR 28 1997

Ms. Kristine M. Hooks
 Agent
 UCB Chemical Corporation
 2001 Jefferson Davis Highway, Suite 1010
 Arlington, VA 22202

Dear Ms. Hooks:

Subject: Thiram 75W
 EPA Registration No. 45728-21
 Application Dated March 21, 1997, Request
 To Amend Registration by Splitting Label
 into Four (4) Labels for Marketing and
 Revising Labeling To Clarify Use Directions
 and Application Rates, and the Addition of
 Latin names of Disease-Causing Fungi

The subject proposed labeling amendments have been reviewed and found acceptable for registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended, provided that you:

1. Add the following Precautionary Statement to each of the four (4) labels:

"Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals".

2. Submit one (1) copy of the final printed labeling before you release the product for shipment under the subject labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the product under the subject labeling constitutes acceptance of these conditions.

Stamped copies of the labeling are enclosed for your records.

Sincerely yours,

BEST AVAILABLE COPY

CONCURRENCES		Joanne I. Miller	
SYMBOL		Product Manager (23)	
SURNAME	.Wilson:Diskette Thiram 03-28-97	Fungicide-Herbicide Branch	
DATE	Enclosures (4)	Registration Division (7505C)	

20/12



Thiram Granuflo®

(The Original 75% Water Dispersible Granules)
Fungicide

ACCEPTED
with COMMENTS
In EPA Letter Dated

ACTIVE INGREDIENT:
Thiram (Tetramethylthiuram disulfide)..... 75.0%
INERT INGREDIENTS: 25.0%
TOTAL 100.0%
EPA Reg. No. 45728-21
EPA Est. No. 32557-BEL-1

MAR 28 1987

KEEP OUT OF REACH OF CHILDREN

CAUTION

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
45728-21

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration, if indicated.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling spray mist. Do not use or store near food or feed.

The consumption of alcoholic beverages increases the toxic effects of Thiram.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Workers and other handlers must wear:

- ◆ Long-sleeved shirt and long pants
- ◆ Waterproof gloves
- ◆ Shoes plus socks
- ◆ Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protection Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Engineering control statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

User should:

- ◆ Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- ◆ Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- ◆ Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

GENERAL PRODUCT INFORMATION

Thiram Granuflo® is a broad-spectrum fungicide for plant disease control in the culture of Apples, Peaches, and Strawberries. Thiram Granuflo® is a broad-spectrum fungicide for use on Non-commercial Turf (parks, golf courses, rights-of-way, commercial landscapes, etc.) to control Dollar Spot, Brown Patch and Fusarium Patch. When used as a Seed Protectant, Thiram Granuflo® will usually increase crop stands and yields by reducing seedling losses from seed decay, damping-off or seedling blights.

UCB's recommendations for the use of this product are based upon tests believed to be reliable. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CHEMIGATION: Do not apply this product through any type of irrigation system.

GENERAL RECOMMENDATIONS

Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use.

For agricultural foliar applications use recommended rates for dilute, concentrate and aerial applications. Ground application is preferred, especially when full canopy of foliage exists. For dilute application, use Thiram Granuflo® at the recommended rate listed in the "per 100 gallons" column of the Specific Applications table. For concentrate ground application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum of 20 gallons per acre. For aerial application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum of 10 gallons per acre.

Applications of Thiram Granuflo® fungicide are most effective when made before infection periods. Under low disease conditions, minimum label rates should be used; however, maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions. If you are not familiar with such conditions, get advice from your state agricultural extension service.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Addition of a spreader/sticker to the finished spray mixture may improve coverage. Make fresh spray mixtures for each application.

3 of 12

MIXING INSTRUCTIONS

To mix, fill the spray tank about 2/3 full of water, and with agitator running, slowly sift in the recommended amount of Thiram Granuflo®. Finish filling the tank with water and continue agitation until thoroughly mixed. Continue agitation and apply immediately.

AGRICULTURAL APPLICATIONS (FOLIAR):

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- ♦ Long-sleeved shirt and long pants
- ♦ Waterproof gloves
- ♦ Shoes plus socks
- ♦ Chemical-resistant headgear for overhead exposure

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

FRUIT CROPS	DISEASES	POUNDS THIRAM GRANUFLO®		DIRECTIONS FOR USE
		per acre	per 100 gal	
APPLES*	Apple Blotch (<i>Phylosticta solitaria</i>) Apple Scab (<i>Venturia inaequalis</i>) Bitter Rot (<i>Glomerella cingulata</i>) Black Pox (<i>Helminthosporium papulosum</i>) Black Rot on fruit and foliage (<i>Botryosphaeria obtusa</i>) Brook's Spot (<i>Mycosphaerella pomi</i>) Bull's Eye Rot (<i>Pezicula malicorticis</i>) (Pacific Northwest Only) Fly Speck (<i>Schizothyrium pomi</i>) Gray Mold Rot (<i>Botrytis cinerea</i>) Sooty Blotch (<i>Gloeodes pomigena</i>) White Rot (<i>Botryosphaeria dothidea</i>)	5.2 to 6.8 lbs	1.3 to 1.7 lbs ⁽¹⁾	Apply at pre-bloom, calyx, early cover sprays and late cover sprays.
	Cedar Apple Rust, Quince Rust (<i>Gymnosporangium</i> spp.)	5.2 to 6.8 lbs	1.3 to 1.7 lbs ⁽¹⁾	Apply at pre-bloom, calyx and first cover. Additionally, apply 2.6 lbs/A (0.65 lb/100 gal) at bloom.
PEACHES*	Brown Rot Blossom Blight and Fruit Rot (<i>Monilinia</i> spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	For Blossom Blight, apply at 3-4 day intervals during bloom to protect blossoms. For Fruit Rot, apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Leaf Curl (<i>Taphrina deformans</i>)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply dormant applications post-harvest in late fall and/or in late winter prior to bud swell, at bloom and petal-fall.
	Rhizopus Rot (<i>Rhizopus</i> spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Scab (<i>Cladosporium carpophilum</i>)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply at petal-fall and repeat 3 to 6 times at 10 to 14 day intervals as needed. Do not apply within 7 days of harvest.
STRAWBERRIES*	Gray Mold (<i>Botrytis cinerea</i>)	3.4 to 4.4 lbs	1.7 to 2.2 lbs ⁽³⁾	Broadcast 3 to 5 applications at 10 day intervals per crop cycle ⁽⁴⁾ beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results. Do not apply within 3 days of picking.

⁽¹⁾ Based on dilute sprays of 400 gallons per acre.

⁽²⁾ Based on dilute sprays of 300 gallons per acre.

⁽³⁾ Based on dilute sprays of 200 gallons per acre.

⁽⁴⁾ Crop cycle is defined as pre-bloom through harvest.

* Not for use in California.

DO NOT graze or feed cover crops from treated orchards.

DO NOT rotate treated crops with crops other than those with registered Thiram uses.

NON-COMMERCIAL TURF APPLICATIONS:

Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

Not for use on residential lawns or turf.

Thiram Granuflo® prevents infestation of common turf diseases such as Dollar Spot, Brown Patch and Fusarium Patch during the Summer growing season and Snow Mold during the cool weather seasons. When used at curative rates, Thiram Granuflo® will cure diseased turf infected with Dollar Spot, Brown Patch and Fusarium Patch.

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

TURF	DISEASES	OUNCES THIRAM GRANUFLO® per 1000 square feet	DIRECTIONS FOR USE
NON-COMMERCIAL TURF*	Dollar Spot (<i>Sclerotinia homeocarpa</i>) Brown Patch (<i>Rhizoctonia Blight</i>) (<i>Rhizoctonia solani</i>)	Preventative Rate: 2.5 to 3.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. beginning in early Spring <u>before</u> disease symptoms appear. Repeat at 7 to 10 day intervals as needed when environmental conditions favor disease development. Do not apply more than 40 oz per 1000 sq.ft. during the Summer growing season.
		Curative Rate: 4.5 to 5.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. after disease symptoms first appear. Repeat at 7 to 10 day intervals as needed. Do not apply more than 40 oz per 1000 sq.ft. during the Summer growing season.
	Fusarium Patch (<i>Fusarium nivale</i>) (Pink Snow Mold and Gerlachia Patch)	Preventative Rate: 8.0 oz	Apply recommended rate in 2 to 5 gallons water/1000 sq. ft. in late Fall immediately before expected permanent snow cover.
		Curative Rate: 3.0 oz	As a follow-up to the late Fall application, apply in Spring after snow melts and runoff has ended.

DO NOT graze or feed treated turf.

*Not for use in California

Seed Protectant Applications:

For control of damping-off, seed decay, seedling blights, covered kernel smut of sorghum and loose smut and bunt of wheat, etc., apply Thiram Granuflo® at recommended dosages as a dry treatment or as a water-based slurry. Apply with other registered seed treatment insecticides and fungicides through commercial slurry or mist-type seed treatment equipment.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels attached to treated seed must state "Treated seed must not be used for or mixed with food or animal feed, or processed for oil." and "Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

Thiram Granuflo® Seed Protectant Dosage Rates

TYPE OF SEED*	OUNCES THIRAM GRANUFLO® per 100 lbs
Barley	2.7
Beans (dry and succulent)	1.3
Beans (Lima)	2.2
Beans (snap)	1.3
Beets (garden)	5.3
Canola	4.3
Cantaloupe, Cucumber, Pumpkin, Squash, Watermelon	3.0
Castor beans	3.0
Chicory	3.5
Conifers	21.5
Corn (field and sweet)	2.0
Cottonseed (reginned, fuzzy and acid-delinted)	3.0
Cowpeas	1.3
Eggplant	4.0
Flax	3.5
Grasses	5.3
Legumes (small-seeded)	5.3
Lentils	2.0

50412

Millet	2.7
Oats	2.2
Okra	4.0
Onions	4.0
Peanuts (shelled and unshelled)	3.0
Peas	2.0
Rice	2.9
Rye	2.5
Safflower	2.7
Sesame	2.0
Sorghum	1.8
Soybeans	2.2
Sugar beets	5.3
Sunflowers	5.4
Triticale	2.2
Tomatoes	4.0
Wheat	2.2
Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Collards, Endive, Kale, Kohlrabi, Lettuce, Mustard, Peppers, Radish, Spinach, Swiss Chard, Turnip	5.3
Ornamental flower seeds, Ornamental foliage plants in the families Araliaceae (scheffleras), Araceae (philodendrons) and Palmaeae (palms)	5.4

*Not for use in California.

STORAGE AND DISPOSAL

Pesticide Storage

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for the use or storage in or around the home. Protect from excess heat.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TERMS OF SALE OR USE

On purchase of this product, buyer and user agree to the following conditions:

Warranty: UCB makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Granuflo® is a Registered Trademark of UCB S.A., a Belgian corporation.

For Emergencies Involving a Spill, Leak, Fire, Exposure, or Accident, contact: CHEMTREC at (800) 424-9300.
For Other Emergencies, phone (770) 801-5248 or (770) 433-6421.

NET WEIGHT:



Chemicals

UCB Chemicals Corporation
2000 Lake Park Drive
Smyrna, Georgia 30080

60712



Thiram Granuflo®

(The Original 75% Water Dispersible Granules)

Agricultural Fungicide

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 28 1997

ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram disulfide)..... 75.0%

INERT INGREDIENTS:..... 25.0%

TOTAL 100.0%

EPA Reg. No. 45728-21

EPA Est. 32557-BEL-1

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration, if indicated.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling spray mist. Do not use or store near food or feed.

The consumption of alcoholic beverages increases the toxic effects of Thiram.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- ♦ Long-sleeved shirt and long pants
- ♦ Waterproof gloves
- ♦ Shoes plus socks
- ♦ Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protection Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering control statements:
When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

User should:

- ♦ Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- ♦ Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- ♦ Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

GENERAL PRODUCT INFORMATION

Thiram Granuflo® is a broad-spectrum fungicide for plant disease control in the culture of Apples, Peaches, and Strawberries.

UCB's recommendations for the use of this product are based upon tests believed to be reliable. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CHEMIGATION: Do not apply this product through any type of irrigation system.

GENERAL RECOMMENDATIONS

Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use.

Use recommended rates for dilute, concentrate and aerial applications. Ground application is preferred, especially when full canopy of foliage exists. For dilute application, use Thiram Granuflo® at the recommended rate listed in the "per 100 gallons" column of the Specific Applications table. For concentrate ground application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum of 20 gallons per acre. For aerial application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum of 10 gallons per acre.

Applications of Thiram Granuflo® fungicide are most effective when made before infection periods. Under low disease conditions, minimum label rates should be used; however, maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions. If you are not familiar with such conditions, get advice from your state agricultural extension service.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Addition of a spreader/sticker to the finished spray mixture may improve coverage. Make fresh spray mixtures for each application.

7012

MIXING INSTRUCTIONS

To mix, fill the spray tank about 2/3 full of water, and with agitator running, slowly sift in the recommended amount of Thiram Granuflo®. Finish filling the tank with water and continue agitation until thoroughly mixed. Continue agitation and apply immediately.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- ♦ Long-sleeved shirt and long pants
- ♦ Waterproof gloves
- ♦ Shoes plus socks
- ♦ Chemical-resistant headgear for overhead exposure

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

FRUIT CROPS*	DISEASES	POUNDS THIRAM GRANUFLO®		DIRECTIONS FOR USE
		per acre	per 100 gal	
APPLES*	Apple Blotch (<i>Phylosticta solitaria</i>)	5.2 to 6.8 lbs	1.3 to 1.7 lbs ⁽¹⁾	Apply at pre-bloom, calyx, early cover sprays and late cover sprays.
	Apple Scab (<i>Venturia inaequalis</i>)			
	Bitter Rot (<i>Glomerella cingulata</i>)			
	Black Pox (<i>Helminthosporium papulosum</i>)			
	Black Rot on fruit and foliage (<i>Botryosphaeria obtusa</i>)			
	Brook's Spot (<i>Mycosphaerella pomi</i>)			
	Bull's Eye Rot (<i>Pezicula malicorticis</i>) (Pacific Northwest Only)			
	Fly Speck (<i>Schizothyrium pomi</i>)			
	Gray Mold Rot (<i>Botrytis cinerea</i>)			
	Sooty Blotch (<i>Gloeodes pomigena</i>)			
	White Rot (<i>Botryosphaeria dothidea</i>)			
	Cedar Apple Rust, Quince Rust (<i>Gymnosporangium</i> spp.)	5.2 to 6.8 lbs	1.3 to 1.7 lbs ⁽¹⁾	Apply at pre-bloom, calyx and first cover. Additionally, apply 2.6 lbs/A (0.85 lb/100 gal) at bloom.
PEACHES*	Brown Rot, Blossom Blight and Fruit Rot (<i>Monilinia</i> spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	For Blossom Blight, apply at 3-4 day intervals during bloom to protect blossoms. For Fruit Rot, apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Leaf Curl (<i>Taphrina deformans</i>)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply dormant applications post-harvest in late fall and/or in late winter prior to bud swell, at bloom and petal-fall.
	Rhizopus Rot (<i>Rhizopus</i> spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Scab (<i>Cladosporium carpophilum</i>)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply at petal-fall and repeat 3 to 6 times at 10 to 14 day intervals as needed. Do not apply within 7 days of harvest.
STRAWBERRIES*	Gray Mold (<i>Botrytis cinerea</i>)	3.4 to 4.4 lbs	1.7 to 2.2 lbs ⁽³⁾	Broadcast 3 to 5 applications at 10 day intervals per crop cycle ⁽⁴⁾ beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results. Do not apply within 3 days of picking.

⁽¹⁾ Based on dilute sprays of 400 gallons per acre.

⁽²⁾ Based on dilute sprays of 300 gallons per acre.

⁽³⁾ Based on dilute sprays of 200 gallons per acre.

⁽⁴⁾ Crop cycle is defined as pre-bloom through harvest.

*Not for use in California.

DO NOT graze or feed cover crops from treated orchards.

DO NOT rotate treated crops with crops other than those with registered Thiram uses.

STORAGE AND DISPOSAL**Pesticide Storage**

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for the use or storage in or around the home. Protect from excess heat.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TERMS OF SALE OR USE

On purchase of this product, buyer and user agree to the following conditions:

Warranty: UCB makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Granuflo® is a Registered Trademark of UCB S.A., a Belgian corporation.

For Emergencies involving a Spill, Leak, Fire, Exposure, or Accident, contact: CHEMTREC at (800) 424-9300.
For Other Emergencies, phone (770) 801-3248 or (770) 433-6421.

NET WEIGHT:



Chemicals

UCB Chemicals Corporation
2000 Lake Park Drive
Smyrna, Georgia 30080



Thiram Granuflo®

(The Original 75% Water Dispersible Granules)

Turf Fungicide

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 28 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
45728-21

ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram disulfide)..... 75.0%

INERT INGREDIENTS:..... 25.0%

TOTAL 100.0%

EPA Reg. No. 45728-21

EPA Est. 32557-BEL-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration, if indicated.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling spray mist. Do not use or store near food or feed.

The consumption of alcoholic beverages increases the toxic effects of Thiram.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

GENERAL PRODUCT INFORMATION

Thiram Granuflo® is a broad-spectrum fungicide for use on Non-commercial Turf (parks, golf courses, rights-of-way, commercial landscapes, etc.) to control Dollar Spot, Brown Patch and Fusarium Patch.

UCB's recommendations for the use of this product are based upon tests believed to be reliable. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CHEMIGATION: Do not apply this product through any type of irrigation system.

GENERAL RECOMMENDATIONS

Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use.

Applications of Thiram Granuflo® fungicide are most effective when made before infection periods. Under low disease conditions, minimum label rates should be used; however, maximum label rates and shortened spray schedules are recommended for severe or threatening conditions. If you are not familiar with such conditions, get advice from your state agricultural extension service.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Addition of a spreader/sticker to the finished spray mixture may improve coverage. Make fresh spray mixtures for each application.

MIXING INSTRUCTIONS

To mix, fill the spray tank about 2/3 full of water, and with agitator running, slowly add in the recommended amount of Thiram Granuflo®. Finish filling the tank with water and continue agitation until thoroughly mixed. Continue agitation and apply immediately.

Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes. Not for use on residential lawns or turf.

Thiram Granuflo® prevents infestation of common turf diseases such as Dollar Spot, Brown Patch and Fusarium Patch during the Summer growing season and Snow Mold during the cool weather seasons. When used at curative rates, Thiram Granuflo® will cure diseased turf infected with Dollar Spot, Brown Patch and Fusarium Patch.

100412

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

TURF	DISEASES	OUNCES THIRAM GRANUFLO® per 1000 square feet	DIRECTIONS FOR USE
NON-COMMERCIAL TURF*	Dollar Spot (<i>Sclerotinia homeocarpa</i>) Brown Patch (<i>Rhizoctonia Blight</i>) (<i>Rhizoctonia solani</i>)	Preventative Rate: 2.5 to 3.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. beginning in early Spring <u>before</u> disease symptoms appear. Repeat at 7 to 10 day intervals as needed when environmental conditions favor disease development. Do not apply more than 40 oz per 1000 sq.ft. during the Summer growing season.
		Curative Rate: 4.5 to 5.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. after disease symptoms first appear. Repeat at 7 to 10 day intervals as needed. Do not apply more than 40 oz per 1000 sq.ft. during the Summer growing season.
	Fusarium Patch (<i>Fusarium nivale</i>) (Pink Snow Mold and Gerlachia Patch)	Preventative Rate: 8.0 oz	Apply recommended rate in 2 to 5 gallons water/1000 sq. ft. in late Fall immediately before expected permanent snow cover.
		Curative Rate: 3.0 oz	As a follow-up to the late Fall application, apply in Spring after snow melts and runoff has ended.

DO NOT graze or feed treated turf.
*Not for use in California.

STORAGE AND DISPOSAL

Pesticide Storage

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for the use or storage in or around the home. Protect from excess heat.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TERMS OF SALE OR USE

On purchase of this product, buyer and user agree to the following conditions:

Warranty: UCB makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding purchase price paid and shall not include incidental or consequential damages.

Granuflo® is a Registered Trademark of UCB S.A., a Belgian corporation.

For Emergencies Involving a Spill, Leak, Fire, Exposure, or Accident, contact: CHEMTREC at (800) 424-9300.
For Other Emergencies, phone (770) 801-3248 or (770) 433-8421.

NET WEIGHT:



Chemicals

UCB Chemicals Corporation
2000 Lake Park Drive
Smyrna, Georgia 30080

11042



Thiram Granuflo®

(The Original 75% Water Dispersible Granules)

Seed Protectant Fungicide

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 28 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram disulfide)..... 75.0%

INERT INGREDIENTS:

25.0%

TOTAL 100.0%

EPA Est. 32557-BEL-1

EPA Reg. No. 45728-21

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration, if indicated.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes skin and eye irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling spray mist. Do not use or store near food or feed.

The consumption of alcoholic beverages increases the toxic effects of Thiram.

ENVIRONMENTAL HAZARDS

Product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

GENERAL PRODUCT INFORMATION

When used as a Seed Protectant, Thiram Granuflo® will usually increase crop stands and yields by reducing seedling losses from seed decay, damping-off and seedling blights.

UCB's recommendations for the use of this product are based upon tests believed to be reliable. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CHEMIGATION: Do not apply this product through any type of irrigation system.

GENERAL RECOMMENDATIONS

For control of damping-off, seed decay, seedling blights, covered kernel smut of sorghum and loose smut and bunt of wheat, etc., apply Thiram Granuflo® at recommended dosages as a dry treatment or as a water-based slurry. Apply with other registered seed treatment insecticides and fungicides through commercial slurry or mist-type seed treatment equipment.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels attached to treated seed must state "Treated seed must not be used for or mixed with food or animal feed, or processed for oil" and "Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

STORAGE AND DISPOSAL

Pesticide Storage

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for the use or storage in or around the home. Protect from excess heat.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

12/12

Thiram Granuflo® Seed Protectant Dosage Rates

TYPE OF SEED*	OUNCES THIRAM GRANUFLO® per 100 lbs
Barley	2.7
Beans (dry and succulent)	1.3
Beans (Lima)	2.2
Beans (snap)	1.3
Beets (garden)	5.3
Canola	4.3
Cantaloupe, Cucumber, Pumpkin, Squash, Watermelon	3.0
Castor beans	3.0
Chicory	3.5
Conifers	21.5
Corn (field and sweet)	2.0
Cottonseed (reginned, fuzzy and acid-defatted)	3.0
Cowpeas	1.3
Eggplant	4.0
Flax	3.5
Grasses	5.3
Legumes (small-seeded)	5.3
Lentils	2.0
Millet	2.7
Oats	2.2
Okra	4.0
Onions	4.0
Peanuts (shelled and unshelled)	3.0
Peas	2.0
Rice	2.9
Rye	2.5
Safflower	2.7
Sesame	2.0
Sorghum	1.8
Soybeans	2.2
Sugar beets	5.3
Sunflowers	5.4
Triticale	2.2
Tomatoes	4.0
Wheat	2.2
Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Collards, Endive, Kale, Kohlrabi, Lettuce, Mustard, Peppers, Radish, Spinach, Swiss Chard, Turnip	5.3
Ornamental flower seeds, Ornamental foliage plants in the families Araliaceae (scheffleras), Araceae (philodendrons) and Palmaceae (palms)	5.4

*Not for use in California.

TERMS OF SALE OR USE

On purchase of this product, buyer and user agree to the following conditions:

Warranty: UCB makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Granuflo® is a Registered Trademark of UCB S.A., a Belgian corporation.

For Emergencies Involving a Spill, Leak, Fire, Exposure, or Accident, contact: CHEMTREC at (800) 424-9300.
For Other Emergencies, phone (770) 801-3248 or (770) 433-6421.

NET WEIGHT:



Chemicals

UCB Chemicals Corporation
2000 Lake Park Drive
Smyrna, Georgia 30080