PLEASE NOTE

This image contains more than one label approved for this product on this date.

45% Please read instructions on	28-2/	,	2-27-200 Form App	Oroved. OMB No. 2	070-0 080	l
\$EPA	Un Environmental	ited States		Registra Amenda Other	ation	271115
	ρ	pplication for	Pesticide - Sec	tion i		
1. Company/Product Numb 45121	_		2. EPA Product Mar			posed Classification
	Granuflo		Pages K	22		None Restricted
1000 /	pplicant Unclude ZIP Cod marial 19 Corps La ber Vorle U , GA 3062	no Hon	· · · · · · · · · · · · · · · · · · ·			FIFRA Section 3(c)(3) nposition and labeling
Check if th	is is a new address		Product Name			
		Se	ction - II		N	OTIFICATION
Amendment - Explain Resubmission in rec	sponse to Agency letter o	lated	Agency let	ed labels in respond ter dated Application. plain below.	se to	C 2 7 2000
Explanation: Use addition of the control of the con	onal page(s) if necessary on for remo fungicide on California letter for cer	ring the Apples, for the Notation	restriction Scaches and tice 98-10 1 ta team t	for the s s traw b.	use o unies lotice	f Thiram in the
	: 	Se	ction - III			
1. Material This Product V						
Child-Resistant Packaging Yes* No * Certification must	Unit Packaging Yes No /o /6. 6 If "Yes" Unit Packaging wgt.	No. per If "Y	Yes No Yes No. per contain			bag andboard
be submitted	10 16. box	4	<u> </u>			
3. Location of Net Content Label	S Information Container	4. Size(s) Retail Con イメノの	• • • • • • • • • • • • • • • • • • • •	5. Location of Li On Lab	al	npanying product
6. Manner in Which Label	is Affixed to Product	Lithograph Paper glued Stenciled	Oth	or		
		Se	ction - IV			
1. Contact Point Comple	te items directly below fo	or identification of inc	dividual to be contacted	i, if necessary, to j		
Name Timothy	1 Dotson	Title	ego la long Ah	airs May.	1 .	tio. (Include Area Code)
/ I certify that the sta	tements I have made on any knowingly false or m	Certification this form and all atte	chments thereto are tr	ue, accurate and 6		é. Dete Application Received (Ştamped)
2. Signature	Alo ton		me la trong A Ho	urs Man	a i por	
4. Typed Name Time thy	1. Do tson	5. Date	12/5/00			••••



VIA CERTIFIED MAIL

December 5, 2000

Mr. Dan Kenny
Product Manager (22)
Fungicide-Herbicides Branch
Office of Pesticide Programs – H7504C
U.S. Environmental Protection Agency
401 M St., S.W.
Washington, D.C. 20460-0001

Subject:

Notification for Removing the Restriction for the Use of Thiram

Granuflo® Fungicide (EPA Reg. No. 45728-21) on Apples, Peaches and

Strawberries in the State of California Per PR-Notice 98-10

Dear Dan:

UCB Chemicals Corporation would like to submit for registration of Thiram Granuflo® Agricultural Fungicide, a sub-label of Thiram Granuflo® Fungicide, to the State of California. In order to do so, we must first remove the restriction for sales of this product in California. UCB Chemicals' supported a self-imposed restriction since the product is not currently registered in California. We are removing this restriction by removing the table end-note for apples, peaches and strawberries denoted by asterisks in the "Specific Applications for Thiram Granuflo®" table.

Please find enclosed 1) one copy of EPA Form 8570-1, Application for Pesticide – Notification, and, 2) one copy of the new label for Thiram Granuflo[®] Fungicide.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR-Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.





Page 2 December 05, 2000 Mr. Dan Kenny

If you have any questions, please call at 770/801-3212.

Sincerely,

Timothy A. Dotson

Regulatory Affairs Manager

U.S. and Canada

Enclosure

Thiram Granuflo®

(The Original 75% Water Dispersible Granules)
Fungicide

NOTIFICATION

ACTIVE INGREDIENT:

.0% DEC 27 **2000**

EPA Reg. No. 45728-21

TOTAL 100.0% EPA Est. No. 32557-BEL-1

KEEP OUT OF REACH OF CHILDREN CAUTION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration, if indicated

PRECAUTIONARY STATEMENTS -

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling spray mist. Do not use or store near food or feed. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

The consumption of alcoholic beverages increases the toxic effects of Thiram.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- *Long-sleeved shirt and long pants
- *Waterproof gloves
- *Shoes plus socks
- *Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protection Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering control statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

User should:

*Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

*Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

*Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

GENERAL PRODUCT INFORMATION

Thiram Granuflo® is a broad-spectrum fungicide for plant disease control in the culture of Apples, Peaches, and Strawberries. Thiram Granuflo® is a broad-spectrum fungicide for use on Turf (parks, golf courses, rights-of-way, commercial landscapes, etc.) to control Dollar Spot, Brown Patch and Fusarium Patch. When used as a Seed Protectant, Thiram Granuflo® will usually increase crop stands and yields by reducing seedling losses from seed decay, damping-off or seedling blights. When used as a Repellent, Thiram Granuflo® can protect trees, vines, nursery stock, shrubs and ornamentals from rabbit, deer and rodent depredation.

UCB's recommendations for the use of this product are based upon tests believed to be reliable. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal Law $\,$ to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CHEMIGATION: Do not apply this product through any type of irrigation system.

GENERAL RECOMMENDATIONS

Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use.

For agricultural foliar applications use recommended rates for dilute, concentrate and aerial applications. Ground application is preferred, especially when full canopy of foliage exists. For dilute application, use Thiram Granuflo® at the recommended rate listed in the "per 100 gallons" column of the Specific Applications table. For concentrate ground application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum of 20 gallons per acre. For aerial application, apply the recommended rate disted in the "per acre" column of the Specific Applications table in a minimum of 10 gallons per acre.

Applications of Thiram Granuflo® fungicide are most effective when made before infection periods. Under low disease conditions, minimum label rates should be used; however, maximum label rates and shortened spray schedules are recommended for severe or threatening disease

5/9

conditions. If you are not familiar with such conditions, get advice from your state agricultural extension service.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Addition of a spreader/sticker to the finished spray mixture may improve coverage. Make fresh spray mixtures for each application.

MIXING INSTRUCTIONS

To mix, fill the spray tank about 2/3 full of water, and with agitator running, slowly sift in the recommended amount of Thiram Granuflo®. Finish filling the tank with water and continue agitation until thoroughly mixed. Continue agitation and apply immediately.

AGRICULTURAL APPLICATIONS (FOLIAR):

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

**Long-sleeved shirt and long pants

**Waterproof gloves

**Shoes plus socks

į

**Chemical-resistant headgear for overhead exposure

Page 2 of 6

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

POUNDS THIRAM GRANUFLO®

FRUIT CROPS	DISEASES	per acre	per 100 gal	DIRECTIONS FOR USE
APPLES	Apple Blotch (Phyllosticta solitaria) Apple Scab (Venturia inaequalis) Bitter Rot (Glomerella cingulata) Black Pox (Helminthosporium papulosum) Black Rot on fruit and foliage (Botryosphaeria obtusa) Brook's Spot (Mycosphaerella pomi) Bull's Eye Rot (Pezicula malicorticis) (Pacific Northwest Only) Fly Speck (Schizothyrium pomi) Gray Mold Rot (Botrytis cinerea)	per acre 5.2 to 6.8 lbs	per 100 gal	Apply at pre-bloom, calyx, early cover sprays and late cover sprays.
	Sooty Blotch (Gloeodes pomigena) White Rot (Botryosphaeria dothidea) Cedar Apple Rust, Quince Rust	5.2 to 6.8 lbs	1.3 to 1.7 lbs ⁽¹⁾	Apply at pre-bloom, calyx and first cover. Additionally, apply
	(Gymnosporangium spp.)	V.2 (V C.O.)	110 10 117 100	2.6 lbs/A (0.65 lb/100 gal) at bloom.
PEACHES	Brown Rot Blossom Blight and Fruit Rot (Monilinia spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	For Biossom Blight, apply at 3-4 day intervals during bloom to protect biossoms. For Fruit Rot, apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Leaf Curl (Taphrina deformans)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply dormant applications post-harvest in late fall and/or in late winter prior to bud swell, at bloom and petal-fall.
	Rhizopus Rot (Rhizopus spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Scab (Cladosporium carpophilum)	3.9 to 5.1 lbs	1.3 to 1.7 lbs ⁽²⁾	Apply at petal-fall and repeat 3 to 6 times at 10 to 14 day intervals as needed. Do not apply within 7 days of harvest.
STRAWBERRIES	Gray Mold (Botrytis cinerea)	3.4 to 4.4 lbs	* 7 to 2.2 lbs**	Broadcast 3 to 5 applications at 10 day intervals per crop cycle fibeginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results. Do not apply within 3 days of picking.

DO NOT graze or feed cover crops from treated orchards.



Page 3 of 6

Thiram Granuflo 10/26/00.TAD(EPAxxx, CDPRxxx)

Based on dilute sprays of 400 gallons per acre.
 Based on dilute sprays of 300 gallons per acre.
 Based on dilute sprays of 200 gallons per acre.
 Crop cycle is defined as pre-bloom through harvest.

TURF APPLICATIONS:

Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

Not for use on residential lawns or turf.

Thiram Granuflo® prevents infestation of common turf diseases such as Dollar Spot, Brown Patch and Fusarium Patch during the Summer growing season and Snow Mold during the cool weather seasons. When used at curative rates, Thiram Granuflo® will cure diseased turf infected with Dollar Spot and Brown Patch and can also prevent Fusarium Patch when two applications are made during the Fall through the Spring season.

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

TURF	DISEASES	OUNCES THIRAM GRANUFLO® per 1000 square feet	DIRECTIONS FOR USE
TURF	Dollar Spot (Sclerotinia homeocarpa) Brown Patch (Rhizoctonia Blight) (Rhizoctonia solani)	Preventative Rate: 2.5 to 3.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. beginning in early Spring <u>before</u> disease symptoms appear. Repeat at 7 to 10 day intervals as needed when environmental conditions favor disease development. Do not apply more than 40 oz per 1000 sq.ft. during the Summer growing season.
		Curative Rate: 4.5 to 5.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. after disease symptoms first appear. Repeat at 7 to 10 day intervals as needed. Do not apply more than 40 oz per 1000 sq.ft. during the Summer growing season.
	Fusarium Patch (Fusarium nivale) (Pink Snow Mold and Gerlachia Patch)	Preventative Rate: 8.0 oz	Apply recommended rate in 2 to 5 gallons water/1000 sq. ft. in late Fall immediately before expected permanent snow cover. As a follow-up to the late Fall application, a Spring application of 3.0 oz/1000 sq. ft. during Spring thaw and after runoff has ended may be necessary to aid in the prevention of Spring disease.

DO NOT graze or feed treated turf.

ANIMAL REPELLENCY APPLICATIONS*:

When used according to label directions, Thiram Granuflo® can protect trees, vines, nursery stock, shrubs and ornamentals from rabbit, deer and rodent depredation. One application will protect treated parts from 1 to 3 months, depending upon weather conditions. Thiram Granuflo® can also be used as a repellent in detering deer from moving into important areas such as airport runways as a result of local feeding and browsing. It can also deter migratory birds such as geese on golf courses and other turf areas. The addition of a latex based sticker prolongs the residual effectiveness of the treatment. Repellency is the result of rendering the treated plant parts distasteful.

Applications To Vines, Tree Trunks and Dormant Plants:

Prepare a slurry of 1 pound of Thiram Granuflo® in 3 quarts of water and apply as a full coverage spray to all portions of the plant to be protected. For greater retention and residual efficacy, add 1 pint of a latex based sticker to the spray solution.

Brush On Application:

Prepare a slurry of 1 pound of Thiram Granuflo® in 3 quarts of water and apply with a paint-brush to stem parts and trunk parts to be protected. For greater retention and residual efficacy, add 1 fluid ounce of a latex based sticker into each quart volume of solution prepared.

Foliar Applications:

Prepare a slurry of 1 pound of Thiram Granuflo® in 6 quarts of water and apply as a full coverage spray to all portions of the plant to be protected. For greater retention and residual efficacy, add 1 pint of a latex based sticker to the spray solution. DO NOT use wetting agents or surfactants as foliar damage may result. For best results, spray when temperatures are greater than 40 F and relative humidity is less than 70 percent. DO NOT USE FRUIT FOR HUMAN CONSUMPTION.

* NOT FOR USE IN CALIFORNIA

Page 4 of 6

• 4 2 4

Thiram Granuflo 10/26/00.TAD(EPAxxx, CDPRxxx)

^{*}Not for use in California

SEED PROTECTANT APPLICATIONS:

For control of damping-off, seed decay, seedling blights, covered kernel smut of sorghum and loose smut and bunt of wheat, etc., apply Thiram Granuflo® at recommended dosages as a dry treatment or as a water-based slurry. Apply with other registered seed treatment insecticides and fungicides through commercial slurry or mist-type seed treatment equipment.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels attached to treated seed must state "Treated seed must not be used for or mixed with food or animal feed, or processed for oil." and "Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

Thiram Granuflo® Seed Protectant Dosage Rates

TYPE OF SEED*	OUNCES THIRAM GRANUFLO® per 100 lbs	
Barley	2.7	
Beans (dry and succulent)	1.3	
Beans (Lima)	2.2	
Beans (snap)	1.3	
Beets (garden)	5.3	
Canola	4.3	
Cantaloupe, Cucumber, Pumpkin, Squash, Watermelon	3.0	
Castor beans	3.0	
Chicory	3.5	
Conifers	21.5 ·	
Corn (field and sweet)	2.0	
Cottonseed (reginned, fuzzy and acid-delinted)	3.0	
Cowpeas	1.3	
Eggplant	4.0	
Flax	3,5	
Grasses	5.3	
Legumes (small-seeded)	5.3	
Lentils	2.0	
Millet	2.7	
Oats	2.2	
Okra	4.0	
Onions	4.0	
Peanuts (shelled and unshelled)	3.0	
Peas	2.0	
Rice	2.9	
Rye	2.5	
Safflower	2.7	
Sesame	2.0	
Sorghum	1.8	
Soybeans	2.2	
Sugar beets	5.3	
Sunflowers	5.4	
Triticale	2.2	
Tomatoes	4.0	
Wheat		
	2.2	
Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower,	5.3	
Collards, Endive, Kale, Kohlrabi, Lettuce, Mustard,	1	
Peppers, Radish, Spinach, Swiss Chard, Turnip		
Ornamental flower seeds, Ornamental foliage plants in the	5.4	
families Araliacea (scheffleras), Aracea (philodendrons) and		
Palmacea (palms)		

*Not for use in California.

Thiram Granuflo 10/26/00.TAD(EPAxxx, CDPRxxx)

Page 5 of 6

STORAGE AND DISPOSAL

Pesticide Storage

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for the use or storage in or around the home. Protect from excess heat.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TERMS OF SALE OR USE

On purchase of this product, buyer and user agree to the following conditions:

Warranty: UCB makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Granuflo® is a Registered Trademark of UCB S.A., a Belgian corporation.


For Emergencies Involving a Spill, Leak, Fire, Exposure, or Accident, contact: CHEMTREC at (800) 424-9300. For Other Emergencies, phone (770) 801-3248 or (770) 433-6421.

NET WEIGHT:



Chemicals

UCB Chemicals Corporation 2000 Lake Park Drive Smyrna, Georgia 30080



• Page 6 of 6

NEXT

LABEL

Please read instructions on reverse before completing form.	12-27-2000	2070 0000		
United States Environmental Protection Washington, DC 20460	Agency	opp Identifier Number ndment 271116		
Application	for Pesticide - Section I			
1. Company/Product Number 45728-21	2. EPA Product Manager **Ban *** **Leng *** **The Company of the	3. Proposed Classification		
4. Company/Product (Name) Thiram Grameflo Fungicile	PM# 22	None Restricted		
5. Name and Address of Applicant (Include ZIP Code), UCB Chemical's Corporation 2000 Loke Park Drive Smyraa, GA 3008	•	ordance with FIFRA Section 3(c)(3) dentical in composition and labeling		
Check if this is a new address	Product Name			
	Section - II	NOTIFICATION		
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.	Final printed labels in res Agency letter dated "Me Too" Application. Other - Explain below.	DEC 2.7 2000		
Explanation: Use additional page(s) if necessary. (For section is Le-submission of notification for non-agricultural use requirements be agricultural use requirements be alrelusion of certification state.	mand Section II.) "Segregating" or separate to under the wifs; or X mant per /K-Notice Section - III	Chibian of non-		
4. Massiel This Product MEN De Posture des	Section - III			
1. Material This Product Will Be Packaged In: Child-Resistant Packaging Yes* No * Certification must Unit Packaging No No No If "Yes" Unit Packaging wgt. container	Water Soluble Packaging 2. Ty Yes No If "Yes" No. per Package wgt container	pe of Container Metal Plastic Glass Paper Other (Specify)		
be submitted	1	Z Other (Specify)		
3. Location of Net Contents Information 4. Size(s) Retailed Label Container 47.10	On	of Label Directions Lebel Labeling accompanying product		
6, Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled Cother				
	Section - IV			
Contact Point (Complete items directly below for identification				
1	Regulatory Attails Mayo.	770-801 - 921 2-		
Certificat I certify that the statements I have made on this form and a I acknowledge that any knowingly false or misleading state both under applicable law.	all attachments thereto are true, accurate a			
	i. Title	, , , , , , , , , , , , , , , , , , , ,		
	Regalatory Attails M.	*		
14. Typed Name Timothy A. Dotson	12/14/00			



VIA CERTIFIED MAIL

December 14, 2000

Mr. Dan Kenny
Product Manager (22)
Document Processing Desk (NOTIF)
Fungicide-Herbicide Branch
Registration Division (7504C)
U.S. Environmental Protection Agency
401 M St., S.W.
Washington, D.C. 20460-0001

Subject:

Re-Submission of the Notification for "Segregating" Agricultural and Non-Agricultural Use Requirements Under the Worker Protection Standard; Addition of Non-Agricultural Use Requirements Box for Thiram Granuflo® Fungicide (EPA Reg. No. 45728-21)

Dear Dan:

The following notification "segregates" or separates the uses for Thiram Granuflo® Fungicide into "Agricultural Use Requirements" and "Non-Agricultural Use Requirements" based on the Worker Protection Standard, in order to distinguish between WPS and non-WPS PPE requirements. Based on the WPS, the non-agricultural uses are already "de-scoped" from the WPS, and thus this notification only provides additional clarification.

Please find enclosed 1) one copy of EPA Form 8570-1, Application for Pesticide – Notification, and 2) one copy of the new Thiram Granuflo® Fungicide label (EPA Reg. No. 45728-21).

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR-Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.





Page 2 Mr. Dan Kenny December 13, 2000

If you have any questions, please call at 770/801-3212.

KADo bon

Sincerely,

Timothy A. Dotson

Regulatory Affairs Manager

U.S. and Canada

Label Code: 12/14/00.TAD(EPA12/14/00)

(The Original 75% Water Dispersible Granules) **Fungicide**

ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram disulfide)......75.0% OTHER INGREDIENTS......25.0% NOTIFICATION

PAL 100.0% DEC 2 7 2000 EPA Est. No. 32557-BEL-1

EPA Reg. No. 45728-21

KEEP OUT OF REACH OF CHILDREN CAUTION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration, if indicated.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling spray mist. Do not use or store near food or feed. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

The consumption of alcoholic beverages increases the toxic effects of

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- *Long-sleeved shirt and long pants
- *Waterproof gloves
- *Shoes plus socks
- *Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining Personal Protection Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering control statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

User should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

GENERAL PRODUCT INFORMATION

Thiram Granuflo® is a broad-spectrum fungicide for plant disease control in the culture of Apples, Peaches, and Strawberries. Thiram Granuflo® is a broad-spectrum fungicide for use on Turf (parks, golf courses, rightsof-way, commercial landscapes, etc.) to control Dollar Spot, Brown Patch and Fusarium Patch. When used as a Seed Protectant, Thiram Granuflo® will usually increase crop stands and yields by reducing seedling losses from seed decay, damping-off or seedling blights. When used as a Repellent, Thiram Granuflo® can protect trees, vines, nursery stock, shrubs and ornamentals from rabbit, deer and rodent depredation.

UCB's recommendations for the use of this product are based upon tests believed to be reliable. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

Not for use on residential lawns or turf.

CHEMIGATION: Do not apply this product through any type of irrigation system.

GENERAL RECOMMENDATIONS

Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use.

For agricultural foliar applications use recommended rates for dilute, concentrate and aerial applications. Ground application, is preferred, especially when full canopy of foliage exists. For dilute application, use Thiram Granuflo® at the recommended rate listed in the "per 100 gallons" column of the Specific Applications table. For concentrate ground application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum of 20 gallons per actet. For aerial application, apply the recommended rate listed in the "per acre" . column of the Specific Applications table in a minimum of 10 gallons person acre.Applications of Thiram Granuflo® fungicide are most effective

when made before infection periods. Under low disease conditions, minimum label rates should be used; however, maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions. If you are not familiar with such conditions, get advice from your state agricultural extension service.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Addition of a spreader/sticker to the finished spray mixture may improve coverage. Make fresh spray mixtures for each application.

MIXING INSTRUCTIONS

To mix, fill the spray tank about 2/3 full of water, and with agitator running, slowly sift in the recommended amount of Thiram Granuflo®. Finish filling the tank with water and continue agitation until thoroughly mixed. Continue agitation and apply immediately.

AGRICULTURAL APPLICATIONS (FOLIAR):

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

**Long-sleeved shirt and long pants

**Waterproof gloves

**Shoes plus socks

**Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL APPLICATIONS:

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Non-Residential Turf Use: Keep children and pets out of treated areas until sprays have dried.

Page 2 of 6

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

POUNDS THIRA	M GRANUFLO _®	
per acre	per 100 gal	DIRECTIONS FOR USE
5.0 to 6.0 lb-	1 2 to 1 7 lbo(1)	Apply of ore bloom, asky, each, payor corrup and late

FRUIT CROPS	DISEASES	per acre	per 100 gal	DIRECTIONS FOR USE
APPLES*	Apple Blotch (Phyllosticta solitaria) Apple Scab (Venturia inaequalis) Bitter Rot (Glomerella cingulata) Black Pox (Helminthosporium papulosum) Black Rot on fruit and foliage (Botryosphaeria obtusa) Brook's Spot (Mycosphaerella pomi) Bull's Eye Rot (Pezicula malicorticis) (Pacific Northwest Only) Fly Speck (Schizothyrium pomi) Gray Mold Rot (Botrytis cinerea) Sooty Blotch (Gloeodes pomigena) White Rot (Botryosphaeria dothidea)	5.2 to 6.8 lbs	1.3 to 1.7 lbs(1)	Apply at pre-bloom, calyx, early cover sprays and late cover sprays.
	Cedar Apple Rust, Quince Rust (Gymnosporangium spp.)	5.2 to 6.8 lbs	1.3 to 1.7 lbs(1)	Apply at pre-bloom, calyx and first cover. Additionally, apply 2.6 lbs/A (0.65 lb/100 gal) at bloom.
PEACHES*	Brown Rot Blossom Blight and Fruit Rot (Monilinia spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs(2)	For Blossom Blight, apply at 3-4 day intervals during bloom to protect blossoms. For Fruit Rot, apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Leaf Curl (Taphrina deformans)	3.9 to 5.1 lbs	1.3 to 1.7 lbs(2)	Apply dormant applications post-harvest in late fall and/or in late winter prior to bud swell, at bloom and petal-fall.
	Rhizopus Rot (Rhizopus spp.)	3.9 to 5.1 lbs	1.3 to 1.7 lbs(2)	Apply at petal-fall, two weeks after petal-fall and in cover sprays at 7 day intervals if rot is apparent on green fruit. Do not apply within 7 days of harvest.
	Peach Scab (Cladosporium carpophilum)	3.9 to 5.1 lbs	1.3 to 1.7 lbs(2)	Apply at petal-fall and repeat 3 to 6 times at 10 to 14 day intervals as needed. Do not apply within 7 days of harvest.
STRAWBERRIES*	Gray Mold (Botrytis cinerea)	3.4 to 4.4 lbs	1.7 to 2.2 lbs(3)	Broadcast 3 to 5 applications at 10 day intervals per crop cycle (4) beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results. Do not apply within 3 days of picking.

⁽¹⁾ Based on dilute sprays of 400 gallons per acre.

DO NOT graze or feed cover crops from treated orchards.

⁽²⁾ Based on dilute sprays of 300 gallons per acre.(3) Based on dilute sprays of 200 gallons per acre.

⁽⁴⁾ Crop cycle is defined as pre-bloom through harvest.

^{*} Not for use in California.

TURF APPLICATIONS:

Thiram Granuflo® prevents infestation of common turf diseases such as Dollar Spot, Brown Patch and Fusarium Patch during the Summer growing season and Snow Mold during the cool weather seasons. When used at curative rates, Thiram Granuflo® will cure diseased turf infected with Dollar Spot and Brown Patch and can also prevent Fusarium Patch when two applications are made during the Fall through the Spring season.

SPECIFIC APPLICATIONS FOR THIRAM GRANUFLO®

TURF	DISEASES	OUNCES THIRAM GRANUFLO $_{f \oplus}$ per 1000 square feet	DIRECTIONS FOR USE
TURF	Dollar Spot (Sclerotinia homeocarpa) Brown Patch (Rhizoctonia Blight) (Rhizoctonia solani)	Preventative Rate: 2.5 to 3.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft. beginning in early Spring before disease symptoms appear. Repeat at 7 to 10 day intervals as needed when environmental conditions favor disease development. Do not apply more than 40 oz per 1000 sq.ft. during the Summer growing season.
		Curative Rate: 4.5 to 5.0 oz	Apply recommended rate in 5 to 10 gallons water/1000 sq. ft, after disease symptoms first appear. Repeat at 7 to 10 day intervals as needed. Do not apply more than 40 oz per 1000 sq.ft, during the Summer growing season.
	Fusarium Patch (Fusarium nivale) (Pink Snow Mold and Gerlachia Patch)	Preventative Rate: 8.0 oz	Apply recommended rate in 2 to 5 gallons water/1000 sq. ft. in late Fall immediately before expected permanent snow cover. As a follow-up to the late Fall application, a Spring application of 3.0 oz/1000 sq. ft. during Spring thaw and after runoff has ended may be necessary to aid in the prevention of Spring disease.

DO NOT graze or feed treated turf.

ANIMAL REPELLENCY APPLICATIONS*:

When used according to label directions, Thiram Granuflo® can protect trees, vines, nursery stock, shrubs and ornamentals from rabbit, deer and rodent depredation. One application will protect treated parts from 1 to 3 months, depending upon weather conditions. Thiram Granuflo® can also be used as a repellent in detering deer from moving into important areas such as airport runways as a result of local feeding and browsing. It can also deter migratory birds such as geese on golf courses and other turf areas. The addition of a latex based sticker prolongs the residual effectiveness of the treatment. Repellency is the result of rendering the treated plant parts distasteful.

Applications To Vines, Tree Trunks and Dormant Plants:

Prepare a slurry of 1 pound of Thiram Granuflo® in 3 quarts of water and apply as a full coverage spray to all portions of the plant to be protected. For greater retention and residual efficacy, add 1 pint of a latex based sticker to the spray solution.

Brush On Application:

Prepare a slurry of 1 pound of Thiram Granuflo® in 3 quarts of water and apply with a paint-brush to stem parts and trunk parts to be protected. For greater retention and residual efficacy, add 1 fluid ounce of a latex based sticker into each quart volume of solution prepared.

Foliar Applications:

Prepare a slurry of 1 pound of Thiram Granuflo® in 6 quarts of water and apply as a full coverage spray to all portions of the plant to be protected. For greater retention and residual efficacy, add 1 pint of a latex based sticker to the spray solution. DO NOT use wetting agents or surfactants as foliar damage may result. For best results, spray when temperatures are greater than 40 F and relative humidity is less than 70 percent. DO NOT USE FRUIT FOR HUMAN CONSUMPTION.

* NOT FOR USE IN CALIFORNIA

Page 4 of 6

^{*}Not for use in California

SEED PROTECTANT APPLICATIONS:

For control of damping-off, seed decay, seedling blights, covered kernel smut of sorghum and loose smut and bunt of wheat, etc., apply Thiram Granuflo® at recommended dosages as a dry treatment or as a water-based slurry. Apply with other registered seed treatment insecticides and fungicides through commercial slurry or mist-type seed treatment equipment.

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels attached to treated seed must state "Treated seed must not be used for or mixed with food or animal feed, or processed for oil." and "Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

Thiram Granuflo® Seed Protectant Dosage Rates

TYPE OF SEED*	OUNCES THIRAM GRANUFLO® per 100 lbs
Barley	2.7
Beans (dry and succulent)	1.3
Beans (Lima)	2.2
Beans (snap)	1.3
Beets (garden)	5.3
Canola	4.3
Cantaloupe, Cucumber, Pumpkin, Squash, Watermelon	3.0
Castor beans	3,0
Chicory	3.5
Conifers	21.5
Corn (field and sweet)	2.0
Cottonseed (reginned, fuzzy and acid-delinted)	3.0
Cowpeas	1.3
Eggplant	4.0
Flax	3.5
Grasses	5.3
Legumes (small-seeded)	5.3
Lentils	2.0
Millet	2.7
Oats	2.2
Okra	4.0
Onions	4.0
Peanuts (shelled and unshelled)	3.0
Peas	2.0
Rice	2.9
Rye	2.5
Safflower	2.7
Sesame	2.0
Sorghum	1.8
Soybeans	2.2
Sugar beets	5.3
Sunflowers	5.4
Triticale	2.2
Tomatoes	4.0
Wheat	2.2
Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower. Collards, Endive, Kale, Kohlrabi, Lettuce, Mustard, Peppers, Radish, Spinach, Swiss Chard, Turnip	5.3
Ornamental flower seeds, Ornamental foliage plants in the families Araliacea (scheffieras), Aracea (philodendrons) and Palmacea (palms)	5.4

*Not for use in California.



STORAGE AND DISPOSAL

Pesticide Storage

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for the use or storage in or around the home. Protect from excess heat.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary tandfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TERMS OF SALE OR USE

On purchase of this product, buyer and user agree to the following conditions:

Warranty: UCB makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Granuflo® is a Registered Trademark of UCB S.A., a Belgian corporation.

For Emergencies Involving a Spill, Leak, Fire, Exposure, or Accident, contact: CHEMTREC at (800) 424-9300. For Other Emergencies, phone (770) 801-3248 or (770) 433-6421.

NET WEIGHT:



Chemicals

UCB Chemicals Corporation 2000 Lake Park Drive Smyrna, Georgia 30080