

45735-25

10/30/2002

1/8

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number
--	---	---	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 45735-25	2. EPA Product Manager 	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Sevin 4L Flowable	PM#	
5. Name and Address of Applicant (Include ZIP Code) Burlington Scientific Corp. 71 Carolyn Blvd. Farmingdale, NY 11735 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: <div style="text-align: center;">NOTIFICATION</div> EPA Reg. No. _____ <div style="text-align: center;">OCT 30 2002</div> Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Change primary product name from "Sevin 4 Flowable" to "Carbaryl 4L Flowable"

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III


1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input checked="" type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	* Certification must be submitted	
If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> _____		
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Thomas C. McEntee	Title Regulatory Manager	Telephone No. (Include Area Code)
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Manager	
4. Typed Name Thomas C. McEntee	5. Date October 3, 2002	

NOTIFICATION
OCT 30 2002

CARBARYL 4L FLOWABLE

PULL UP 

**Controls Insects on
Vegetables, Lawns, Ornamentals and Fruits**

Active Ingredient:		
Carbaryl (1-Naphthyl N-methylcarbamate)		42.85%
Inert Ingredients		57.15%
TOTAL:		100.00%
This product contains 4 pounds Carbaryl per Gallon		

EPA REG. NO. 45735-25
EPA Est. NO. 74134-CHN-001, 16-VA-1₂
Subscript following Est. No. Corresponds to first digit in Lot No.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See booklet for statement of practical treatment and precautionary statements

Burlington Scientific Corp.
71 Carolyn Blvd.
Farmingdale, NY 11735

CARBARYL 4L

FLOWABLE

Controls Insects on
Vegetables, Lawns, Ornamentals and Fruits

Active Ingredient	
Carbaryl (1-Naphthyl N-methylcarbamate)	42.85%
Inert Ingredients	57.15%
TOTAL:	100.00%

This product contains 4 pounds Carbaryl per gallon

EPA REG. NO. 45735-25
EPA Est. NO. 74134-CHN-001, 16-VA-1₂
Subscript following Est. No. Corresponds to first digit in Lot No.

KEEP OUT OF REACH OF CHILDREN
CAUTION

10/3/02
① of 6 3/8

Statement of Practical Treatment.

This product is an N-methyl carbamate and is a cholinesterase inhibitor

If Swallowed: Drink one to two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If Inhaled: Remove victim to fresh air.

If On Skin: Wash with soap and water.

If In Eyes: Flush eyes with water.

Note to Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor.

Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

Burlington Scientific Corp.

71 Carolyn Blvd.

Farmingdale, NY 11735

Notice: Buyer assumes all responsibility for safety and use not in accordance with label directions.

For information on this product (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Tele-communication Network at 1-800/858-7378.

Precautionary Statements**Hazards to Humans and Domestic Animals**

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Do not take internally.

Personal Protective Equipment:**Applicators and Other Handlers Must Wear:**

- Long-Sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Chemical resistant head gear for overhead protection

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water.

Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

4/8

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: Coveralls, Waterproof gloves, and Shoes plus socks, and chemical resistant head gear for overhead exposure.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

Labeling for the Carbaryl 4L FLOWABLE

MIXING INSTRUCTIONS: To insure a uniform product, gently **SHAKE CONTENTS BEFORE POURING**. Be sure sprayer tank, spray tips and screens are clean before use. Fill the sprayer with $\frac{1}{2}$ to $\frac{3}{4}$ the desired amount or clean water. Pour the recommended amount of product into the tank and complete filling to desired level with water. Before and frequently during use, agitate the spray mixture in the tank to prevent settling. It is recommended that spray equipment be thoroughly cleaned after each day's use.

Vegetables

Carbaryl 4L Flowable will control a wide range of vegetable insect pests. Begin applications when insects first appear and repeat applications to maintain control but not more often than once every 7 days. Do not apply to any crop not listed on this label. Coverage of both upper and lower leaf surfaces is essential for good control. Use sufficient gallonage to obtain control.

Apply 1 quart per acre to control: Alfalfa caterpillar, bean leaf beetle, cucumber beetle, Colorado potato beetle, harlequin bug, Japanese beetle, leafhopper, meal worms, Mexican bean beetle, red-necked peanutworm, squash bug, three-cornered alfalfa hopper, thrips, velvetbean caterpillar and Western bean cutworm.

Apply 1 $\frac{1}{2}$ quarts per acre to control: armyworms, aster leafhopper, corn earworm, cutworms, grasshopper, pea leaf weevil, pea weevil, spittlebug and yellow striped armyworm.

Apply 2 quarts per acre to control: Asparagus beetle, chinch bug, corn rootworms, cowpea curculio, European corn borer, fall armyworm, imported cabbageworm, lace bug, lima beanpod borer, lygus bug, stink bug, sap beetle, Southwestern corn borer, tarnished plant bug, tomato fruitworm, tomato hornworm and tomato pinworm

Carbaryl 4L Flowable may be applied to the vegetables listed below. Follow the time intervals between last application and harvest indicated by the number in parenthesis "()" following the crop name. Asparagus (1), Beans, including cowpeas, crowder or Southern peas, dry beans, green beans, lima beans, navy beans, snap beans (succulent 3, dry and hay 21, forage 14), broccoli (3), Brussels sprouts (3), beets (root 7, tops 14), cabbage (3), cauliflower (3), Chinese cabbage (14), collards (14), carrots (7), corn (sweet 2; sweet, field and pop: forage & silage 14, grain and fodder 48), cucumbers (3) dandelion (14), endive (escarole) (14) eggplant (3), Hanover salad (14), horseradish (7), kale (14), kohlrabi (3), leaf and head lettuce (14), mustard greens (14), melons (3), okra (3), peanuts (14), peas (succulent 3, dried and hay 21, forage 14), peppers (3), potatoes (7), pumpkin (3), radish (7), rutabaga (7), squash (3), salisfy (roots 7, tops 14), spinach (14) and tomatoes (3).

FORAGE AND FIELD CROPS

Apply when insects or their damage appear. Repeat applications at 7 to 14 day intervals, if necessary, to maintain control. Thorough and uniform coverage is essential for effective control.

APPLY 1 QUART PER ACRE TO CONTROL: blister beetle, Mexican bean beetle, alfalfa caterpillar, bean leaf beetle, cucumber beetle, green cloverworm, Japanese beetle, leafhopper, three-cornered alfalfa hopper, thrips, velvetbean caterpillar, striped blister beetle, red-necked peanutworm, flea beetles, grape colaspis and cereal leaf beetle.

APPLY 1 $\frac{1}{2}$ QUART PER ACRE TO CONTROL: Armyworm, corn earworm, alfalfa worm, webworm, lygus bug, chinch bug, sorghum midge, Southwestern corn borer, beet weevil, Egyptian alfalfa weevil larvae, essex skipper.

APPLY 2 QUARTS PER ACRE TO CONTROL: whitefringed beetle adult, painted lady (thistle caterpillar), sweet potato weevil, leafhopper, stink bug, saltmarsh caterpillar, stem weevil, sweet potato hornworm, tarnished plant bug, white grub (green June Beetle), cutworm, yellowstriped armyworm, sunflower beetle, tortoise beetle, leafroller.

part
ALI
On
FOI
pre
PE/
RIC
PRI
RE:
SO
Trea
are
gall
SO
HE:
SU
SU
SW
WH
TO:
To c
inju
hav
hon
Use
allo
CH
Do
beg
AD:
To c
adj:
and
kill:
OR
Car
upp
nec
arm
woi
gall
fulk
bud
leaf

5/8

parenthesis "(" following each crop. Follow specific directions for crops where listed.

ALFALFA (7) AND CLOVERS (7): For alfalfa weevil larvae, if pre-treatment damage is extensive, cut alfalfa and treat the stubble. On dense growth, use 25 to 40 gallons of water per acre with ground equipment to insure adequate coverage.

FORAGE GRASSES (7) PASTURE (14) AND RANGELAND (14): To control thrips in grasses grown for seed, use high spray pressure to improve penetration into boot.

PEANUTS (14)

RICE (14): CAUTION! May kill shrimp and crabs. Do not use in areas where these are important resources. **DO NOT APPLY PROPANIL HERBICIDES WITHIN 15 DAYS BEFORE OR AFTER APPLICATION OF THIS PRODUCT OR PLANT INJURY WILL RESULT.** MISSISSIPPI DELTA AND TEXAS.

SORGHUMS (Milo and grain sorghum) (forage 14, grain & fodder 21): Direct spray into forming heads for optimum insect control. Treat for sorghum midge when 25 to 30 percent of the heads have emerged from the boot and are in bloom. Repeat 7 days later if adults are still active. A third application may be necessary in late planted sorghum or if midge are abundant. For chinch bugs, use high gallonage ground application directed at the base of plants.

SOYBEANS (Succulent 3, Dried and Hay 21, Forage 14), DO NOT APPLY A COMBINATION OF Carbaryl 4L AND 2,4-DB HERBICIDE TO SOYBEANS.

SUGAR BEETS (28 for roots and forage)

SUNFLOWER (60): Do not use in California

SWEET POTATO (7): Apply as a foliage spray. Full coverage of plant is essential. Do not use in California.

WHEAT (Forage 7, Grain & Straw 21)

TOBACCO

To control tobacco flea beetle in plant beds, apply 2 tablespoons/gallon or 1 quart/50 gallons per 100 square yards. To prevent plant injury, avoid excessive applications. To control green June beetles grubs, use 1 ½ quarts per 100 gallons and apply to areas that larvae have uprooted by sprinkling as a drench with 50 to 100 gallons per square yards. In fields to control budworms, fall armyworm, flea beetle, hornworm, Japanese beetle, June beetle and shuckfly, apply 1 to 2 quarts per acre. Use lower rate on young plants (up to knee high). Use at least 10 gallons of prepared spray per acre. Begin treatments when worms are small. If late season applications are necessary, allow three days before pruning or cutting.

CHESTNUTS (14)

Do not use in California. To control chestnut weevil, apply 2 or 3 quarts per 100 gallons of water. Make 4 applications at weekly intervals beginning in late July for adult chestnut weevil control. Last application should be made prior to shuck split.

ADULT MOSQUITO CONTROL

To control adult mosquito infestations in pastures, rangelands, yards, parks, recreational areas, logging camps, military posts and adjacent forested lands or wastelands, apply ½ to 1 quart per 100 gallons of water (1/2 to 1 tablespoons per gallon). Treat shrubbery and areas where adult mosquitoes congregate. Treat with pressure ground sprayers when adult mosquitoes are active. Caution! May kill shrimp and crabs. Do not use in area where there are important resources. **OBSERVE BEE CAUTION.**

ORNAMENTALS AND LAWNS

Carbaryl 4L Flowable will control the insects listed below. Apply 1 quart to 100 gallons of water and spray thoroughly covering both upper and lower leaf surfaces. Begin applications when insects, or their damage first appear and repeat in 7-10 day intervals if necessary, to maintain control. The addition of DRAGON® Spreader Sticker may improve residual control. Ants, apple aphid, armyworm, azalea, leafminer, bagworms, birch leaf miner, blister beetle, boxelder bug, boxwood leafminer, browntail moth, cankerworms, catalpa sphinx, chiggers, cooley spruce gall aphid, cutworms, cypress tip moth, Douglas Fir Tussock moth, Eastern Spruce gall aphid, Elm leaf aphid, Elm leaf beetle, Elm leaf spanworm, triophid mites, European Pine shoot moth, fall armyworm, flea beetle, fuller rose beetle, gall midges, gall wasps, green striped mapleworm, grasshoppers, gypsy moth, hackleberry hipplegall maker, molly bud moth, molly leafminer, jackpine budworm, budworm, Japanese beetle, Jeffrey pine needleminer, June beetles, lace bug, leafhoppers, leafrollers, locust borer, Maple leafcutter, mealy bugs, mimosa webworm, Nantucket Pine tip moth, Oak leafminers, Oak leaf skeletoniser,

4 6/8

oakworm complex, Oleander caterpillar, olive ash borer, orange stripe oakworm, orange tortrix, periodical cicada, pine sawfly, pine spittlebug, pitch pine tip moth, plant bug, insectia hornworm, flyids, puss caterpillar, lumpy oakworm, rose aphid, rose chafer, rose slug, saddled prominent, jarflies (exposed), scale insects, sowbugs, spiny elm caterpillar, springtails, spruce budworm, spruce needleminer, subtropical pine tip moth, tent caterpillars, thorn bugs, thrips (exposed), ticks, walnut caterpillars, webworms, Western Hemlock looper, Western Spruce budworm, willow leaf beetles, and yellow poplar weevil.

LAWN AND RECREATIONAL AREAS: Carbaryl 4L Flowable will control the following insects on lawns and recreational areas: ants, armyworms, bluegrass billbug, centipedes, chiggers, cutworms, earwigs, Essex skipper, European chafer, fall armyworm, fiery skipper, fleas, grasshoppers, June beetles, leafhoppers, Lucerne moth, millipedes, mosquitoes, sod webworms (lawn moths), sowbugs, springtails, ticks, yellowstriped armyworm, chinch bugs, and white grubs. Mix Carbaryl 4L Flowable with water according to directions given in the following table and apply as a coarse (large droplet) spray using a knapsack type, hose attachment applicator or other similar equipment. For best results, mow lawn and apply after watering or rain. If it is impractical to use the quantity of water listed in the table, apply in at least one gallon of water per 1,000sq. ft. and sprinkle with the recommended amount of water after application to insure good penetration of turf. Repeat 2-3 weeks later if necessary. Use the higher rate listed to control chinch bugs and white grubs. To treat 500 sq. ft., use 3 1/2 to 5 oz. Carbaryl 4L Flowable to 15 gallons of water, treat 1,000 sq. ft. with 7 1/2 to 10 oz. per 30 gallons of water, treat 2,500 sq. ft. with 1/2 - 2/3 quarts to 75 gallons of water, treat 5,000 sq. ft. with 1-1 1/2 quarts to 150 gallons of water.

TREE FRUIT, NUT CROPS, SMALL FRUIT

Carbaryl 4L Flowable will control many insect pests on fruit and nut trees. Use sufficient gallonage to obtain coverage. Follow the specific directions for each crop. The time interval between last application and harvest is indicated by the number in parenthesis "()" following each crop. Follow locally issued spray schedules for timing and frequency of applications in your area.

TREE FRUIT

APPLE Thinning: Apply 1/2 to 1 pint per 100 gallons of water in a full coverage dilute spray between 10 to 25 days after full bloom. If the factors such as tree age, variety, nutrition, previous crop, pruning, bloom and degree of set favor excessive fruit thinning with Carbaryl 4L Flowable, exercise caution to avoid possible yield reduction. The above rate is for easily thinned varieties including Cortland, Grimes, Jonathan, Macintosh, Orleans, Rome Beauty, Puritan, Red Delicious, Winesap, Yellow Newton, For more difficult to thin varieties including Baldwin, Ben Davis, Duchess, Early Macintosh, Golden Delicious, Lady Apple, Northern Spy Rhode Island Greening, Steele Red, Turley, Wealthy, Yellow Transparent and York Imperial, use 1 to 2 pints per 100 gallons of water.

APPLE, PEARS (3): Carbaryl 4L Flowable will control the insects on apples and pears, apple aphid, apple maggot, apple mealybug, apple rust mite, apple sucker, bagworm, California pear slug, (pear sawfly), codling moth, Eastern tent caterpillar, European apple sawfly, eyespotted, bud moth, forbes scale, fruittree leafroller, green fruitworm, Japanese beetle, lecanium scales, lesser appleworm, lygus bugs, orange tortrix, oystershell scale, pear curculio, red-banded leafroller, rosy apple aphid, San Jose scale, tarnished plant bug, tentiform leafminer, white apple leafhopper, wooly apple aphid, and yellow headed fire worm. Apply 1 quart per 100 gallons. Apple mealy bugs, apple aphid, codling moth, and white apple leafhopper may be controlled with 1 pint east of Rocky Mountains only. For psylla (coristel), apply when eggs hatch or young nymphs are present. For optimum scale control, apply when crawlers are present. To avoid undesirable apple thinning, delay use until at least 30 days after full bloom.

CITRUS FRUITS (5): Carbaryl 4L Flowable will control the following insects on citrus, avacado leafroller, black scale, brown soft scale, California orangedog, California Red Scale, Citricola scale, citrus cutworm, citrus rootweevil, citrus snow scale, fruittree leafroller, orange tortrix, western tussock moth, West Indian sugarcane borer (adult) and yellow scale. Apply 1 quart per 100 gallons. Do not apply less than 10 gallons of diluted spray mixtures per mature tree. May be mixed with petroleum oils commonly used on citrus.

PEACHES, PLUMS, PEARS, CHERRIES (3), APRICOTS, NECTARINES (3): Carbaryl 4L Flowable will control the following insects on the above fruits, apple pademas, black cherry aphids, brown soft scale, cherry fruitworm, cherry maggots, cucumber beetle, European earwig, eyespotted bud moth, forbes scale, fruittree leafhopper, green fruitworm, Japanese beetle, June beetle, Lecanium scales, lesser peachtree borer, mealy plum aphid, olive scale, oystershell scale, orange tortrix, Oriental fruit moth, peach twig borer,

per
mo
wh
NU
PE
eq
wel
API
Poc
pop
unil
the
To
Eur
dia
qua
SM
BR
spr
(7):
fri
anc
Jap
ske
(7):
ber
PO
CH
mit
of s
typ
on :
bui
De
sp
far
inc

5/8

periodical cicada, plum curculio, prune leafhopper, red-banded leafhopper, rose chafer, San Jose scale, tarnished plant bug, Tussock moth, variegated leafhopper. Apply 1 quart per 100 gallons of water. Spray for thorough coverage. For optimum scale control, apply when crawlers are present. For lesser peachtree borer control, spray limbs and tree trunk thoroughly weekly during moth flight.

NUT CROPS

PECANS (14): To control the following insects, apply 10 ounces per medium sized tree in approximately 20 gallons of water. This is equivalent of 3 1/8 quarts per acre based on 10 trees per acre. Black margined aphid, fall webworm, hickory shuckworm, lesser webworm, pecan leaf phylloxera, pecan nut-casebearer, pecan spittlebug, pecan weevil, twig girdler and walnut caterpillar. **SOIL APPLICATION FOR PECAN WEEVIL:** When foliar application of Carbaryl 4L Flowable is not feasible due to lack of proper equipment. Poorly accessible tree location or other factors, a soil application method may be used to provide suppression of adult pecan weevil population. Apply 5 ounces Carbaryl 4L Flowable per tree, to the soil under tree in sufficient water (10 gallons minimum per tree) to uniformly cover the entire area beneath the tree canopy 10 feet beyond the dripline. Make applications every 7-10 days from the time that the pecan shells begin to harden (early August) until the adult weevils are not longer present (about September 15). **WALNUTS (14):** To control the following insects, apply 1 pint per 100 gallons of water and spray the entire tree thoroughly, Calico scale, codling moth, European fruit lecanium, fiberworm, fruit leafroller, and frosted scale. For codling moth, apply first spray when average cross sectional diameters of developing nuts are 1/3 to 1/2 inch. Repeat during middle or late June as needed. To control European earwig, apply 2 quarts per 100 gallons of water and spray tree to point of runoff.

SMALL FRUIT CROPS

BRAMBLES (Blackberries, raspberries, dewberries, boysenberries, loganberries) (7): Apply 1-2 quarts per 100 gallons water and spray thoroughly to control: European raspberry aphid, Japanese beetle, leafhopper, rose chafer, snow tree cricket. **BLUEBERRIES (7):** Apply 1 1/2 quarts per 100 gallons water three weeks before harvest and repeat as necessary to control: blueberry maggot, cherry fruitworm, cranberry fruitworm, European fruit lecanium and Japanese beetle. **GRAPES (7):** Apply 2 quarts per 100 gallons of water and spray thoroughly to control: cutworms, eight spotted forester, European fruit lecanium, grape berry moth, grape leafhopper, Japanese beetle, June beetle, orange tortrix, omnivorous leafroller, red-banded leafroller, saltmarsh caterpillar, western grapeleaf skeletonizer, and western yellowstriped armyworm. Apply before first broad leafroller larvae emerge from rolls. **STRAWBERRIES (7):** Apply 1-2 quarts per 100 gallons to control: Flea beetles, meadow spittlebug, omnivorous leaflier (strawberry fruitworm), strawberry leafroller, and strawberry weevil. Carbaryl may injure Early Dawn and Sunrise varieties on Delmarva Peninsula.

POULTRY INSECT CONTROL

CHICKENS, DUCKS, GEESE, GAME BIRDS, PIGEONS AND TURKEYS: To control chicken mites, fleas, lice and northern fowl mites, use as direct mist spray on birds by: (1) misting with electric fog machine. Mix 10 ounces in 1 gallon of water. Use 1 1/2 gallons of spray mixture per 1,000 hens in cages, on litter or on slotted floor. Repeat in 4 weeks if necessary, or (2) spray with knapsack or cylinder type compressed air sprayers. Mix 6 ounces in 5 gallons of water. Apply 1 gallon of spray mixture per 100 hens in cages, on litter or on slotted floor. Repeat in 4 weeks if necessary. Direct mist spraying for chicken mites and fleas is a supplement to spraying roofs and buildings for control of these pests. Do not apply to poultry and game birds within 7 days of slaughter.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of on site or at an approved water disposal facility. Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose on in sanitary landfill or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Acti
Car
Iner
TO
This

6
8/8