

NET CONTENTS- 1 U.S. GALLON  
(128 FL.OZ.)

LIQUID  
SOIL FUNGICIDE

# ISOBAC 20

U.S. PATENT 3,346,447

ACTIVE INGREDIENTS: Mono-Sodium Salt Of 2,2-Methylenebis  
(3,4,6-Trichlorophenol) 20.0 %  
INERT INGREDIENTS 80.0 %

**WARNING** KEEP OUT OF REACH  
OF CHILDREN


See Back Panel for Additional Cautions

ALCO LABORATORIES, INC.

2000 W. 10TH AVE.  
MILWAUKEE, WIS. 53224

E.I. EST NO. 34881-TN-1

E.P.A. REG. NO. 15382-6



ACCEPTED

15 APR 1975

UNDER THE FEDERAL INSECTICIDE  
ACT AND THE FEDERAL FUNGICIDE ACT  
FOR THE YEAR 1974  
45989-4

RECEIVED  
MAR 26 1975

### DIRECTIONS FOR USE:

For control of damping-off, sore shin and soil-borne diseases attacking cotton seedlings.

- (1) In areas where cotton is planted deep (1 inch to 2½ inches), apply Isobac 20 at rates of 3 ozs. to 6 ozs. in 5-15 gals. water per acre in seed furrow at planting time.
- (2) In areas where cotton is planted shallow (½ to 1 inch), apply Isobac 20 at rates of 3 ozs. to 6 ozs. per acre in a 10-15 inch band over seeded row at planting time in 10-15 gallons of water per acre. Use at above rates with banded applications of herbicides used at planting time.
- (3) In all areas make additional applications if conditions warrant as plants emerge or after emergence at rates of 2 ozs. to 6 ozs. per acre in 2-10 gals. water per acre. Use 2 oz. rate in narrow banded sprays. Use 6 oz. rate for broadcast applications. May be applied in combination with early insecticide or post-emergence herbicide applications.

Isobac has been designed for use and is compatible with most herbicides and insecticides commonly used on cotton.

For control of boll-rot of cotton

Begin control program by using Isobac 20 in lay-by herbicide application at rate of 8 fluid ounces per acre to reduce inoculum potential.

Continue program by spraying Isobac when first bolls are formed. Use 4 fluid ounces Isobac per acre in aircraft or ground application systems. Continue sprays on 5-7 day intervals or oftener (3-4 day intervals) in periods of high humidity or in rainy periods. Isobac may be applied in combination with insecticides in low volume or ultra low volume aircraft systems; in conventional ground equipment use 15-25 gallons water per acre and direct spray toward bolls. Do not use within seven days of harvest.

#### WARNING

May be fatal if swallowed. Do not get in eyes or on skin. In case of contact wash thoroughly with water. For eyes get medical attention. Do not contaminate feed or foodstuffs. Do not breathe dust or spray mist.

Keep out of any body of water. Apply this product only as specified on this label.

#### WARRANTY

Seller makes no warranty, expressed or implied, including the warranties of merchantability and/or fitness for particular purpose, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material, when such use and/or handling is contrary to label instructions.



KALO LABORATORIES, INC.

2210 N. GREEN HILL ROAD, KANSAS CITY, MISSOURI 64117