

46-173-7

PM 25

BEST AVAILABLE COPY

1439-221

CONTROL No. 1

NET CONTENTS

GALLONS

DIRECTIONS FOR USE CHEM RICE is a selective herbicide for use in rice fields. It is effective against a wide range of grasses and weeds. It is applied as a spray or as a granular material. It is recommended that local authorities following through testing.

TIMING AND DOSAGE CHEM RICE is applied to the rice field at the time of planting. It is applied at a rate of 4 to 6 lbs. per acre. It is applied in a single application. It is applied in a single application. It is applied in a single application.

TO AVOID RESIDUES AT HARVEST DO NOT APPLY AFTER THE SEEDLING STAGE OF RICE. DO NOT EXCEED A TOTAL OF 4 POUNDS ACTIVE PER ACRE PER SEASON.

METHOD OF SEEDING CHEM RICE may be used in any method of seeding. It is applied to the seedbed or to the rice field. It is applied to the seedbed or to the rice field. It is applied to the seedbed or to the rice field.

WATER MANAGEMENT BEFORE TREATMENT Dry Planting. Plant in a moist seedbed or in a field that is well watered. Muddy Water Planting. The planting flood is usually drained 1 to 3 days after seeding. Clear Water Planting. Planting flood may be deep or shallow. It is lowered as the rice is pegged. Maintain a shallow flood or intermittent flooding until the rice is well rooted.

ALL METHODS OF PLANTING FLUSH THE FIELDS AS OFTEN AS NEEDED TO PREVENT DRYING AND CRUETING AND TO AID THE EMERGENCE OF RICE AND GRASS DURING PERIODS OF HIGH TEMPERATURE DRYING WINDS AND LOW RAINFALL. IT IS ESPECIALLY IMPORTANT TO FLUSH OFTEN ENOUGH TO KEEP THE GRASS GROWING ACTIVELY AND THUS MORE SUSCEPTIBLE TO THE OPTIMUM DOSAGE OF CHEM RICE. ALL OR MOST OF THE WATER SHOULD BE OFF THE FIELD AT THE TIME OF SPRAYING TO ALLOW FULL COVERAGE OF THE GRASSES AND WEEDS.

WATER MANAGEMENT AFTER TREATMENT TREATED FIELDS SHOULD ALWAYS BE FLOODED BEFORE A SECOND INFESTATION OF GRASS HAS A CHANCE TO DEVELOP BEYOND THE ONE LEAF STAGE. Start flooding in 12 to 24 hours in warm sunny weather. If treatment has been made on active growing grass in moist soil. During cool, cloudy weather or if treatment is made on slow growing grass in dry soil delay flooding until 2 to 5 days after treatment. It is recommended that the grass be completely submerged and maintain the flood at least 2 inches deep. It is recommended that the rice be submerged temporarily. At the rice is submerged temporarily. At the rice is submerged temporarily.

FERTILIZER Fields to be treated with CHEM RICE may be fertilized before or at planting time. Place nitrogenous fertilizer 2 to 4 inches deep to minimize loss during periods when the fields are drained.

EPA Reg No 11449-2

CHEM-SECT BRAND

CHEM RICE

POST EMERGENCE GRASS AND WEED KILLER

Active Ingredient 3,4-dichloropropionanilide 35% Inert Ingredient 65% 100%

Label is 4 lbs. of the active ingredient per gallon

WARNING KEEP OUT OF REACH OF CHILDREN

Avoid breathing spray mist. Harmful if swallowed, absorbed through the skin. Do not get in eyes or on skin or in clothing. In case of contact, flush skin and eyes with plenty of water, for eyes get medical attention. Do not take internally. May cause skin irritation. Do not store near food or feed. Do not contaminate streams, lakes or ponds with this material. Do not store at temperatures below 15°F.

CONSULT STATE AGRICULTURAL EXPERIMENTAL STATION OR EXTENSION SERVICE WEED SPECIALISTS FOR FURTHER RECOMMENDATIONS ON DOSAGE, TIMING, WATER MANAGEMENT AND CULTURAL PRACTICES TO MEET LOCAL CONDITIONS.

This product is not for use in streams, lakes or ponds with this material.

PATENTS PENDING

APPLICATION EQUIPMENT All CHEM RICE is applied to the ground and is applied to the ground. It is applied to the ground. It is applied to the ground. It is applied to the ground.

AIRCRAFT CHEM RICE is applied to the ground and is applied to the ground. It is applied to the ground. It is applied to the ground. It is applied to the ground.

GROUND SPRAYERS Standard low pressure herbicide sprayers equipped with nozzles are satisfactory. The sprayer which delivers a medium fine drop spray per acre at 40 to 50 psi at 100 feet ground speed is not in excess of 100 gals. per acre. In most rice fields, special care is needed to avoid skip to avoid leaving the boom too high so the spray fails to reach the weeds. OPERATING CONDITIONS Do not apply with winds or ground equipment with winds over 10 mph. A cross wind during application of 10 mph or more may be treated when grass is either dry or if rain threatening in 4 hours.

DRIFT HAZARD TO OTHER CROPS CHEM RICE causes most crops except grasses. Avoid drift or accidental applications from burning aircraft. Avoid drift or accidental applications from burning aircraft. Avoid drift or accidental applications from burning aircraft. Avoid drift or accidental applications from burning aircraft.

LIMITATIONS AND TOLERANCES FOR Crop Rice 55 days or before tillering Recommended use does not result in residues at

BEST AVAILABLE COPY

ACCEPTED Jul 14 1972

1439-229

BLUE SPRUCE COMPANY

519 SOUTH MAPLE AVENUE • BASKING RIDGE NEW JERSEY, U.S.A.

CABLE ADDRESS TREE