

Faint, illegible text at the top left of the page, possibly bleed-through from the reverse side.



SANI-PLEX

Disinfectant-Cleaner-Sanitizer-Fungicide-Midlevelstate-Virucide* —
Deodorizer for Hospitals, Institutional and Industrial Use
Effective in hard water up to 400 ppm hardness (calculated as CaCO₃) in the presence of 5% serum contamination

Precautionary Statements

Hazards to Humans and domestic animals

DANGER

Keep out of reach of children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed or absorbed through the skin. Avoid contamination of food.

STORAGE AND DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL

PESTICIDE, SPRAY MIXTURE OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS. OR BURY IN A SAFE PLACE

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES

NET CONTENTS

Active Ingredients

Octyl Decyl Dimethyl Ammonium Chloride	6.510%
Diethyl Dimethyl Ammonium Chloride	3.255%
Didecyl Dimethyl Ammonium Chloride	3.255%
(10% G + 50% C + 40% C + 10%) Dimethyl Benzyl Ammonium Chloride	8.680%
Inert ingredients	78.300%
	100.000%

ACCEPTED
46269-4
SEP 22 1982
U.S. Environmental Protection Agency
Fungicide and Herbicide Division
as an active ingredient for the pesticide
registered under
EPA Reg. No.

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
EPA Registration No. 46269-4 EPA Establishment No. 39183-DE-01

MANUFACTURED BY: Quip, Inc., 407 E. Ayre St., Newport, DE 19804
FOR SALE FOR USE AND STORAGE BY MAINTENANCE PERSONNEL ONLY