



SANI-PLEX

Disinfectant-Cleaner-Sanitizer-Fungicide-Mildewstate-Virucide*
Deodorizer for Hospitals, Institutional and Industrial Use
Effective in hard water up to 400 ppm hardness (calculated as CaCO₃) in the presence of 5% serum contamination

Active Ingredients

Octyl Decyl Dimethyl Ammonium Chloride	6.510%
Decyl Dimethyl Ammonium Chloride	3.255
Didecyl Dimethyl Ammonium Chloride	3.255
(Alkyl C-14, 50%, C-12, 40%, C-10, 10%) Dimethyl Benzyl Ammonium Chloride	8.680%
Inert Ingredients	78.300%
	100.000%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
EPA Registration No. 46269-4 EPA Establishment No. 39183-DE-01

MANUFACTURED BY: Quip, Inc., 407 E. Ayre St., Newport, DE 19804
FOR SALE FOR USE AND STORAGE BY MAINTENANCE PERSONNEL ONLY

Directions for Use GENERAL CLASSIFICATION

SANI-PLEX is a disinfectant-cleaner-sanitizer-fungicide-mildewstate-virucide which is effective in hard water up to 400 ppm hardness in the presence of 5% serum contamination.

Apply SANI-PLEX to walls, floors and other hard non-porous surfaces such as tables, chairs, counters, etc. by spraying and scrubbing with a cloth, mop or mechanical spray device so as to thoroughly wet the surface. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution of SANI-PLEX when the solution becomes visibly dirty.

Disinfection: To disinfect hard non-porous surfaces, add 0.5 oz. per gallon of water. Treated surfaces must remain wet for 10 minutes. At this use level, SANI-PLEX is also virucidal and fungicidal and fungistat.

Sanitizing: To sanitize porous or non-porous food contact surfaces, add 0.5 oz. per 5.5 gallons of water. Treated surfaces must remain wet for 60 seconds.

0.5 oz. per gallon use-level: The broad spectrum activity of SANI-PLEX has been evaluated and found to be effective against the following organisms by the AOAC Use Dilution test:

- Flavobacterium aeruginosa
- Staphylococcus aureus
- Salmonella choleraesuis
- Escherichia coli
- Streptococcus pyogenes
- Klebsiella pneumoniae
- Enterobacter aerogenes
- Salmonella schottmuelleri
- Streptococcus faecalis
- Shigella dysenteriae
- Brevibacterium ammoniagenes
- Salmonella typhi

AOAC Fungicidal Test

SANI-PLEX is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities, etc.

Mold and Mildew Control: At 0.5 oz. per gallon, SANI-PLEX will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard non-porous surfaces (as indicated in general instructions above). Allow to dry on surface and repeat when mildew growth returns.

Virucidal Performance: AT 0.5 oz. per gallon use-level, SANI-PLEX was evaluated and found to be effective against the influenza A, Brazil Virus on inanimate environmental surfaces.

Sanitizing-Non-Food Contact Surfaces (such as floors, walls, tables, etc.) At 0.5 oz. per 5.5 gallon use-level, SANI-PLEX is an effective sanitizer against Staphylococcus aureus and Klebsiella pneumoniae on hard porous and non-porous environmental surfaces.