
SUPER K-GRO

DURSBAN

LAWN INSECT SPRAY



as anisoide that the p register to en EPA Rug No 4451

SHATTEPPROOF

CONTROLS: Ants, Fleas, Grasshopper, Red Fire Ants, and Grubs in Lawns.

ACTIVE INGREDIENTS:

*Chlorpyrifos [0,0-diethy] 0-(3,5,6-trichloro-2- pyridyl) phosphorothioate]
**INERT INGREDIENTS
TOTAL
*Contains .40 lb. of Chlorpyrifos per gal. *Dursban ^R - Trademark of the Dow Chemical Company U.S. Patent No. 3,244,586

**Contains xylene range aromatic solvent.


Keep out of reach of children

CAUTION

See back panel for additional precautionary statements. Read entire label.

NET CONTENTS 1 PINT

(16 FL. OZ.) .473 LITER



IMPORTANT: TO AVOID DAMAGE OR POOR RESULTS, CAREFULLY READ ALL DIRECTIONS AND PRECAJTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes, and clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use in food areas of food handling establishments, restaurants or other places where food is commercially prepared or processed. Do not allow spray to come in contact with food, feed or water supplies. Food utensils should not be used after use with pesticides. Keep children and pets off treated areas until spray has dried.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one or two glasses of water. Do not induce vomiting. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF IN EYES: Flush eyes with water continuously for 15 minutes and get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with plenty of pater. Call a physician if irritation develops.

IF INHALED: Remove victim to fresh air. If appropriate, give artificial respiration. Get medical attention.

NOTE TO A PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is a antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, ponds or streams. Do not apply where run(ff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or to residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

WARNING

Do not use or store near heat or open flame.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Super K-Gro Dursban^R Lawn Insect Spray is a phosphate insecticide which provides control of insects attacking home lawns and other ornamental and recreational turf grass areas.

HOW TO USE: Shake well before using. Read entire label. Apply this product only as specified on this label. Do not spray vegetable gardens or other plants used for food purposes.

NOTE: 2 Tbsp. = 1 fl. oz. = 6 tsp. = 1/8 cup

ĺ

INDOOR PESTS: (Home invading pests inside dwellings). For spot treatment of Ants, Cockroaches, Clover Mites, Crickets, Firebrats, Silverfish and Spiders: mix 6 1/2 Tbsp. (3 1/3 oz.) in 1/2 gal. of water to reduce light infestations, and 13 1/3 Tbsp. (6 2/3 oz.) to reduce heavy infestations. Apply as a coarse spray or with a paint brush in corners, cracks in walls, along baseboards, beneath sinks, stoves and around plumbing.

For Ants: Apply to ant trails, around doors and windows.

*NOTE: When using this product on carpets or fine woods, test a small area for discoloration.

OUTDOOR PESTS: For centrol of Ants, Brown Dog Ticks, Chiggers, Chinch Bugs, Cockroaches, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Flies, Grasshoppers, Sod Webworms, Millipedes, Mosquitoes, Sowbugs and Spiders: mix 2/3 Tbsp. (1/3 oz.) in 1 gal. of water. Apply as a spray or with a paint brush to outside surfaces of buildings, porches, screens, window frames, eaves, patios and garages. Repeat as needed.

Outdoor Control Of Fleas And Ticks; In yards: Mix 3 Tbsp. (1 1/2 fl. oz.) of this product in 6 gals. of water to cover 200 sq. ft. Apply as a coarse, low pressure spray, and repeat as necessary. Thoroughly water immediately after treatment to wash insecticide into the treated area. For best results, the lawn should be moist at time of treatment.

In Kennels And Dog Houses: Mix 1 1/2 cups (12 2/3 oz.) in 1 gallon of water and apply to resting areas, walls and floors. Old bedding of pets should be removed and replaced with dian fresh bedding after treatment. Remove pets before spraying. Do not treat pets with this product. Keep children and pets off treated areas until spray has dried. Around Buildings: To help prevent infestation of building by Ants, Cockroaches, Crickets, Earwigs, Millipedes and Sowbugs: mix 2/3 Tbsp. in 1 1/2 gals. of water to cover 125 sq. ft. Apply thoroughly as a coarse spray to treat a band of soil 6-10 ft. wide around and adjacent to the building and 2-3 ft. high on the foundation. Pests Of Lawns. For control of Ants, Armyworms, Brown Dog Ticks, Chiggers, ChinchBugs, CloverMitestMsleBolek@tscKetsketsickets, Cutworms, Dichondra Flea Beetla, Earwigs, Fleas, Grasshoppers,

Sod Webworms: mix 1 3/4 Tbsp. (1 1/4 oz.) in 3 3/4 gals. of water to cover 125 sq. ft. Spray when pests appear, repeat as necessary. Apply as coarse spray using a hose-end sprayer. Thoroughly water immediately after treatment except as noted. For best results, lawn should be moist at time of treatment. One application usually will control Chinch Bugs for up to 8 weeks. For Sod Webworms, delay watering or mowing the treated area for 12-24 hours after treatment.

FOR TURFGRASS: For control of (Hyperodes) Weevil: mix 3 2/3 Tbsp. (1 3/402.) in 3 3/4 gals. of water. Spray in mid April and again in mid-May and follow recommended directions. For control of White Grubs (European Chafer and Japanese Beetle larvae only): mix 1 3/4 - 3 2/3 Tbsp. (1 - 1 3/4 oz.) in 3 3/4 gals of water. Spray when grubs are young and actively feeding near the soil surface usually late July and August. Immediately after spraying, irrigate the treated area with 1/2-2 inches of water. Consult your local Agricultural Extension Service Specialist for proper time to treat in your area for Hyperodes Weevil and White Grubs.

NOTE: Damaged lawn and ornamental turf should be fertilized to stimulate re-growth and recovery from insect injury.

FOR FIRE ANTS: Mix 4 3/4 Tbsp. (2 1/3 oz.) per gal. of water. Apply 1 gal. of solution as a gentle rain to each Fire Ant mound using a sprinkler can. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12.5 sg. ft.). For best results, apply in cocl weather at 65-80°F or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity and treat all colonies which may not as yet have constructed a mound. Keep children and pets off treated areas until spray has dried. Avoid tracking pesticides into the home via contaminated footwear.

ORNAMENTALS: (Flowers, shrubs, vines, shade and flowering trees and evergreens) - For control of Aphids, Bagworms, Eastern Tent Caterpillars, Fall Webworms, Grasshoppers, Orangestripped Oakworm, Yellow necked Caterpillar, Mites, Spittlebugs, and Whiteflies, mix 7 1/4 Tbsp.* (3 2/3 fl. oz.) in 4 1/2 gals. of water. Spray upper and lower leaf surfaces and infested limbs and trunk. Avoid spraying to the point of excessive runoff. Treat when pests appear and repeat at 7-10 day intervals if

.....

needed. NOTE: DO NOT USE ON AZALEAS, CAMELLIAS, POINSETTIAS, ROSE BUSHES OR VARIEGATED IVY BECAUSE OF POSSIBLE INJURY. For control of Cutworms, Leafhoppers, Mealybugs, Mimosa Webworms, Redhumped Caterpillar and Thrips (exposed), mix 7 1/4 Tbsp.** (3 2/3 oz.) in 2 1/2 gals. of water. For control of Ash and Lilac Borers; Oak, Pine Needle, Tea and White Peach Scale Crawlers, mix** (1 3/4 oz.) in 1 1/2 gals. of water.

(

*Equivalent to 2 oz. of chlorpyrifos per 50 gals. of water **Equivalent to 1 oz. of chlorpyrifos per 12 1/2 gals of water.

STORAGE & DISPOSAL

STORAGE: Store in original container. Store in cool, dry place preferably in locked storage area inaccessible to children. Store away from heat or open flame.

DISPOSAL: Do not reuse empty containers. Securely wrap original container in several layers of newspaper and discard in trash.

MANUFACTURED for Kmart Corporation, Troy, MI 48084

REGISTERED AND MANUFACTURED BY Celex Corporation, Plymouth, MI

48170 EPA REG. NO. 46515-2 EPA EST. NO. 46515-MI-1