

46515-8 PM-17 12/26/89

SUPER K-GRO

TOMATO & VEGETABLE INSECT SPRAY





3

KILLS: APHIDS, CUCUMBER BEETLES, MEXICAN BEAN BEETLES, TOMATO FRUITWORMS AND OTHER LISTED INSECTS.

**ACTIVE INGREDIENTS:** 0.02% Pyrethrins..... \*Piperonyl Butoxide, Technical..... 0.20% \*Equivalent to 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds. Keep Out of Reach of Children CAUTION See back panel for additional precautionary statements Read entire label. ..... NET CONTENTS 1 PT. 8 FL. OZ. Not Reviewed. Registrant claims to be in accordance with Amended 12-1-89

Not Reviewed. Registrant claims to be in accordance with Heregistration of Petticide Product Guidance for Phase 2 Response, Page 2.12, Active to inert Change in Status.

## IMPORTANT: TO AVOID DAMAGE OR POOR RESULTS, CAREFULLY READ ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or ab orbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

### STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Wash affected areas with soap and water. Get medical attention if irritation develops.

NOTE TO PHYSICIAN: Emergency information - call 1-800/858-7378

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO USE: Shake well before using. Adjust nozzle to a fine spray. Hold sprayer 12-15 inches from plant and spray, thoroughly covering both upper and lower leaf surfaces. Repeat weekly throughout the growing season when insects are present. Reapply after rain. For best results, spray early in the morning when air is still and temperatures are cool. Regular light sprayings at 2-3 day intervals are generally more effective than single heavy treatments.

USE ON THE FOLLOWING VEGETABLES: Asparagus, Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collard, Corp. Cranberries, Eggplant, Kale, Lettuce, Mustard Greens, Peppers, Potatoes, Radishes, Spinach, Tomatoes and Turnips.

KILLS THE FOLLOWING INSECTS: Aphids, Cabbage Looper, Cabbageworms, Colorado Potato Beetles, Corn Earworms, Cucumbet Beetles, Diamond Moth Larvae, Flea Beetles, Harlequin Buds, Imported Cabbageworms, Japanese Beetles, Leafhoppers, Mexican Bean Beetles, Potato Beetles, Tomato Fruitworms, Stink Bugs, Webworms, Whiteflies and Wireworms.

NOTE: For Corn Earworms, set nozzle to stream and spray inte ... corn silk. Make 3 to 4 applications at 2-3 day intervals.

# STORAGE & DISPOSAL

Store in original container, preferably in a locked storage area inaccessible to children.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

NOTE: Buyer assumes all responsibility for safety and use not in accordance with directions.

MANUFACTURED IN THE U.S.A. FOR KMART CORPORATION, TROY, MICHIGAN 48084

REGISTERED AND MANUFACTURED BY CELEX CORPORATION, PLYMOUTH, MICHIGAN 48170

EPA Est. No. 46515-MI-1

EPA Reg. No. 46515-8

E St

