

PA 17 46515-8 19 144
MAR 19 1991

Celex Corporation
c/o Ms. Joyce B. Pazianos
Pazianos Associates
1338 G Street SE.
Washington, DC 20003

Dear Ms. Pazianos:

Subject: Label Amendments
Super K-Gro Tomato & Vegetable Insect Spray
EPA Registration No. 46515-8
Your Submission Dated March 7, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Richard F. Mountfort
Product Manager (10)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure


BEST AVAILABLE COPY

62474; I; A-4; Mountfort; L10-3; KEV RIC; 03/14/91; PERM; CL; WO; DD; CL
CONCURRENCES

SYMBOL	SURNAME	DATE						

SUPER
K-GRO

TOMATO & VEGETABLE
INSECT SPRAY



KILLS: APHIDS, CUCUMBER BEETLES, MEXICAN BEAN BEETLES,
TOMATO FRUITWORMS AND OTHER LISTED INSECTS.

PROVIDES RAPID KILL OF INSECTS!

ACTIVE INGREDIENTS:

Pyrethrins.....	0.02%
*Piperonyl Butoxide, Technical.....	0.20%
INERT INGREDIENTS:	99.78%

*Equivalent to 0.16% (butylcarbityl
(6-propylpiperonyl) ether and 0.04%
compounds.

ACCEPTED
MAR 29 1991
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 46515-8

Keep Out of Reach of Children

CAUTION

See back panel for additional precautionary statements.
Read entire label.

NET CONTENTS 1 PT. 8 FL. OZ.

2-27-91

BEST AVAILABLE COPY

SUPER K-GRO TOMATO & VEGETABLE INSECT SPRAY

IMPORTANT: TO AVOID DAMAGE OR POOR RESULTS, CAREFULLY READ ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call physician immediately.
IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.
IF ON SKIN: Wash affected areas with soap and water. Get medical attention if irritation develops.

NOTE TO PHYSICIAN: Emergency information - call 1-800/858-7378

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO USE: Shake well before using. Adjust nozzle to a fine spray. Hold sprayer 12-15 inches from plant and spray, thoroughly covering both upper and lower leaf surfaces. Repeat weekly throughout the growing season when insects are present. Apply when insects or their damage first appear. Reapply after rain. For best results, spray early in the morning when air is still and temperatures are cool. Regular light sprayings at 2-3 day intervals are generally more effective than single heavy treatments.

USE ON THE FOLLOWING VEGETABLES: Asparagus, Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collard, Corn, Cranberries, Eggplant, Kale, Lettuce, Mustard Greens, Peppers, Potatoes, Radishes, Spinach, Tomatoes and Turnips.

KILLS THE FOLLOWING INSECTS: Aphids, Cabbage Looper, Cabbageworms, Caterpillars, Colorado Potato Beetles, Corn Earworms, Cucumber Beetles, Diamond Moth Larvae, Flea Beetles, Harlequin Bugs, Imported Cabbageworms, Japanese Beetles, Leafhoppers, Mexican Bean Beetles, Potato Beetles, Tomato Fruitworms, Tomato Hornworm, Stink Bugs, Webworms, Whiteflies, and Wireworms.

BEST AVAILABLE COPY

NOTE: For Corn Earworms, set nozzle to stream and spray into corn silk. Make 3 to 4 applications at 2-3 day intervals.

STORAGE & DISPOSAL

Store in original container, preferably in a locked storage area inaccessible to children.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.


NOTE: Buyer assumes all responsibility for safety and use not in accordance with directions.

MADE IN THE U.S.A. FOR KMART CORPORATION, TROY, MICHIGAN 48084

REGISTERED AND MANUFACTURED BY CELEX CORPORATION,
PLYMOUTH, MICHIGAN 48170

EPA Est. No. 46515-MI-1

EPA Reg. No. 46515-8



MAR 19 1991

Celex Corporation
c/o Ms. Joyce B. Pazianos
Pazianos Associates
1338 G Street SE.
Washington, DC 20003

Dear Ms. Pazianos:

Subject: Label Amendments
Super K-Gro Tomato & Vegetable Insect Spray
EPA Registration No. 46515-8
Your Submission Dated March 7, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

BEST AVAILABLE COPY

RFM
Richard F. Mountfort
Product Manager (10)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

62474; I; A-4; Mountfort; L10-3; KEV/IC; 03/14/91; PERM; CL; WO; DD; CL
CONCURRENCES

SYMBOL							
NAME							

320-1 (12-70)

OFFICIAL FILE COPY

JUN 17 1991

Ms. Joyce Pazianos
Pazianos Associates
1338 G Street SE.
Washington, DC 20003

Dear Ms. Pazianos:

Subject: Alternate Formulas: Cellex Corporation
Super K-Gro Tomato & Vegetable Insect Spray
EPA Registration No. 46515-8
Super K-Gro Whitefly & Mealybug Insect Killer
EPA Registration No. 46515-23
Super K-Gro House Plant Insect Killer
EPA Registration No. 46515-27
Your Submissions Dated January 5 and 10, 1990

The formula statements referred to above are acceptable as alternate formulas under the respective registrations and have been included in our files.

Sincerely yours,

BEST AVAILABLE COPY

RFM
Richard F. Mountfort
Product Manager (10)
Insecticide-Rodenticide Branch
Registration Division (H7505C)


63317:I:Mountfort:L10-7:KEVRIC:6/12/91:7/10/91:EK:wo:ek

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

SUPER
K-GRO

TOMATO & VEGETABLE
INSECT SPRAY



KILLS: APHIDS, CUCUMBER BEETLES, MEXICAN BEAN BEETLES,
TOMATO FRUITWORMS AND OTHER LISTED INSECTS.

PROVIDES RAPID KILL OF INSECTS!

ACTIVE INGREDIENTS:

Pyrethrins..... 0.02%

*Piperonyl Butoxide, Technical..... 0.08%

INERT INGREDIENTS:

*Equivalent to 0.16% (butylcarbitol
(6-propylpiperonyl) ether and 0.08%
compounds.


ACCEPTED
MAR 29 1991
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 46515-8

Keep Out of Reach of Children

CAUTION

See back panel for additional precautionary statements.
Read entire label.

NET CONTENTS 1 PT. 8 FL. OZ.



SUPER K-GRO TOMATO & VEGETABLE INSECT SPRAY

IMPORTANT: TO AVOID DAMAGE OR POOR RESULTS, CAREFULLY READ ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call physician immediately.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash affected areas with soap and water. Get medical attention if irritation develops.

NOTE TO PHYSICIAN: Emergency information - call 1-800/858-7378

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO USE: Shake well before using. Adjust nozzle to a fine spray. Hold sprayer 12-15 inches from plant and spray, thoroughly covering both upper and lower leaf surfaces. Repeat weekly throughout the growing season when insects are present. Apply when insects or their damage first appear. Reapply after rain. For best results, spray early in the morning when air is still and temperatures are cool. Regular light sprayings at 2-3 day intervals are generally more effective than single heavy treatments.

USE ON THE FOLLOWING VEGETABLES: Asparagus, Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collard, Corn, Cranberries, Eggplant, Kale, Lettuce, Mustard Greens, Peppers, Potatoes, Radishes, Spinach, Tomatoes and Turnips.

KILLS THE FOLLOWING INSECTS: Aphids, Cabbage Looper, Cabbageworms, Caterpillars, Colorado Potato Beetles, Corn Earworms, Cucumber Beetles, Diamond Moth Larvae, Flea Beetles, Harlequin Bugs, Imported Cabbageworms, Japanese Beetles, Leafhoppers, Mexican Bean Beetles, Potato Beetles, Tomato Fruitworms, Tomato Hornworm, Stink Bugs, Webworms, Whiteflies and Wireworms.

BEST AVAILABLE COPY

NOTE: For Corn Earworms, set nozzle to stream and spray into corn silk. Make 3 to 4 applications at 2-3 day intervals.

STORAGE & DISPOSAL

Store in original container, preferably in a locked storage area inaccessible to children.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.


NOTE: Buyer assumes all responsibility for safety and use not in accordance with directions.

MADE IN THE U.S.A. FOR KMART CORPORATION, TROY, MICHIGAN 48084

REGISTERED AND MANUFACTURED BY CELEX CORPORATION,
PLYMOUTH, MICHIGAN 48170

EPA Est. No. 46515-MI-1

EPA Reg. No. 46515-8



AUG 29 1991

Joyce B. Pazianos
Celex Corporation
377 Amelia St.
Plymouth, MI 48170

Dear Ms. Pazianos:

Subject: Alternate Formulations
Super K-Gro Tomato & Vegetable Insect Spray
EPA Reg. No. 46515-8
Super K-Gro Rose & Floral Insect Killer
EPA Reg. No. 46515-9
Super K-Gro Whitefly & Mealybug Insect Killer
EPA Reg. No. 46515-23
Super K-Gro House Plant Insect Killer
EPA Reg. No. 46515-27
Submissions: August 13, 1991

The formula statements referred to above are conditionally acceptable as alternate formulas under the respective registrations and have been included in our files. In order to support these alternate formulations, you must complete an acceptable eye irritation study on one of these formulations (representative of all four) and submit to EPA for review. In addition, submit label changes for all four products if the results of the study warrant.


Sincerely,

Richard F. Mountfort
Product Manager 10
Insecticide & Rodenticide Branch
Registration Division (H7505C)

RSB

SUPER
K-GRO

TOMATO & VEGETABLE
INSECT SPRAY



KILLS: APHIDS, CUCUMBER BEETLES, MEXICAN BEAN BEETLES,
TOMATO FRUITWORMS AND OTHER LISTED INSECTS.

PROVIDES RAPID KILL OF INSECTS!

ACTIVE INGREDIENTS:

Pyrethrins..... 0.02%

*Piperonyl Butoxide, Technical..... 0.20%

INERT INGREDIENTS: 99.78%

*Equivalent to 0.16% (butylcarbityl
(6-propylpiperonyl) ether and 0
compounds.

ACCEPTED
MAR 19 1991
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 46515-8

Keep Out of Reach of Children

CAUTION

See back panel for additional precautionary statements.
Read entire label.

NET CONTENTS 1 PT. 8 FL. OZ.

2-27-91

BEST AVAILABLE COPY

SUPER K-GRO TOMATO & VEGETABLE INSECT SPRAY

IMPORTANT: TO AVOID DAMAGE OR POOR RESULTS, CAREFULLY READ ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call physician immediately.
IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.
IF ON SKIN: Wash affected areas with soap and water. Get medical attention if irritation develops.

NOTE TO PHYSICIAN: Emergency information - call 1-800/858-7378

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO USE: Shake well before using. Adjust nozzle to a fine spray. Hold sprayer 12-15 inches from plant and spray, thoroughly covering both upper and lower leaf surfaces. Repeat weekly throughout the growing season when insects are present. Apply when insects or their damage first appear. Reapply after rain. For best results, spray early in the morning when air is still and temperatures are cool. Regular light sprayings at 2-3 day intervals are generally more effective than single heavy treatments.

USE ON THE FOLLOWING VEGETABLES: Asparagus, Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collard, Corn, Cranberries, Eggplant, Kale, Lettuce, Mustard Greens, Peppers, Potatoes, Radishes, Spinach, Tomatoes and Turnips.

KILLS THE FOLLOWING INSECTS: Aphids, Cabbage Looper, Cabbageworms, Caterpillars, Colorado Potato Beetles, Corn Earworms, Cucumber Beetles, Diamond Moth Larvae, Flea Beetles, Harlequin Bugs, Imported Cabbageworms, Japanese Beetles, Leafhoppers, Mexican Bean Beetles, Potato Beetles, Tomato Fruitworms, Tomato Hornworm, Stink Bugs, Webworms, Whiteflies and Wireworms.

BEST AVAILABLE COPY

NOTE: For Corn Earworms, set nozzle to stream and spray into corn silk. Make 3 to 4 applications at 2-3 day intervals.

STORAGE & DISPOSAL

Store in original container, preferably in a locked storage area inaccessible to children.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

) NOTE: Buyer assumes all responsibility for safety and use not in accordance with directions.

MADE IN THE U.S.A. FOR KMART CORPORATION, TROY, MICHIGAN 48084

REGISTERED AND MANUFACTURED BY CELEX CORPORATION,
PLYMOUTH, MICHIGAN 48170

EPA Est. No. 46515-MI-1

EPA Reg. No. 46515-8

