

PM 23 46575-16  
Pg. 10/3

**SUPER K-GRO**  
**FENCE & GRASS**  
**EDGER**  
**FORMULA II**

**READY  
TO USE**

**ACCEPTED**  
**JUN 10 1986**  
U.S. Environmental Protection Agency  
Federal Insecticide, Fungicide, and Rodenticide Act  
as amended  
the pesticide  
reg. no. 46515-16  
EPA

**NO MORE EDGING, CLIPPING OR CHIPPING GRASS AND WEEDS.**

**KEEPS GRASS FROM CREEPING INTO UNWANTED AREAS.**

**CONVENIENT APPLICATOR TOP**

**TOP KILLS:  
GRASSES AND WEEDS ON CONTACT. USE AROUND SHRUBS,  
TREES, DRIVEWAYS, SIDEWALKS, PATIOS AND FENCES.**

**WILL NOT STERILIZE SOIL.**

<b>ACTIVE INGREDIENTS:</b>	<b>By Lb.</b>
Diquat dibromide (6,7-dihydrodipyrido (1,2-1:2',1'-c)pyrazinedium dibromide) .....	0.23%
<b>INERT INGREDIENTS</b> .....	<u>99.77%</u>
<b>TOTAL</b> .....	<b>100.00%</b>

**Contains 0.01 lbs. Diquat Cation per gal.  
0.02 lbs. Salt per gal.**

**KEEP OUT OF REACH OF CHILDREN**

**WARNING**

**See back panel for additional precautionary statements. Read entire label.**

**NET CONTENTS 1 GALLON LIQUID**

pg. 2 of 3

**IMPORTANT:** To avoid damage or poor results, carefully read all directions and precautionary statements.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING:** Keep out of reach of children. Do not get on skin. May be fatal if swallowed, inhaled or absorbed through skin. May cause eye and skin irritation. Do not breathe vapors or spray mist. Do not allow spray to contaminate food, feed or water supplies. Keep children and domestic animals off treated areas until spray has thoroughly dried.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of milk, egg whites or gelatin solution. If these are unavailable, drink 1 or 2 glasses of water.

**IF IN EYES:** Flush eyes with water for 15 minutes and get medical attention.

**IF ON SKIN:** Remove contaminated clothing and wash affected area with plenty of water. Call a physician if irritation develops.

*correct comment in 5/16/66 letter*

**IF INHALED:** Remove victim to fresh air. If appropriate, give artificial respiration. Get medical attention.

**NOTE TO PHYSICIAN:** Emergency information - call 1-800/858-7378.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Do not apply when weather conditions favor drift from treated areas.

P. 101

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SUPER K-GRO FENCE & GRASS EDGER, FORMULA II** is a Ready-To-Use non-selective herbicide that top kills weeds and grasses on contact. This product becomes quickly inactivated on contact with soil and will not sterilize the soil when used as directed and it will not stain paved or graveled areas. Apply as directed to control unwanted vegetation around shrubs, trees, buildings, patios, in brick and gravel walks, and next to fences, driveways, sidewalks and curbs.

**HOW TO USE:** Shake well before using. Remove cap from applicator top and be certain that applicator top is firmly on jug. Invert jug and adjust stream of product by changing the angle and height at which the bottle is held. For edging around fences, lawns and sidewalks and top killing vegetation around shrubs, trees and buildings, hold the container 2 to 6 inches from the area to be treated depending upon the desired width of the spray pattern. For controlling weeds in unconfined areas such as gravel walks and driveways or patios, apply 1 gallon to 200 square feet. Holding the jug about waist high and away from the body to avoid contact with skin and clothing, walk slowly making sure that all undesirable vegetation is thoroughly covered. For spot treating, handle the jug so that the spray hits only the area to be treated.

**SUPER K-GRO FENCE & GRASS EDGER, FORMULA II** may be applied anytime during the year when weeds are actively growing. For best results, apply during warm, sunny weather. Wilting will occur in 1 to 3 days and complete browning of vegetation in 5 to 7 days. Some hard to control weeds may require more than one application. ✓

**IMPORTANT:** Apply when air is calm to avoid spray contacting desirable vegetation, green bark, broken bark, buds or leaves of desirable plants as damage may result. Do not track spray from treated areas to untreated areas. Do not apply product around plants to be used for food or feed.

#### STORAGE & DISPOSAL

**STORAGE:** Protect from freezing. Store in original container in locked storage area inaccessible to children. To prevent cross-contamination, do not store near pesticides, fertilizers, seeds, food or feed.

**DISPOSAL:** Do not reuse empty container. Securely wrap original container in several layers of newspaper and discard in trash.

**NOTE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

MANUFACTURED IN U.S.A. by Starb Corporation, Troy, MI 48064

REGISTERED AND MANUFACTURED BY CELEX CORPORATION, Plymouth, MI 48170

EPA Est. No. 66515-MI-1

PA Reg. No. 66515-16