

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)
CAUTION**

HARMFUL IF SWALLOWED: Inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. In the home, all food processing services and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquanums before spraying.

PHYSICAL OR CHEMICAL HAZARDS

Extinguish open flame before space spraying. This formulation is combustible. Do not use, pour, spill, or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water.

Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. For containers of 1 gallon or less. Do not reuse empty container. Wrap and put in trash.

GENERAL: Consult federal, state or local disposal authorities for approved alternate procedures.



0-7 # 41577-1
11-17

**PYRA-FOG 6
CONTACT AND SPACE SPRAY**

Kills the accessible exposed stages on contact:
Granary Weevil, Rice Weevil, Cadelle, Confused Flour Beetle, Sawtooth Grain Beetle, Angoumois Grain Moth, Indian Meal Moth, Cigarette Beetle, Drugstore Beetle, Lesser Grain Borer, Tobacco Moth, Red Flour Beetle, Fruit Flies, Grain Mites, Vinegar Flies, Mediterranean Flour Moth, House Flies, Gnats, Cockroaches, Silverfish, Spiders, Ants, Crickets, Mushroom Flies, Fungus Gnats, Horn Flies, and Stable Flies.

ACTIVE INGREDIENTS

Pyrethrins	0.60%
Piperonyl Butoxide, Technical	3.00%
Petroleum Distillate	96.40%
	100.00%
*Equivalent to 2.4% of (Butylcarbityl) (6-Propylpiperonyl) Ether and 0.6% of related compounds.	

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of water. Get medical attention if irritation persists.

MANUFACTURED BY:

Dickson Chemical Co., Inc.

STUTTGART, ARKANSAS 72160

EPA Reg. No. 46579-7

EPA Est. 46579 AR-1

NET CONTENTS:

Gallons

Authorized for use in "Fly-free Product Areas" of Official Establishments Operating Under the Meat Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants, Flour Mills, Rice Mills, Feed Processing Plants, Empty Grain Bins, Grain Elevators, Grainaries, Shipholds, Barges, boxcars, Grain Trucks, Warehouses, Feed and Seed Stores, Grocery Stores, Restaurants, Hotels, Breweries, Canneries and Bakeries.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

SPACE SPRAY: Close room or area to be treated. Shut off air conditioner and ventilator system. Seal all open vents. Remove pets, birds and cover fish aquariums. Apply undiluted as a fine mist of aerosol size particles through manual, electrical or other powered applicators. Direct the mist toward the ceiling and upper corners of the area to be treated. Fill the area with the mist at the rate of 1 FL ounce per 1,000 cubic feet leave area closed for 30 minutes. Ventilate thoroughly before re-entry. Sweep up and destroy dead insects.

In boxcars, ship cargo holds, barges and trucks, cover any loose or unpackaged finished foods with paper or plastic tarps. Apply space spray at a rate of 1 fl. ounce per 1,000 cubic feet of space. Close all doors and openings before applying spray and leave closed for 30 minutes after total dosage has been applied. Ventilate thoroughly before re-entry. Sweep up and destroy dead insects.

CONTACT SPRAY: To kill all listed crawling insects. Adjust spray to produce large wetting type spray particles. Spray insects directly whenever possible. Direct spray to contact insects in cracks and crevices in floors, around pipes, beams, behind cabinets, around machinery and other harborages. Spray these infested areas until wet, repeat as necessary. Avoid contamination of feed or food stuffs.

SPACE SPRAY: For use in barns and poultry houses. To control House Flies, Stable Flies & Horn Flies, apply as a fine mist over the livestock in stanchions or holding pens or over the poultry in cages or in pens. Use a fogger or mist generator, so spray is directed and applied above the animals and direct spray to center of the area where animals are housed. Apply 1 oz. per 1,000 cu. ft., so that the spray mist is distributed uniformly throughout the entire area sprayed. Do not ventilate the area for at least 30 minutes after spray has been applied. If used in Dairy Barns, aerate the barn thoroughly before milking. Repeat daily, morning and afternoon or as required. Wash teats of dairy animals before milking.

IN MUSHROOM PRODUCTION AND PROCESSING: To control mushroom flies or fungus gnats in growing houses, apply at a rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2,000 cubic feet of space.

NON WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

WYNN

2011

100 + 100 77
r 17
1 2000

ACCEPTED
FEB 14 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
Reg. No. **46579-**