

## DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. For containers of 1 gallon or less - Do not reuse empty container. Wrap and put in trash.

**GENERAL:** Consult federal, state or local disposal authorities for approved alternate procedures.

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants, Flour Mills, Rice Mills, Feed Processing Plants, Empty Grain Bins, Grain Elevators, Granaries, Shipholds, Barges, Boxcars, Grain Trucks, Warehouses, Feed and Seed Stores, Grocery Stores, Restaurants, Hotels, Breweries, Canneries, Bakeries, Cattle Barns, Horse Barns, Poultry Houses, Swine Houses, Zoos, Mushroom Production and Processing, and Outdoor Application.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

10.00%  
81.00%  
100.00%  
Equivalent to 4.80% of (Butylcarbyl) (6-Propylpiperonyl) Ether and 1.20% of related compounds.

KEEP OUT OF REACH OF CHILDREN

# CAUTION

SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

## STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Do not induce vomiting because of aspiration hazard. Contact a physician immediately.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**IF ON SKIN:** Wash with plenty of water. Get medical attention if irritation persists.

MANUFACTURED BY:

*Dickson Chemical Co., Inc.*  
STUTT GART, ARKANSAS 72160

EPA Reg. No. 46579-B  
EPA EST. 46579-AR-1

NET CONTENTS

Gallons

Infested stored products should be fumigated or treated by other effective methods using all approved product intended for this purpose. Use a dosage of one-half fluid ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application will kill accessible, exposed stages of those stored product pests listed above. Sweep up and destroy fallen insects.

## INDOOR USE AS A SPACE SPRAY, DILUTED:

For use in conventional mechanical logging equipment cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

## SURFACE SPRAY:

To insure maximum control of crawling insects, dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well and apply as a coarse wetting spray using approximately 1 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in transit protection, clean and spray truck beds, boxcars, and ships hold before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

To control INDIAN MEAL MOTHS, and ANGOUMOIS GRAIN MOTHS in stored grain apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early fall. In case of severe infestation, break up webbing with rake before spraying and make a second application in two weeks.

As an aid in control of MOSQUITOES, GNATS, WASPS AND SKIPPERS: treat walls, ceilings, mouldings, screens, door and window frames, beams, light cords and similar resting places.

**ANIMAL QUARTER USE** (Cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray diluted for use in conventional mechanical logging equipment to kill

exposed or accessible stages of listed insects.

Before using, exposed food products and packaging materials must be covered or removed. After use, equipment materials must be thoroughly rinsed with potable water before operations are resumed. Use as a space spray only when plant is not in operation.

## OUTDOOR GROUND APPLICATION, UNDILUTE:

To control ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS. For best results, apply when there is a light breeze of approximately 5 mph and temperature is cool (70 or less). With a ULV spraying device spray at a dosage rate of 5 ounces per minute upwind of the area to be controlled. If the equipment is truck mounted, keep the speed approximately 5 mph with a swath width of no more than 300 feet.

To control MOSQUITO LARVAE, dilute at the rate of 1 ounce per 50 gallons of oil and apply at the rate of 20 to 30 gallons per acre. Apply as a coarse wetting spray to 1 border or on shallow water of wet lands, swamps and marsh where mosquito larvae are known to breed.

## TEMPORARY REDUCTION OF ANNOYANCE FROM FLIES, MOSQUITOS and SMALL FLYING MOTHS OUTDOORS.

The above mentioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theatres, golf courses, municipal parks, playgrounds and recreational areas). Direct application into tall grass, shrubbery and around lawns where the pests may hover or rest, apply while air is still. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary.

In additional outdoor areas (corral, feed lots, swine lots, zoos), cover water drinking fountains and animal feed belf use. Fill area with mist, directing application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. In zoos avoid exposure of reptiles to the product. Repeat as necessary.

## NON WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.