

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND (DOMESTIC ANIMALS)
CAUTION**

HARMFUL IF SWALLOWED. Inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Remove pets, birds and cover fish aquariums before spraying. In case of contact, wash immediately with soap and water. Wash arms, hands and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of operational area while applying the insecticide, or working in treated areas during posted period. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water when disposing of equipment washwaters. Remove pets and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS

Keep container closed when not in use

Keep away from heat and open flame

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. For containers of 1 gallon or less - Do not reuse empty container. Wrap and put in trash.

GENERAL: Consult federal, state or local disposal authorities for approved alternate procedures.

Designed for Use in Establishments Operating Under the Meat, Poultry, Shell Egg, Grading and Egg Products Inspection Programs and in Other Food Processing Plants, Flour Mills, Rice Mills, Feed Processing Plants, Emph. Grain Bins, Grain Elevators, Granaries, Shipholds, Barges, Boxcars, Grain Trucks, Warehouses, Feed and Seed Stores, Grocery Stores, Restaurants, Hotels, Breweries, Canneries, Bakeries, Cattle Barns, Horse Barns, Poultry Houses, Swine Houses, Zoos, Mushroom Production and Processing, and Outdoor Application.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist



**PYRA-FOG MAGNUM 3
CONTACT AND SPACE SPRAY**

Kills the accessible exposed stages on contact: Granary Weevil, Rice Weevil, Cadelle, Confused Flour Beetle, Sawtooth Grain Beetle, Angoumois Grain Moth, Indian Meal Moth, Cigarette Beetle, Drugstore Beetle, Lesser Grain Borer, Tobacco Moth, Red Flour Beetle, Fruit Flies, Grain Mites, Vinegar Flies, Mediterranean Flour Moth, Houseflies, Gnats, Cockroaches, Silverfish, Spiders, Ants, Crickets, Mushroom Flies, Fungus Gnats, Horn Flies, Stable Flies, Cheese Skippers, Mosquito Larvae, Mosquitoes, and Small Flying Moth.

ACTIVE INGREDIENTS

Pyrethrins	1.00%
Piperonyl Butoxide Technical	6.00%
% Octyl Bicycloheptene Dicarboximide	10.00%
<i>†† Insect Ingredients</i>	81.00%
	100.00%
<i>*Equivalent to 4.80% of (Butylcarbityl) (6-Propylpiperonyl) Ether and 1.20% of related compounds.</i>	
<i>†† Contains Retrokum Distillate</i>	

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting because of aspiration hazard. Contact a physician immediately.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of water. Get medical attention if irritation persists.

MANUFACTURED BY

Dickson Chemical Co., Inc.

STUTT GART, ARKANSAS 72160

EPA Reg. No. 46579-8
EPA EST. 46579-AR-1

NET CONTENTS:

Gallons

Not Reviewed. Registrant claims to be in accordance with Reregistration or Pesticide Product Guidance for Phase 2 Response. REEVALUATION USE AS A SPACE SPRAY

UNDILUTED:

Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at a rate of 1/2 ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. Sweep up and destroy fallen insects.

DRIED FRUIT PRODUCTS in storage or being processed for control of exposed listed insects. Remove and destroy infested products. As noted above disperse this product as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use one-half ounce per 1,000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate the area before re-entry. Sweep up and destroy fallen insects.

STORED FOOD: (Stored in multi-wall bags or cloth bags) in warehouses, storage rooms, flour mills, grain elevators, granaries. Disperse this product first by directing spray into corners, under pallets, and around stacks of packaged foods hitting as many insects as possible and then treat as a space spray.

Infested stored products should be fumigated or treated by other effective methods using all approved product intended for this purpose. Use a dosage of one-half fluid ounce per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application will kill accessible, exposed stages of those stored product pests listed above. Sweep up and destroy fallen insects.

INDOOR USE AS A SPACE SPRAY, DILUTED:

For use in conventional mechanical fogging equipment cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating system equipment. Dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

SURFACE SPRAY:

To insure maximum control of crawling insects, dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well and apply as a coarse wetting spray using approximately 1 gallon of spray mixture to 750 square feet of surface. Spray walls, floor and other surfaces of bins and storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Treat unloading, handling and processing areas and inside conveying processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in transit protection clean and spray truck beds, boxcars, and ship's hold before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

To control INDIAN MEAL MOTHS, and ANGOUMOIS GRAP. MOTHS in stored grain apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In case of severe infestation, break up webbing with rake before spraying and make a second application in two weeks.

As an aid in control of MOSQUITOES, GNATS WASPS AND SKIPPERS: treat walls, ceilings, moldings, screens, door and window frames, beams, light cords and similar resting places.

ANIMAL QUARTER USE (Cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray diluted for use in conventional mechanical fogging equipment to kill

MOSQUITOES, SMALL FLYING MOTHS, AND GNATS, dilute 1 part concentrate plus 29 parts oil (refined kerosene) and mix well. Apply at the rate of 2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray towards the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs. Wash teats of dairy animals before milking.

IN MUSHROOM PRODUCTION AND PROCESSING:

To control exposed insect in growing houses, dilute one part of concentrate plus 29 parts oil (refined kerosene) and mix well and apply at a rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and should be directed slightly upward over them. Circulating fans should be left in operation but for best results, refrigeration should be turned off. In packaging and canning plants use at a rate of one-half ounce per 1,000 cubic feet of space.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS:

Dilute one part concentrate with 3 parts base oil (refined kerosene) and use with a ULV spraying device at the rate of one ounce per 1,000 cubic feet of room space. Apply the spray into all cracks and crevices in wood work, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Contact as many insects as possible. Repeat as necessary for control of exposed or accessible stages of listed insects.

Before using, exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be thoroughly rinsed with potable water before operations are resumed. Use as a space spray only when plant is not in operation.

OUTDOOR GROUND APPLICATION, UNDILUTED:

To control ADULT MOSQUITOES, GNATS AND SMALL FLYING MOTHS. For best results, apply when there is a light breeze of approximately 5 mph and temperature is cool (75° or less). With a ULV spraying device spray at a dosage rate of 5 ounces per minute up-wind of the area to be controlled. If the equipment is truck mounted keep the speed to approximately 5 mph with a swath width of no more than 300 feet.

TEMPORARY REDUCTION OF ANNOYANCE

from FLIES, MOSQUITOS and SMALL FLYING MOTHS OUTDOORS. The above-mentioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theatres, golf courses, municipalities, parks, playgrounds and recreational areas). Direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary.

In additional outdoor areas (corrals, feed lots, wire lots and zoos), cover water drinking fountains and animal feed before use. Fill area with mist, directing application into tall grass, shrubbery and around lawns where these pests may hover or rest apply while air is still. Avoid wetting foliage. In zoos, avoid exposure of reptiles to the product. Repeat as necessary.

NON WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.