

300 | 179,824
17 |

Mr. Jeffery A. Stone
Chemstar Corporation
2501 NW. 7th Street
Miami, FL 33147

Dear Mr. Stone:

Subject: Chemstar Chemaquacide
EPA Registration No. 46597-1
Your Submission of August 24, 1984

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.


Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

RD:W.E.Nelson:DCR-45294:WANG-0574K:KIM:Raven:479-2013:9/14/84:Del.9/30/84

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							



Chemstar Corp.

2501 N.W. 75th Street • Miami, Florida 33147 • Phone: (305) 836-6066
U.S.A. (Toll Free) 1-800-327-0777 • Fla. (Toll Free) 1-800-432-2804

CHEMSTAR CHEMAQUACIDE AQUEOUS INSECTICIDE

For use in institutions, hotels, restaurants, supermarkets,
Food Processing Plants, Canned and Livestock
USDA APPROVED For use in Meat, Poultry, and Egg Grading
and Egg Processing Areas

Apply with Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS

Pyrimins	0.1%
Equiperthrin	1.0%
Petroleum Distillate	0.1%
INERT INGREDIENTS	98.8%

Equivalent to 8% cyfluthrin, 6% cyfloperoxy, ether and
2% related compounds

EPA Reg. No. 415-07-1 EPA Est. No. 415-07-1

SEPT 10 1984

46597-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional cautions

DIRECTIONS FOR USE

The use of this product in food processing establishments where the plant is not in operation. Food should be removed from contact with surfaces should be covered during treatment and ventilated before reoccupying. For maximum effectiveness, fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, mites, granary weevils, rice weevils, confused flour beetles, spider beetles, cigarette beetles, drug store beetles, etc. Use a good sprayer adjusted to a coarse wet spray. Apply to crevices, under pallets, around containers of food, shelves and drawers. For silverfish spray books, surface application only is to be used spray for 750 square feet of surface.

SPACE SPRAYING: Close doors and windows. Use sprayer machine adjusted to deliver a fine mist or fog. Control house flies, skipper flies, wasps, hornets, Angoumois grain moth, and tobacco moths. Spray at an angle to the surface of the entire area at a rate of 0.5 to 1.0 ounces per 1000 cubic feet.

FOGGING: To kill exposed insects in all possible stages of development. Apply at a rate of 20,000 to 40,000 cubic feet of space per gallon.

ON ANIMALS: To protect cattle and horses from house flies, apply a light mist sufficient to wet the surface of the animal. Apply at a rate of 2 ounces per adult cow or horse daily or at intervals necessary to give continuous protection. To control blood-sucking lice, apply to the neck and spray to the base of the hair. Repeat every 2 weeks.

