46781-10

DEC = 6 2007

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

1216/200

United States Environmental Protection Office of Pesticide Programs

Metrex Research Corporation 1717 West Collins Avenue Orange, CA 92867

Attention: Wendy Garman, Manager Regulatory Affairs

Subject: Premicide EPA Registration No. 46781-10 Notification Dated November 5, 2007

This will acknowledge receipt of your notification to update the Basic formulation by adding a new facility to the list of producers and to make minor changes to the product label, submitted under the provisions of FIFRA Section 3(c)(9). Based on a review of the submitted material, the following comments apply.

The Notification dated November 5, 2007 is in compliance with PR Notice 98-10 and is acceptable. The information has been added to your file.

If you have any questions concerning this letter, please contact Martha Terry at (703) 308-6217.

Sincerely

Marshall Swindell Product Manager (33) Regulatory Management Branch 1 Antimcrobials Division (7510P)

Enclosure

Flease read instructions on reverse before comprising fo	prm. Form Ap	proved. OMB No. 2070-0060, Approvel expires 2-28-95				
United Environmental Pro Washington,	otection Agency	Registration OPP Identifier Number Amendment ✓				
Арр	lication for Pesticide - Sec	ction I				
ompany/Product Number 2. EPA Product Manager 3. Proposed Classification 6781 / 46781-10 M. Swindell Vone Restricted						
 Company/Product (Name) Metrex Research Corporation / Premicide 	PM# 33					
5. Name and Address of Applicant <i>(Include ZIP Code)</i> Metrex Research Corporation 1717 West Collins Avenue Orange, CA 92867	(b)(i), my product to:	veiw. In accordance with FIFRA Section 3(c)(3) is similar or identical in composition and labeling				
	Section - II					
Amendment - Explain below. Resubmission in response to Agency letter dated V Notification - Explain below.	Agency let	Final printed labels in repsonse to Agency letter dated "Me Too" Application. Other - Explain below.				
added to the enclosed proposed Premicide label. Pak Technologies, 8000 W Good Hope Rd., Milwauker This notification is consistent with the provisions of PR Notice Statement of Formula of this product. I understand that it is a	e, WI 53223 e 98-10 and EPA regulations at 40 CFR 152.46, a a violation of 18 U.S.C. Section 1001 to willfully ma	oducers on page 3. Additionally, the EPA Est. No. 51435-WI-004 was nd no other changes have been made to the labeling or the Confidential ake any false statement to EPA. I further understand that if this notification RA and I may be subject to enforcement action and penalties under				
	Section - III					
	Water Soluble Packaging Yes No If "Yes" No. per Itainer Package wgt containe					
3. Location of Net Contents Information 4. Size Label Container 6. Manner in Which Label is Affixed to Product	Lithograph Othe	S. Location of Label Directions On Label On Labeling accompanying product				
	Section - IV					
1. Contact Point <i>(Complete items directly below for iden</i> Name Wendy Garman	ntification of individual to be contacted, Title Manager, Regulatory Affairs	s 714-516-7602				
C I certify that the statements I have made on this f I acknowledge that any knowlinglly false or mislea both under applicable law.						
Afrely Garman	3. Title Manager, Regulatory Affairs					
4. Typed Name Wendy Garman	5. Date	7				

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

.



November 5, 2007

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) US Environmental Protection Agency Room S-4900, One Potomac Yard 2777 S. Crystal Drive Arlington, VA 22202-4501

Re: NOTIFICATION TO REVISE THE CONFIDENTIAL STATEMENT OF FORMULA (BASIC FORMULATION)

Product: Company: Premicide, EPA Reg. No. 46781-10 Metrex Research Corporation

Dear Sir or Madam:

Enclosed is the proposed Confidential Statement of Formula (Basic Formulation) of Premicide, EPA Reg. No. 46781-10. The following additional manufacturing facility has been added to the list of producers. Please also note that there have been no changes to any of the nominal (column 13b) or certified limit values (columns 14a and 14b).

> Pak Technologies 8000 W Good Hope Rd. Milwaukee, Wl 53223

Additionally, the EPA Est. No. 51435-WI-004 was added to the enclosed proposed Premicide label. It is requested that this label be stamped for acceptance as the most current label for the product.

The following items are being submitted in support of this amendment:

- Two copies of the Application for Pesticide Form (Section 1)*
- One copy of the <u>current</u> Premicide CSF (Basic Formulation), dated 10/10/06 (Section 2)*
- Two copies of the proposed Premicide CSF (Basic Formulation), dated November 5, 2007 (Section 3)*
- One copy of the proposed Premicide label with changes highlighted (Section:4)*

* Please treat the information contained in Sections 1-4 as proprietary and confidential as covered in the provisions of FIFRA Section 10.

If you have any questions regarding this submission, please contact Orlando Tadeo, Jr. or me at any of the following numbers/addresses:

Phone Numbers: Fax Number: Email addresses: Company Address:

i

O. Tadeo, Jr. (714) 516-7419 or W. Garman (714) 516-7602 (714) 516-7472 orlando.tadeo@sybrondental.com or wendy.garman@sybrondental.com Metrex Research Corporation 1717 West Collins Avenue Orange, CA 92867

Sincerely,

Wendy Garman Manager, Regulatory Affairs

Enclosures

(

PREMICIDE Sanitizer/Solidifier

The efficacy of this product has been demonstrated in the presence of 100% blood serum.

ACTIVE INGREDIENTS:

Glutaraldehyde	
OTHER INGREDIENTS:	
Total	100.00%

KEEP OUT OF REACH OF CHILDREN DANGER

	FIRST AID
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
lf on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
lf inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
	ncy information on product, call Metrex at 1-800-841-1428, Monday through Friday, between 6 a.m. and 4 p.m. Pacific Time. 4 p.m., call Chemtrec 24-Hour Emergency Service at 1-800-424-9300.
	uct container or label with you when calling a poison control center or doctor, or going for treatment.
Prot	NOTE TO PHYSICIAN: bable mucosal damage may contraindicate the use of gastric lavage.
	See side panel for additional precautionary statements.

Net Contents: XX Solidifies up to XXXXcc

> Page 1 of 7 ver. 10.29.2007

[Side Panel]

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals **DANGER**

Corrosive. Causes irreversible eye damage. Causes skin irritation. Harmful if inhaled. Harmful if swallowed. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Causes asthmatic signs and symptoms in hyper-reactive individuals.

- Do not get in eyes, on skin or on clothing.
- Avoid breathing vapor.
- Do not swallow.
- Wear goggles, protective clothing and rubber gloves.
- Wash thoroughly with soap and water after handling.
- Remove contaminated clothing and wash before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry, well-ventilated area.

Pesticide Disposal: Dispose of waste according to Federal, State and local regulations. **Container Disposal:** Do not reuse container. Triple rinse. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR SINGLE USE ONLY

[Side Panel A -The following verbiage will appear only on the 500cc, 1200cc, 1500cc, 2000cc and 3000cc bottles]

Fast-acting encapsulant which contains scented dry glutaraldehyde to sanitize and solidify liquid medical waste and laboratory, human and animal wastes. Antibacterial. Controls odor formation by destroying odor-causing bacteria.

The efficacy of this product has been demonstrated in the presence of 100% blood serum.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To help prevent the blood from clotting inside the suction canister, immediately dispense PremiCide following the end of surgery or after the suction canister has been completely filled.

- Wear hand, eye and face protection when handling infectious waste and using PremiCide.
- Shake and squeeze bottle to loosen any settled material.
- Following surgery, open the appropriate port and place the PremiCide cap onto the open port and push down to secure proper fitting.
- Following surgery, turn off suction and remove the patient tubing. Close all ports and activate suction. Firmly grasp the base of the cap and pull upward on the bottle. Suction will pull the PremiCide powder into the canister.
- Once the PremiCide bottle is empty, tap the side of the bottle to remove any remaining powder. Push down on the bottle to close the system. Once the liquid waste is solidified, remove the bottle by gripping the base of the PremiCide cap and lift up. Replace the port cap on the suction canister. To assure thorough treatment of the suction canister, invert solidified canister and dispose of according to institutional procedures and Federal, State and local regulations.

For product or technical information, contact Metrex Customer Care at **800-841-1428** or visit our website at **www.metrex.com**.

EPA Reg. No. 46781-10

EPA Est. No. 72075-WI-001 46781-MI-001 51435-WI-004

Reorder No: 00-0000

Packed For: [or] Manufactured By: Metrex Research Corporation 28210 Wick Road Romulus, Michigan 48174 [Side Panel B - The following verbiage will appear only on the 10000cc multi-dose bottle]

Fast-acting encapsulant which contains scented dry glutaraldehyde to sanitize and solidify liquid medical waste and laboratory, human and animal wastes. Antibacterial. Controls odor formation by destroying odor-causing bacteria.

The efficacy of this product has been demonstrated in the presence of 100% blood serum.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To help prevent the blood from clotting inside the suction canister, immediately dispense PremiCide following the end of surgery or after the suction canister has been completely filled.

- Wear hand, eye and face protection when handling infectious waste and using PremiCide.
- Shake and squeeze bottle to loosen any settled material.
- Unscrew cap, remove safety seal and recap.
- Open a port in the top of the suction canister or provide access to the liquid waste if other device is used. Remove red cap.
- Insert PremiCide nozzle into opening. Gently squeeze the bottle, injecting PremiCide into the canister. Do not allow the tip of the bottle to touch the liquid waste. Recap port and replace red cap on bottle. Each graduation solidifies 1,000cc's.
- Allow sufficient time for encapsulation before handling.

For product or technical information, contact Metrex Customer Care at **800-841-1428** or visit our website at **www.metrex.com**.

EPA Reg. No. 46781-10

EPA Est. No. 72075-WI-001 46781-MI-001 51435-WI-004 Reorder No: 00-0000

Packed For: [or] Manufactured By: Metrex Research Corporation 28210 Wick Road Romulus, Michigan 48174 [The following optional verbiage will appear on a sticker that may be packaged with the product]

Chemically Treated Solid Non-Infectious Medical Waste

Page 5 of 7 ver. 10.29.2007 [The following verbiage only appears on labeling for PremiCide CA, Alternate Brand Name of PremiCide]

[Front of sticker label]

PremiCide

Determination of Canister Disposal (see procedure on back of label)

Canister No.

Calculation

A. Suction canister EBL (Note Amount) _____ cc
B. Total cc fluid in suction canister (Note Amount) _____ cc
C. Total % blood in suction canister (A/B x 100 = C) _____ %

Refer to "C" and circle only one (1) of the following:

White BagRed Bag(≤20% blood in canister)(> 20% discard as medical waste)

Not for use in post-surgery body cavity drainage.

[Back of sticker label]

Procedure for disposal of suction canister when using PremiCide

Important Note: -

Following the completion of surgery it is important to determine the amount of blood contained in a suction canister used during a surgical procedure. The amount of blood in the canister must be determined to properly dispose of the canister.

Instructions:

- 1. Following the completion of the procedure, the participating anesthesiologist shall calculate the blood content of each suction canister used during surgery. The calculation shall be made using the information generated by the anesthesiologist when calculating the estimated blood loss.
- 2. The estimate shall be recorded on this PremiCide disposal label, which is then affixed to the suction canisters used during surgery. The label shall be completed as follows:
 - Item A: Record the "EBL" or "Estimate Blood Loss" made by the anesthesiologist in total cc's. Exclude all estimates for laps, spills, or other areas where blood may have been lost but was not captured in suction canisters.
 - Item B: Record the total amount of fluid in cc's in the suction canister. This will include all fluids (blood, irrigants, or other fluids everything in the canister).
 - Item C: Calculate A / B x 100 = "C" (Estimate of total percent blood in canister): Example:

"A"	/	"B"	х	100	.=	"C"
300 cc EBL in canister	/	3000 cc total fluid in canister			=	10% blood

Page 6 of 7 ver. 10.29.2007 [The following verbiage only appears on labeling for PremiCide CA, Alternate Brand Name of PremiCide]

- 3. Affix label to suction canister.
- 4. If Item "C" is less than or equal to 20% circle "White Bag" method of disposal. If Item "C" is greater than 20% circle "Red Bag" method of disposal.

For product or technical information, contact your local Metrex representative at 1-800-841-1428 or visit our website at www.metrex.com.

Page 7 of 7 ver. 10.29.2007