

Easy-Open Bottom
SEE DIRECTIONS ON BUTT FLAP

ACCEPTED
SEP 15 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No. 46946-59

OUTER
Fungicide

wettable powder in water soluble bags

ACTIVE INGREDIENT

Triphenyltin hydroxide* 47.5%

INERT INGREDIENTS 52.5%

100.0%

*Equivalent to 15.3% Metallic tin

KEEP OUT OF REACH OF CHILDREN
DANGER

STATEMENT OF PRACTICAL TREATMENT

IF INHALED—Remove victim to fresh air and apply respiration if indicated.
IF ON SKIN—Wash promptly with soap and water. Rinse thoroughly.
IF IN EYES—Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

A RESEARCH DISCOVERY OF DUPHAR B.V. AMSTERDAM HOLLAND
Manufactured or sold under one or more of the following U.S. Patents: 3,113,069,
3,140,977, 3,248,283, 3,268,395, 3,395,228, 3,445,575, and other Patents Pending

EPA Reg. No. 46946-59 4-82 EPA Est. No. 148-MS-1
NET CONTENTS 2 LBS. (907 GRAMS)

THE AGRICULTURAL CHEMICAL COMPANY, INC.
CROP PROTECTION DIVISION

DO NOT OPEN UNTIL READY TO USE,
DO NOT USE LESS THAN FULL 2LB. BAG.

IMPORTANT — How To Use This Package:

1. Lift bottom flap and remove hooklet.
2. Follow bag opening instructions and allow water soluble inner bag to drop directly into spray tank.
3. Destroy outer bag.

This product intended for large acreage use only.

SEE FOLDER IN BOTTOM FLAP FOR USE INSTRUCTIONS

A FUNGICIDE FOR CONTROL OF
• EARLY & LATE BLIGHT ON POTATOES
• LEAFSPOT ON SUGAR BEETS & PEANUTS
• SCAB & CERTAIN OTHER DISEASES ON PECANS
• LEAFSPOT AND ALTERNARIA ON CARROTS

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with or inhalation of spray mist. Avoid contamination of foodstuffs and feed. Wear clean protective clothing and bathe after each exposure.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

BEFORE USE, READ DIRECTIONS, WARNING AND NOTICE STATEMENTS CAREFULLY. Consult agricultural extension or agricultural experiment station specialists regarding timing recommendations.

GENERAL INFORMATION

Full coverage is necessary for best results. Diluted spray should be directed to all parts of the plant uniformly. Application rates increased according to size of plants. Use the lowest application rates in protective schedules. Higher rates later in the season or during high infection periods.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

MIXING

Before adding DU TER to the spray tank, partially fill with water. Add the desired amount of DU TER according to the rates listed elsewhere. After mixing thoroughly, add remaining water to spray tank. Continuous agitation within the spray tank is essential. If used in combination with liquid products, place DU TER in tank first and when completely wetted, add emulsifiable concentrate or other liquid product. Wash out spray tank thoroughly after use and dispose of washings by burying away from water supplies. Observe all cautions and limitations on labeling of all products used in mixtures.

APPLICATION METHODS

GROUND: Mix with water for a total solution of 10-100 gallons per acre.

AERI: Mix with water for a total solution of 3-20 gallons per acre. **CAUTION:** In non mobile systems, application by irrigation system must be equipped with safety valves or other devices to prevent back-siphoning of pesticide into water source. Any irrigation water treated with this product should be held on the treated area until it is absorbed by the soil. It should not be turned into the soil water or fed back into the irrigation system. **Irrigation Equipment:** Tank connected to the suction side of irrigation pump, hand sprayer or other pressurized equipment connected to the main irrigation line may be used for injection. Good agitation must be provided in injection tank.

CALIBRATION AND USE IN SET SYSTEMS AND OTHER NON-MOBILE SYSTEMS:

1. Measure acreage covered by sprayers.
2. Operate irrigation system and injection equipment at normal pressures.
3. Fill tank of injection equipment.
4. Adjust flow from injection equipment to use contents over a period of 10 to 30 minutes.
5. For application of DU TER, mix desired amount of product for acreage to be covered (see chart for rates) into quantity of water used during prior calibration and operate entire system.

1. Amount of dye established during calibration (DU TER) should be injected during the irrigation or as a separate application. 2. Slug test equipment after treatment is completed and continue to operate irrigation system until DU TER has been cleared from the last sprinkler head. Refer to dye method discussed below.

CALIBRATION AND USE IN CENTER-PIVOT AND OTHER MOBILE SYSTEMS:

1. Operate system and injection equipment at normal pressures.
2. Fill tank of injection equipment.
3. a) For systems other than center pivot, operate entire system for one complete trip, measure the required amount of water used from injection tank and acreage covered. b) For center pivot systems, the water flow, speeds, acreage and other calculations of the system are given in the manual or should be determined.
4. For application of DU TER, mix desired amount of product for acreage to be covered, use chart for rate, mix quantity of water used during prior calibration and inject into system continuously for one complete coverage of the field.
5. If irrigation is to be stopped after DU TER has been applied, shut off the injection equipment after treatment is completed and continue to operate irrigation system until DU TER has been cleared from the last sprinkler head. Refer to the method discussed below.
6. With mobile systems, irrigation may be continued after injection has been stopped.

DYE CALIBRATION

A recommended procedure for establishing the time when the dye is cleared from stopping the injection of DU TER to its clearance from the last sprinkler is as follows:

1. During calibration of injection equipment as described above, insert a water soluble Food, Drug and Cosmetic approved dye into the irrigation line at the injection point and record the time it takes for the dye to be released from the last sprinkler head.
2. Collection of water from the last sprinkler in a glass jar will assist in observing when dyed water reaches that point.



USE DIRECTIONS

Do not detach from container

4-82

CROP DISEASE	RATE A	REMARKS	PRECAUTIONS
Affect to Cercospora botrytis	4-8 oz	Application should begin when weather conditions favor disease development and continue at 7-day intervals until the ground is covered with a 1/4 inch of rain.	Do not use tops as feed for livestock. Do not add surfactants, spreaders or stickers to sprays. Do not graze dairy or meat animals in treated areas. Do not apply within 14 days of harvest.
POTATOES Early and Late Blight Botrytis gray mold	5-10 oz (6-10 oz through irrigation) ground min 15 gpa aerial 3-10 gpa	Application should begin when weather conditions favor disease development and continue at 7-day intervals until the ground is covered with a 1/4 inch of rain. (Irrigation only for Botrytis gray mold.)	Do not add surfactants, spreaders or stickers. Do not graze dairy or meat animals in treated areas. Do not apply within 7 days of harvest. May be used in combination with wettable powder and emulsifiable concentrate insecticides; however, before using with emulsifiable concentrates, it is suggested that the mixture be tested for phytotoxic response under actual use conditions. For helicopter application fly high enough so as to not whip the vines.
PECAN Scab Brown Leaf spot Downy Spot Powdery Mildew Liver Spot Sooty Mold Leaf Blotch	75-15# or 3-2-9-6 oz per 100 gallons— ground (1-3# per 500 gallon tank)	Application should be at pre-pollination stages when the young leaves are unfolding and a second application made when the small nuts are forming. Repeat at 2-4 week intervals as needed. To maintain control use lower rate for first two applications or until disease becomes severe or during dry weather. Use higher rate during wet weather or during severe disease infection.	Do not apply after nuts have started to open. Do not graze dairy or meat animals in treated areas. EC-R may be applied in combination with emulsifiable concentrate or wettable powder insecticides.
CARROTS Cercospora Leafspot Alternaria blight	4-8 oz	Applications should begin 6 weeks after planting or as soon as first signs of disease appear. Continue at 7-day intervals or as needed for control.	Do not use tops as feed for livestock. Do not add surfactants, spreaders or stickers to sprays. Do not graze dairy or meat animals in treated areas. Do not apply within 14 days of harvest.
PEANUTS Cercospora Leafspot (& Spider Mite Suppression)	5-6 oz	Application should begin approximately 6 weeks after planting or as soon as first signs of leafspot appear. Spray applications should continue on a 10-14 day schedule.	Do not allow hogs to feed on peanuts in treated fields. Do not use vines as feed. Do not graze dairy or meat animals in treated fields. Do not add surfactants, spreaders or stickers to sprays. Do not mix with emulsifiable concentrate or liquid formulations. Do not apply within 14 days of harvest.

- A FUNGICIDE FOR CONTROL OF
- EARLY & LATE BLIGHT ON POTATOES
 - LEAFSPOT ON SUGAR BEETS & PEANUTS
 - SCAB & CERTAIN OTHER DISEASES ON PECANS
 - LEAFSPOT AND ALTERNARIA ON CARROTS

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with or inhalation of spray mist. Avoid contamination of foodstuffs and feed. Wear clean protective clothing and bathe after each exposure.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

BEFORE USE READ DIRECTIONS, WARNING AND NOTICE STATEMENTS CAREFULLY. Consult agricultural extension or agricultural experiment station specialists regarding timing recommendations.

GENERAL INFORMATION:

Full coverage is necessary for best results. Diluted spray should be directed to all parts of the plant uniformly. Application rate increased according to size of plants. Use the lowest application rates in protective schedules, higher rates later in the season or during high infection periods.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITION: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinseate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

Easy-Open Bottom
SEE DIRECTIONS ON BUTT FLAP

ACCEPTED

SEP 15 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this pesticide is registered under EPA Reg. No. 46846-59

wettable powder in water soluble bags

ACTIVE INGREDIENT

Triphenyltin hydroxide* 47.5%

INERT INGREDIENTS 52.5%

*Equivalent to 15.3% Metallic tin 100.0%

KEEP OUT OF REACH OF CHILDREN
DANGER

STATEMENT OF PRACTICAL TREATMENT

IF INHALED - Remove victim to fresh air and apply respiration if indicated.
IF ON SKIN - Wash promptly with soap and water. Rinse thoroughly.
IF IN EYES - Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

A RESEARCH DISCOVERY OF DUPHAR B.V. AMSTERDAM, HOLLAND
Manufactured or sold under one or more of the following U.S. Patents: 3,113,069, 3,140,977, 3,248,263, 3,268,395, 3,395,228, 3,445,579, and other Patents Pending

EPA Reg. No. 46846-59 4-82 EPA Lst No. 148-MS 1
NET CONTENTS 2 LBS. (907 GRAMS)

THE UNIVERSITY OF MICHIGAN
CROP PROTECTION DIVISION

DO NOT OPEN UNTIL READY TO USE,
DO NOT USE LESS THAN FULL 2LB. BAG.

IMPORTANT — How To Use This Package:

1. Lift bottom flap and remove hooklet.
2. Follow bag opening instructions and allow water soluble inner bag to drop directly into spray tank.
3. Destroy outer bag.

This product intended for large acreage use only.

SEE FOLDER IN BOTTOM FLAP FOR USE INSTRUCTIONS

A FUNGICIDE FOR CONTROL OF
- EARLY & LATE BLIGHT ON POTATOES
- LEAF SPOT ON SUGAR BEETS & PEANUTS
- SCAB & CERTAIN OTHER DISEASES ON PECANS
- LEAF SPOT AND ALTERNARIA ON CARROTS

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with the inhalation of spray mist. Avoid contamination of foodstuffs and feed. Wear clean protective clothing and bathe after each exposure.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Do not spray directly into any body of water. Do not spray where runoff is likely to occur. Do not spray where it will drift into water. Do not spray where it will drift into water.

BEFORE USE, READ DIRECTIONS, WARNING AND NOTICE STATEMENTS CAREFULLY. Consult agricultural local extension or agricultural experiment station specialists regarding timing and recommended rates.

GENERAL INFORMATION

Full coverage is necessary for best results. Directed spray should be directed to all parts of the plant uniformly. Coverage indicated according to size of plants. Use the lowest application rates in protective schedules. Higher rates may be used in the season of during high infection periods.

DIRECTIONS FOR USE

For use on potatoes, sugar beets, peanuts, pecans, and carrots.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not burn, mix with other pesticides, or use in any way not specified on the label.
PESTICIDE DISPOSAL: Pesticide spray mixtures or residues that may be used on the label may be disposed of according to Federal, State, and local laws and regulations. Resource Conservation and Recovery Act.
CONTAINER DISPOSAL: Do not reuse or refill this container. Use and Federal, State, and local laws and regulations. Resource Conservation and Recovery Act.

CROP/DISEASE	RATE/A	REMARKS	USE CHART
SUGARBEETS Leavespot (Cercospora beticola)	4-10 oz	Applications should begin when leavespot conditions appear or when the disease is in the area and repeat at 10-14 day intervals.	Do not feed beet tops to livestock. Do not add surfactants, spreaders or stickers to DU TEH sprays. Do not graze dairy or meat animals in treated areas. Do not apply within 14 days of harvest.
POTATOES Early and late blight	5-10 oz (6-10 oz through ground-min sprayer)	Applications should begin when weather conditions favor disease development and continue on a 7-day schedule (USC combination with wettable powder and emulsifiable concentrate insecticides, acaricides, or stickers. Do not graze dairy or meat animals in treated areas. Do not apply within 7 days of harvest. May be applied in combination with emulsifiable concentrate insecticides.	Do not add surfactants, spreaders, or stickers. Do not graze dairy or meat animals in treated areas. Do not apply within 14 days of harvest. Do not use tops as feed for livestock. Do not add surfactants, spreaders or stickers to sprays. Do not mix with emulsifiable concentrate or liquid formulations. Do not apply with a 10-14 day schedule.
PEANUTS Cercospora leafspot (A. Spoor blight)	5.8 oz	Applications should begin approximately 6 weeks after planting or as soon as first signs of leafspot appear. Spray applications should continue on a 10-14 day schedule.	Do not add surfactants, spreaders or stickers to sprays. Do not mix with emulsifiable concentrate or liquid formulations. Do not apply with a 10-14 day schedule.
CARROTS Alternaria leafspot (Lewy Blight)	4.8 oz	Applications should begin 6 weeks after planting or as soon as first signs of disease appear. Continue at 7-day intervals (trial) or as needed for control.	Do not use tops as feed for livestock. Do not graze dairy or meat animals in treated areas. Do not apply within 14 days of harvest.
PEANUT Sud Brown Leaf Damping Spot Powdery Mildew Late Blight Lewy Spot Sooty Mold	75.1 oz 0.2-2.9-6 0.2 per 100 gallons 1.3-3 per 500 gallon tank)	Applications should be at pre-planting stages when the young leaves are unfolding and a second application made when the small nuts are forming. Repeat at 2-4 week intervals as needed to maintain control, or until disease becomes severe, or during dry weather or first higher rain during wet weather or during severe disease infections.	Do not add surfactants, spreaders, or stickers. Do not graze dairy or meat animals in treated areas. Do not apply within 7 days of harvest. May be applied in combination with emulsifiable concentrate insecticides.



USE DIRECTIONS
Do not detach from
container

4-82

WARNING:
When mixing DU TEH in the spray tank, partially fill the water with the desired amount of DU TEH. Then add the concentrate. Mix thoroughly. Add the water to the tank. Do not mix DU TEH with other pesticides. Do not mix DU TEH with other pesticides. Do not mix DU TEH with other pesticides.


APPLICATION METHODS
GROUND ...
AIR ...
SPRAYER ...

CALIBRATION AND USE IN SET SYSTEMS AND OTHER NON MOBILE SYSTEMS:
1. Measure the spray system's output ...
2. Measure the spray system's output ...
3. Adjust the spray system's output ...
4. Adjust the spray system's output ...
5. Adjust the spray system's output ...

...
...
...

CALIBRATION AND USE IN CENTER-PIVOT AND OTHER MOBILE SYSTEMS:
1. Operate system at normal operating pressure ...
2. Adjust the spray system's output ...
3. Adjust the spray system's output ...
4. Adjust the spray system's output ...
5. Adjust the spray system's output ...

DYE CALIBRATION
...
...
...



USE DIRECTIONS
 Do not detach from container
 4-82

MOBILE SYSTEMS
 CALIBRATION AND USE IN CENTER PIVOT AND OTHER MOBILE SYSTEMS

APPLICATION METHODS
 SPRINKLER IRRIGATION

NON-MOBILE SYSTEMS
 CALIBRATION AND USE IN SET SYSTEMS AND OTHER NON-MOBILE SYSTEMS

DYE CALIBRATION

USE CHART

CROP/DISEASE	RATE/A	REMARKS	PRECAUTIONS
SUGARBEETS Leafspot (Cercospora beticola)	4-10 oz	Application should begin when leafspot conditions appear or when the disease is in the area and repeat at 10-14 day intervals.	Do not feed beet tops to livestock. Do not add surfactants, spreaders or stickers to DU-TER sprays. Do not graze dairy or meat animals in treated areas. Do not apply within 14 days of harvest.
TOES Early and Late Blight Botrytis gray mold	5-10 oz (6-10 oz through irrigation) ground-min 15 gpa aerial 3-10 gpa.	Application should begin when weather conditions favor disease development and continue on a 7-day schedule. (Use sprinkler irrigation only for Botrytis gray mold.)	Do not add surfactants, spreaders, or stickers. Do not graze dairy or meat animals in treated areas. Do not apply within 7 days of harvest. May be used in combination with wettable powder and emulsifiable concentrate insecticides; however, before using with emulsifiable concentrates, it is suggested that the mixture be tested for phytotoxic response under actual use conditions. For helicopter application fly high enough so as to not whip the vines.
PECAN Scab Brown Leaf-spot Downy Spot Powdery Mildew Liver Spot Sooty Mold Leaf Blotch	7.5-15 # or 3.2-9.6 oz per 100 gallons ground (1.3# per 500 gallon tank)	Applications should be at pre-pollination stages when the young leaves are unfolding and a second application made when the small nuts are forming. Repeat at 2-4 week intervals as needed. To maintain control, use lower rate for first two applications, or until disease becomes severe, or during dry weather. Use higher rate during wet weather or during severe disease infections.	Do not apply after shucks have started to open. Do not graze dairy or meat animals in treated areas. DU-TER may be applied in combination with emulsifiable concentrate or wettable powder insecticides.
CARROTS Cercospora Leafspot Alternaria blight	4-8 oz.	Applications should begin 6 weeks after planting or as soon as first signs of disease appear. Continue at 7-day interval or as needed for control.	Do not use tops as feed for livestock. Do not add surfactants, spreaders or stickers to sprays. Do not graze dairy or meat animals in treated areas. Do not apply within 14 days of harvest.
PEANUTS Cercospora Leafspot (A Spider Mite Suppression)	5.8 oz	Application should begin approximately 6 weeks after planting, or as soon as first signs of leafspot appear. Spray applications should continue on a 10-14 day schedule.	Do not allow hogs to feed on peanuts in treated fields. Do not use vines as feed. Do not graze dairy or meat animals in treated fields. Do not add surfactants, spreaders or stickers to sprays. Do not mix with emulsifiable concentrate or liquid fertilizers. Do not apply within 14 days of harvest.

ICK & CO., Inc.
Way, New Jersey

PRECAUTIONARY STATEMENTS - Environmental Hazards

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal FIFSA law to use this product in a manner inconsistent with its labeling.

ELM TREES - 1 Year Treatment - Aids in the control of Dutch Elm Disease:

Preventive Treatment - For each 5 inches of trunk diameter inject 1 fluid ounce ARBOTECT 20-S in 40 fluid ounces (1 1/2 quarts) of water to 2 fluid ounces ARBOTECT 20-S in 80 fluid ounces (1 1/4 quarts) of water. Use the higher levels of ARBOTECT 20-S under high disease pressure situations.

Preventive applications should be made when leaves approach full size, usually in late May or June.

Therapeutic Treatment - For each 5 inches of trunk diameter inject 2 fluid ounces of ARBOTECT 20-S in 80 fluid ounces (2 quarts) of water to 4 fluid ounces of ARBOTECT 20-S in 160 fluid ounces of water. Use the higher levels of ARBOTECT 20-S under high disease pressure situations.

Therapeutic applications should be made as soon as the current year infection are seen, usually in late June through August.

For optimum disease control, preventive treatment is recommended. When a tree shows more than 5% crown symptoms, treatment may not be effective. Treatment should be used in conjunction with an insect control and sanitation program for ring of diseased limbs in order to obtain best results. Trees that are 5 inches or less in diameter at chest height should not be treated.

Place injection sites at near ground level (up to 3 to 10 inch intervals) and the trunk with a maximum hole diameter of 1/2 inch using a minimum of 3 or 4 equally spaced injection points per tree.

ELM TREES - 3 Year Treatment - For Preventive and Therapeutic Treatment of Dutch Elm Disease:

Inject 12 fluid ounces of ARBOTECT 20-S for each 5 inches of trunk diameter. Dilute each 20 fluid ounces of ARBOTECT 20-S with 1 gallon of water. Inject into any exposed root flares, below ground, once every three years. The maximum diameter of the injection hole should be 1/2 inch. Do not use this treatment if trees are less than 10 inches in diameter. When a tree shows more than 5% crown symptoms, treatment may not be effective. Treatment should be used in conjunction with an insect control and sanitation program for ring of diseased limbs in order to obtain best results.

SYCAMORE TREES - Aids in the control of Sycamore Anthracnose:

For each 5 inches of trunk diameter, inject 4 fluid ounces of ARBOTECT 20-S diluted with 80 to 160 fluid ounces of water (one part ARBOTECT 20-S to between 20 and 40 parts of water).

For best results, injections should be made in late summer or early fall in each of two consecutive years. Repeat treatments may be necessary if the disease recurs.

Place injection sites at 3 to 10 inch intervals around the trunk with a maximum hole diameter of 1/2 inch using a minimum of 3 or 4 equally spaced injection points per tree. Injection sites may be placed in root flares or in low ground level on the trunk as near ground level as possible. Trees that are 5 inches or less in diameter at chest height should not be treated. If pressure injectors will be used, do not exceed 100 psi.

Do not dilute ARBOTECT 20-S with highly alkaline water as a precipitate may form.

ARBOTECT 20-S is to be used by trained arborists and others trained in injection techniques and in the identification of Dutch elm disease and sycamore anthracnose.

STORAGE AND DISPOSAL

Prohibitions - Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse container.

Pesticide Disposal - Pesticide spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

Container Disposal - Triple rinse (or equivalent). Then dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

General - Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

NOTICE TO USER - Buyer assumes all risks of use and handling which are at variance in any way with the directions herein. There are no warranties which extend beyond the description of this label.

8492804