

**WARNING
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid prolonged breathing of spray mist.

If swallowed, consult a physician immediately. Note to physician: Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry, secure area in original container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER: Triple rinse (or equivalent). Then offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other approved state and local procedures.

ANIMAL BUILDINGS- Horse Stables, Dairy Barns, Auction and Show Barns (Flies, Gnats and Mosquitoes) Apply 0.5% VAPONA Insecticide as a fog or mist. 1 quart of a 0.5% solution will treat 8000 cubic feet (40x20x10). In dairy barns, before milking, the fog or mist can be applied over and around each animal to kill flies. Before applying, reduce air movement as much as possible by closing doors, windows and other openings. Do not use this solution for fogging or misting purposes in areas where animals have received a direct application within 8 hours. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Do not apply regularly to calves under 6 months

PM 15



47000-23

1/130/90
Directions for Use, Continued

**READY-TO-USE
0.5% VAPONA® INSECTICIDE
Spray or Fogging Solution**

ACTIVE INGREDIENTS:

**Dichlorvos (2,2-Di-chlorovinyl Dimethyl Phosphate 0.465%
Related Compounds 0.035%

*INERT INGREDIENTS: 99.500%
Total 100.000%

*Contains petroleum hydrocarbons
**Equivalent to 0.5% VAPONA® Insecticide

Keep Out of Reach of Children

WARNING

See Side Panel for Additional Cautions

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. If possible vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply respiration if needed.

If on skin: Wash immediately with soap and water.

If in eyes: Flush eyes for at least 15 minutes with water. Call a physician immediately.

SEE NOTE TO PHYSICIAN AND ANTIDOTE STATEMENT.

Net Contents

EPA Reg. No. 47000-23

EPA Est. No. 47000-IA-1

Not Reviewed. Registrant claims to be in accordance with registration of Pesticide Product Guidance for Phase 2 Response. Page 2.12, Active to Inert Change in Status.

Manufactured by:
CHEM TECH, LTD.
Des Moines, IA 50321



of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

OTHER USES - ANIMAL BUILDINGS
MILK SHEDS - OUTDOOR AREAS - (Flies, Gnats, Mosquitoes, Roaches, Fleas, Ants, Snowbugs, Brown Dog Ticks) Apply 0.5% VAPONA Insecticide as a coarse wet spray at the rate of one quart per 1000 square feet. For milk sheds, stables, livestock barns, loafing sheds, pig pens, and outdoor areas treat around doorways, feed-storage rooms, alleyways and window sills. Do not contaminate milk or milking utensils. In poultry houses, apply to manure, window sills, outside of penned enclosures and on floor of feed storage rooms.

Treat outdoor living areas, picnic grounds, backyard areas, patios, loading docks, outdoor latrines, parking areas, refuse areas around service stations, open-air drive-ins, ice cream stands, garbage collection and disposal areas.

In treating dog kennels apply to outside dog runways, window sills and ledges. For control of fleas and brown dog ticks increase dosage to two quarts 0.5% solution per 1000 square feet.

For control of fly maggots in maggot breeding areas such as garbage dumps and manure piles, apply at the rate of 1-2 quarts 0.5% solution per 100 square feet.

Before applying in enclosed areas, reduce air movement as much as possible by closing doors windows and other openings. Do not treat animals or humans. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Repeat as necessary.

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.