

47000-33
PM17
10F1

Received by Notification
per P.R. Notice 88-6
11/7/88

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed. Do not induce vomiting. Contact physician. If in eyes flush with water and contact physician. If on skin or clothing, remove clothing. Wash skin with soap and water. Avoid contamination of milk and milk handling equipment.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For the protection of Dairy Animals from biting flies and the control of Houseflies in Barns and Milk Houses.

Protects against Hornflies, Stableflies, Houseflies, Gnats, Mosquitoes and Blood Sucking Lice.

A temporary treatment to relieve cattle from annoyance caused by face flies.



DY-FLY II

DAIRY SPRAY

ACTIVE INGREDIENTS	100.0%
Pyrethrins	0.1%
Technical Piperonyl Butoxide**	1.0%
Petroleum Distillate	98.9%

**Equivalent to 8.80% of (butylcarbityl) (8-propylpiperonyl) ether and 0.20% of related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional cautions.

NET CONTENTS:

Manufactured by:
CHEM TECH, LTD.
DES MOINES, IA. 50313



DIRECTIONS (Cont.)

To protect dairy stock from hornflies, houseflies, stableflies, gnats and mosquitoes, spray after milking in the barn using 1 to 2 ounces per head, not to exceed 8 ounces per animal per day. The spray should be applied as a light mist primarily to envelop or hit the flies as they attempt to leave the cows.

For blood-sucking lice on cattle, apply the spray lightly to infested areas, not over 2 ounces per animal, and use a stiff brush on such areas to get the spray to the base of the hair. Treatment for lice should not be repeated more often than once every three weeks.

Space treatment in barns, houses and milk parlors for effective control of flies. Direct mist toward the upper portions of the enclosure, filling the air with fine spray particles, using approximately 2 ounces of material for each 1,000 cubic feet of space. Whenever possible, keep doors and windows closed for 10 minutes after application.

Face fly spray — Apply to face of animal in morning while stanchioned, just before releasing to pasture. Use a spray which produces large wetting droplets. Apply sufficient material to wet the face of the animal, but do not exceed 1.5 ounces per animal. Repeat daily, or as necessary.

NOTICE

Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA REG. NO. 47000-33

EPA EST. NO. 47000-1A-01

1.85