12-07-1990 UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

5285111

DEC 7 1990

Mr. James R. Melton Chem-Tech, Ltd. 4515 Fleur Drive #303 Des Moines, IA 50321

Dear Mr. Melton:

Subject: Revised Label Formula Dy-Fly II Dairy Spray EPA Registration No. 47000-33 Your Submission Dated June 6, 1989

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

As new labels are prepared, delete "ACTIVE INGREDIENTS - 100%" from the ingredient statement. You are correctly declaring 98.4 % inert ingredients in this formulation. The formula statement dated June 5, 1989 is acceptable and has been included in our files for this product.

Sincerely yours,

Platon to

Phil Hutton Product Manager (17) Insecticide-Rodenticide Branch Registration Division (H7505C)

Enclosure

51299:I:A-4:Schroeder:C.Disk:KENCO:09/20/90:CL:ejh

CONCURRENCES									
SYMBOL									
SURNAME						Į			
DATE									
EPA Form 1320	0-1 (12-70)						OFFICI	AL FILE COPY	

CAUTION (12 PT) PRECAUTIONARY STATEMENTS (SPT) HAZARDS TO HUMANS & DOMESTIC ANIMALS (SPT) Harmful if swallowed. Do not induce vomiting. Contact physician. If in eyes flush with water and contact physician. If on skin or clothing, remove clothing. Wash skin with soap and water. Avoid contamination of milk, milk handling equipment and utensils.

ENVIRONMENTAL HAZARDS This product is toxic to fish. Keep out of water. Do not contaminate water when disposing of equipment washwaters.


PHYSICAL AND CHEMICAL HAZARDS Do not use, pour, spill or store near heat or open flame.

STATEMENT OF PRACTICAL TREATMENT IF SWALLOWED:Call a physician or Poison Control Center. Do not induce vomiting. This product contains petroleum solvent Aspiration may be a hazard. Call a physician at once. INHALED: Remove to fresh air. Use artificial respiration if needed. Contact a physician. IN EYES: In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. ON SKIN: Wash with soap and water immediately. Get medical attention if irritation persists.

RED

Blue

DEC 7199





ACTIVE INGREDIENTS	100.0%
Pyrethrins	0.1%
Technical Piperonyl Butoxide**	1.0%
INERTS	98.9%

**Equivalent to 0.80% of (butylcarbityl) (6-propylpiperonyl) ether and 0.20% of related compounds.


KEEP OUT OF REACH OF CHILDREN (SF) CAUTION (2177) See side panel for additional cautions. (S1777)

NET CONTENTS:

MANUFACTURED BY: CHEM-TECH, LTD. 4515 Fleur Dr. Des Moines, IA 50321

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For the protection of Dairy Animals from biting flies and the control of Houseflies in Barns and Milk Houses.

Protects against Hornflies, Stableflies, Houseflies, Gnats, Mosquitoes and Blood Sucking Lice.

A temporary treatment to relieve cattle from annoyance caused by face flies.

DIRECTIONS (Con't.)

To protect dairy stock from hornflies, houseflies, stableflies, gnats and mosquitoes, spray after cattle enter the barn using 1 to 2 ounces per head, not to exceed 2 ounces per animal per day. The spray should be applied as a light mist primarily to envelop or hit the flies as they attempt to leave the cows.

For blood-sucking lice on cattle, apply the spray lightly to infested areas, not over 2 ounces per animal, and use a stiff brush on such areas to get the spray to the base of the hair. Treatment for lice should not be repeated more often than once every three weeks.

Space treatment in barns, houses and milk parlors for effective control of flies. Direct mist toward the upper portions of the enclosure, filling the air with fine spray particles, using approximately 2 ounces of material for each 1,000 cubic feet of space. Whenever possible, keep doors and windows closed for 10 minutes after application.

Face fly spray — Apply to face of animal in morning while stanchioned, just before releasing to pasture. Use a spray which produces large wetting droplets. Apply sufficient material to wet the face of the animal, but do not exceed 1.5 ounces per animal. Repeat daily, or as necessary.



NOTICE

Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons o property arising out of use or handling and accepts the produc on these conditions.

EPA REG, NO, 47000-33

EPA EST. NO. 47000-1A-0

6/89

