

47006-54
 PM 15
 09/12/70
 1/31/70

CAUTION
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid prolonged breathing of spray mist.

If swallowed, consult a physician immediately. Avoid overspraying which will result in wetting the animals' hides. Avoid contamination of milk and milk handling equipment.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

PHYSICAL AND CHEMICAL HAZARDS
DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry, secure area in original container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER: Triple rinse (or equivalent). Then offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other approved state and local procedures.



DAIRY CATTLE
SPRAY

Kills and repels horn flies, stable flies, mosquitoes and gnats. When properly applied, the repellent effect against above insects will last two days.

ACTIVE INGREDIENTS:

| | |
|---|-----------------|
| Pyrethrins | .05% |
| *Piperonyl butoxide, technical | .10% |
| **N-octyl bicycloheptene dicarboximide | .18% |
| DI-n-propyl isocinchomeronate | .20% |
| Dichlorvos (2,2-dichlorovinyl dimethyl phosphate) | .465% |
| Relate Compounds | .035% |
| INERT INGREDIENTS: | 98.990%+ |
| TOTAL | 100.000% |

+Contains petroleum distillate

*Consists of 0.08% (butylcarbityl) (6-propylpiperonyl) ether and 0.02% other related compounds

**MGR 264 Insecticide Synergist U.S. Pat. No. 2,473,512

MGR Repellent 326 U.S. Pat. No. 2,757,120

Equivalent to .50%w of Vapona™ Insecticide "Shell Trademark"

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE ADDITIONAL CAUTIONS ON SIDE PANEL

NET CONTENTS

MANUFACTURED BY:
CHEM-TECH, LTD.
 Des Moines, IA 50313



DIRECTIONS FOR USE

To protect livestock from attack by flies, horn flies, mosquitoes and gnats, apply to pen, particularly cracks and crevices. If adequate control is obtained.

In a complete fly control program in combination with residual insecticide.

To develop a residual repellency to stable flies and gnats on dairy cows, apply 1 to 2 oz. of each animal for a three day period. The spray should be applied to the head, neck, legs, belly and shoulders. Avoid areas where the animal will be milked. Avoid wetting the hide. Spray in the morning and evening. Repeat every second day to disturb the animal. Be sure the teats are dry prior to milking. Calves should be sprayed instead of attempting to build up a residual.

For control of lice on livestock, spray the livestock with Cattle Spray. Immediately rub the spray into the infested areas. Repeat frequently than once every three weeks.

To kill flies, mosquitoes and gnats in the barn, doors and windows and direct the spray should remain closed for 10 to 20 minutes. Repeat as necessary. A spray per 1,000 cubic feet. Do not use on animals have received direct spray with water, feed or foodstuffs, milk or milk products.

NOTICE

Seller's guarantee shall be limited to the product as shown on the label. The buyer assumes any liability for use or handling and accepts the product as shown.

EPA REG. NO. 47000-54

