

FRUIT TREES, VEGETABLES, FLOWERS, AND ORNAMENTALS Apply 0.03% Lindene solution as directed below. Prepare 0.03% solution by adding 1 tablespoonful Lindene Concentrate to 2 1/2 gallons of water or 1 pint to 40 gallons of water.

PEACHES, PLUMS Aphids Leafhoppers Thrips Use 1 gallon of 0.03% spray per 100 square feet. Do not apply within 60 days of harvest.

TREES OR SHRUBS	BORER AND OTHER INSECTS	DOSAGE	TIME TO APPLY
Gr (English and Scotch) Birch Laurel	Leaf Miners	1 tablespoonful per 1 gal. water	Apply when small miners appear; repeat in mid-July. Apply when small miners appear in early July.
Pines	Pine Bark Beetles	1 pint 5 gal. water	Spray trunks thoroughly. Wet trunks to run off.
Pine Spruces	White Pine Weevil	1 Tablespoonful per 1 gal. water	Apply in April when buds begin to swell. Spray upper parts of trees thoroughly.
Taxus	Reddish-brown Bark Beetle Taxus Weevil	1 Tablespoonful per 1 gal. water	Apply in May or early June in the trunk and around base of trees, sealing the bark and wetting the ground surface.
Deciduous Rhododendron-Like	Deciduous Bark Beetle Rhododendron Bark Weevil	1 Tablespoonful per 1 gal. water	Apply in May or early June in the trunk and around base of trees, sealing the bark and wetting the ground surface.

TREES OR SHRUBS	BORER AND OTHER INSECTS	DOSAGE	TIME TO APPLY
Apples Cherries Pearches	Fruit Tree Bark Beetles Rat Hole Saw Borer Round-headed Borer	1 Tablespoonful per 1 gal. water 3 pints 12 gal. water	Apply in late April, June and early July. Apply in June, early ground. Seal bark in trunk. Do not seal in the crown.

Maple, Ash, Sycamore, Elm, etc.	Apple Bark Beetle Round-headed Borer	1 pint 5 gal. water 1 Tablespoonful per 1 gal. water	Apply in late April, June and early July. Apply in June, early ground. Seal bark in trunk. Do not seal in the crown.
Pearches	Pear Tree Bark Beetle	1 pint 5 gal. water	Apply in late April, June and early July. Apply in June, early ground. Seal bark in trunk. Do not seal in the crown.

PECANS To control Pecan Phyloxera and Shothole Curculio mix 29 ounces with 100 gallons of water. Thoroughly wet the foliage at approximately 30 gallons of spray per tree. Do not apply more than 450 gallons of finished spray per acre (1 gal. 6 ozs. gallons of the product). Apply by air blast sprayer when the pecan buds are beginning to burst. This is early to mid April. Make only one application. **DO NOT GRAZE TREATED AREAS**. Do not apply within 210 days of harvest.

TOMATOES Aphids Use 1 gallon of 0.03% spray per 100 square feet. Do not apply after fruit starts to form.
CABBAGE, CELERY, COLLARDS, CUCUMBERS, LETTUCE, OKRA, PEPPERS, SQUASH, MELONS, TOMATOES White Grubs Cutworms Wireworms For preplant soil application only. Use 2 gallons for each 250 square feet. Work into soil thoroughly before or during planting. Do not treat land to be planted in root crops.
BULBS (SNAPDRAGON, GLADIOLUS, HYACINTH, LILIES, LILIES, NARCISSUS, TULIPS) Aphids, Flea Beetles, European corn borer, Iris borer, Thrips. Apply when insects appear. Repeat as necessary.
FLOWERS (CARNATIONS, CHRYSANTHEMUMS, COLEUS, DAPHNES, GARDENIA, GERANIUMS, HIBISCUS, NASTURTIUMS, PANSIES, PETUNIAS, SWEET PEAS, ZINNIA) Aphids, Flea beetles, Leafhoppers, Thrips. Apply when insects appear. Repeat as necessary.
ORNAMENTALS (AZALEAS, CAMELLIAS, HEATHER, LILACS, RHODODENDRONS, ROSES) Aphids, Black vine weevil, Cucumber beetles, Thrips, White Fly. Apply when insects appear. Repeat as necessary.

TICKS AND CHIGGERS ON VEGETATION A minimum dosage of 10 pounds of actual Lindene per acre for control of ticks and chiggers on vegetation. Lawns, recreation areas, camp grounds, golf courses. Mix thoroughly in 25 gallons of water and apply at the rate of 20 to 25 gallons per acre, or about 1 gallon to 1000 square feet. Hand or knapsack type hand sprayers may be used for small areas, but power equipment will be most satisfactory for large acreages. Treat ground under and around parts of plants with this spray.

RECIPE IS LIMITED AVAILABLE