



VMI LIVESTOCK, POULTRY & PREMISE SPRAY

RABON® 50% WETTABLE POWDER INSECTICIDE

| | |
|--|-----------|
| ACTIVE INGREDIENT: | BY WEIGHT |
| 2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate | 50.0% |
| INERT INGREDIENTS | 50.0% |
| Total | 100.0% |

TAKE TIME

OBSERVE LABEL
DIRECTIONS

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid breathing spray mist and dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

See Other Cautions on Side Panel

NET CONTENTS _____

Packed by
ECONOMY PRODUCTS CO., INC.
SHENANDOAH, IOWA 51401

| Insect | Problem Areas | % Solution | Remarks |
|------------|---|------------|--|
| Flies | Dairy barns, poultry houses, swine barns — other animal buildings. | 2.0% | For dry whitewashed wood or concrete block surfaces 1 gal. of solution/500 sq. ft. |
| | | 1.0% | For unpainted wood or painted concrete block surfaces 1 gal. of solution/500 sq. ft. |
| | | 1.0% | For masonite or galvanized sheet metal surfaces ½ gal. of solution/500 sq. ft. |
| Meat flies | Poultry droppings, manure piles, garbage pites, under feed troughs. | 1.0% | Apply 1 gal. of mixed solution 100 sq. ft. Penetrate problem area first time—repeat every 7-10 days thereafter. |
| Ticks | Campgrounds, backyards, picnic areas, recreational parks, other outdoor living areas. | 0.5% | Spray infested areas thoroughly (approximately 25 gals. of mixed solution/acre). Spray along foot paths and roadsides leading to such areas. |

ACCEPTED
 FEB 13 1975
 ORDER NO. _____
 PURCHASE AND RETURN
 FOR ECONOMIC TOILET
 TO ATTACHED COMMENTS.

DILUTION TABLE

| Amount of Rabon® Wettable Powder | AMOUNT OF WATER FOR | | | |
|-------------------------------------|---------------------|------------------|------------------|------------------|
| | 0.33% Solution | 0.5% Solution | 1.0% Solution | 2.0% Solution |
| ¼ lb. | 9 Gal. | 6 Gal. | 3 Gal. | 1½ Gal. |
| ½ lb. | 36 Gal. | 25 Gal. | 12½ Gal. | 6 Gal. |
| 4 lb. | 75 Gal. | 50 Gal. | 25 Gal. | 12½ Gal. |
| 8 lb. | 150 Gal. | 100 Gal. | 50 Gal. | 25 Gal. |

PRECAUTIONS IN USING

ENVIRONMENTAL CAUTIONS: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DISPOSAL PRECAUTIONS: Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg. No. 3770-311
EPA Est. 3770-NB-1

3653B-PBEP0

Page 1 of 2

| Animal | Insect | % Solution | Remarks |
|-------------|------------------|------------|---|
| Beef Cattle | Horn Flies, Lice | 0.35% | Apply as coarse spray. Use between ½ and 1 gallon of spray per animal depending on size and hair coat. There is no withholding period from last application to slaughter. |
| | Long Star Ticks | 0.5% | |
| Swine | Lice | 0.5% | Apply as a coarse spray using 1 to 2 quarts per head to thoroughly wet the animal. Repeat in 2 weeks if necessary. |

POULTRY USE DIRECTIONS

| Insect | Type of Housing | % Solution | Remarks |
|---|------------------------------|------------|---|
| Lice & Mites | Wire Cages | 0.5% | Apply directly to birds (1 gal./100 birds). spray vent and fluff areas from below. Repeat when necessary. Do not repeat more often than every 14 days. For individual bird treatment apply 1 ounce of the mixture per bird. |
| <p>NOTE: For maximum lasting control of the northern fowl mite, penetration of the feathers around the vent area is absolutely essential. Use power sprayer at 100-125 psi at NO LESS THAN RECOMMENDED RATE. More attention must be given each individual bird when using low-pressure equipment. Treat roosters carefully and thoroughly to avoid reinfestation in breeding flocks.</p> | | | |
| | Floor Management Litter | 0.5% | Apply 1-2 Gal./1,000 sq. ft. evenly for penetration to litter surfaces. Also apply thoroughly to walls, roosts, cracks, crevices and interiors. Spray birds lightly. |
| | Floor Management Litter | 50% WP | Treat evenly and thoroughly using 2.5 oz./100 sq. ft. |
| | Floor Management Dust Box | 50% WP | Mix evenly throughout top layer of box contents using 2.5 oz./50 birds. |
| | Floor Management Roost Paint | 1.0% | Treat with brush or spray thoroughly, particularly cracks and crevices using 1 pt./100 ft. |
| Fowl Tick (Blue Bug) | <u>All Types</u> | 1.0% | Apply 1 gal./100-150 sq. ft. thoroughly to walls, ceiling, floor cracks and crevices with a power sprayer. |