

**CAUTION
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS &
DOMESTIC ANIMALS**

Harmful if swallowed or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply over uncovered milk or food products. Do not use in areas where fish or reptiles may be exposed to the spray.

When using this product as a space spray in food processing plants, foods must be removed or covered during treatment. Apply only when the facility is not in operation. At home and in food processing establishments, cover or remove food prior to treatment. Cover food contact surfaces, benches, utensils, equipment, etc., during treatment or thoroughly clean prior to use with an effective cleaning compound followed by a potable water rinse.

In hospitals or other medical facilities, remove patients prior to treatment. Ventilate treated area for two hours prior to returning patients.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

PHYSICAL OR CHEMICAL HAZARDS

Extremely flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F. may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with the label.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry, secure area in original container.

DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER: Do not reuse. Do not incinerate or puncture. Wrap empty container and put in trash collection.

Hold container upright, nozzle away from you and press actuator button spraying as directed.

To Treat Animals: To protect cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

In Animal Quarters & Milk Rooms: For rapid control of houseflies, mosquitoes, gnats, wasps and flying moths close windows and doors and spray at the rate of one second per 1,000 cubic feet of space. For best results keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is



**DY-FLY
DAIRY
AEROSOL
PRESSURIZED**

KILLS & REPELS

House Flies, Horn Flies, Stable Flies, Horse Flies, Face Flies, Deer Flies, Gnats, Mosquitoes, Small Flying Moths (millers), Roaches, Waterbugs, Sowbugs, Spiders, Centipedes, Ants, Crickets, Wasps and Hornets.

For use in Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food Handling Plants, Dairies, Milk House, Cattle, Horse, Poultry, and Swine Quarters. For use on cattle, horses and swine.

**FOR USE ON CATTLE, IN ANIMAL
QUARTERS, AND MILK ROOMS**

**Non-Aqueous, Low Oil Pyrethrin +
Piperonyl Butoxide Insecticide**

ACTIVE INGREDIENTS:

Pyrethrins 0.5%
*Piperonyl Butoxide 5.0%

INERT INGREDIENTS: 94.5%

*Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and 1.0% related compound.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

Contents under pressure
See additional precautions on side panel.

NET WEIGHT:

Distributed by:
CHEM-TECH, LTD.
DES MOINES, IA 50321



ACCEPTED
18 FEB 1992
Under the Federal Insecticide,
Fungicide, and Plant Growth Regu-
lators Act, for the pesticide
No. 47000-69

HOME AND FOOD PROCESSING AREA USE: For flies, mosquitoes, small flying moths (millers), and gnats: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated areas. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed. **Roaches, Waterbugs, Sowbugs, Spiders and Centipedes:** spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary. **ANTS:** spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

STORED PRODUCT INSECTS (accessible stages): Flying Mediterranean flour moths, Indian meal (chocolate) moths, angoumois grain moths: Close doors and windows. Direct aerosol mist to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be resting. Use a minimum treatment of six to eight seconds per 1000 cubic feet of space. Keep area closed for 30 minutes after each treatment. Repeat treatment weekly or as flying moths appear. Infested stored food products should be fumigated or treated with an approved product.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: For poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN AND STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. In animal application, lightly mist over backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horses face or eyes. Repeat application daily or as necessary.

NOTICE TO USER: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the products on these conditions.

EPA Reg No. 47000-69
EPA Est No. 47000-IA-01

47000-69

02/15/1992

111