

47000-72

11/04/2003

Page 1 of 3

ACCEPTED

NOV 4 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 47000-72



CT-75
Aerosol Insecticide

Especially Formulated For:

Dairy Farms (including Milk House, Milking Parlor, Loafing Sheds and Holding Lot), Poultry and Hog Operations, Horse Barns, and Stables, Beef Cattle Operations, Kennels, Catteries, Food Storage Areas, Meat Packing and Food Processing Plants, Homes, Hotels, Stores and Restaurants.

KILLS: House Flies, Horn Flies, Stable Flies, Gnats, Mosquitoes, Wasps, Small Flying Moths, Fruit Flies, Adult Fleas, Grain Weevils, Flour Beetles, Chocolate Moths, Carpet Beetles, Drug Store Beetles, Roaches (Waterbugs), Spiders, Ants, Silverfish, and Crickets.

ACTIVE INGREDIENTS:

Pyrethrins.....	0.7%
Technical Piperonyl butoxide*.....	5.0%
INERT INGREDIENTS**.....	94.3%
TOTAL	100.00%

*Equivalent to 4.0% (butylcarbityl)(6-propylpiperonyl) ether and 1.0% related compounds.

**Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN

WARNING

FIRST AID

- If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advise.
- If Swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
- If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise.
- If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advise.

For emergency information contact: National Pesticide Information Center at 1-800-858-7378 between 6:30 am and 4:30 Pacific Time. After 4:30 call your poison control center.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

Harmful if swallowed or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. When using this product as a space spray in food areas, do not apply while food processing is under way. Except in federally inspected meat and poultry plants, careful crack and crevice application for inspection purposes is permissible at all times. After space spraying in meat packing plants, bakeries and other food processing plants wash with an appropriate cleaning agent and rinse with potable water where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and birds. Do not apply directly to water.

PHYSICAL OR CHEMICAL HAZARDS

Extremely Flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130 F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TV's, computers, etc.

USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry place away from children, preferably a secure storage area.

CONTAINER DISPOSAL: Do not puncture or incinerate! If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

AGRICULTURAL USES

For control of House Flies, Horn Flies, Stable Flies, Gnats, Mosquitoes, Small Flying Moths.

Dairy Farm Use: MILKING PARLOR AND MILK ROOM—Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed tight, double the dosage. Stand in middle of room. Direct fog upward and spray in all directions. Seconds may be counted saying one-thousand, two-thousand, etc. Keep room closed for 15 minutes after treatment. Example: a room 20 feet long, 12 feet wide, and 8 feet high requires 2 seconds if closed tight, 4 seconds if not. Do not remain in treated areas. Ventilate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

STANCHION BARN USE—Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

ANIMAL USE—From approximately 2 foot distance, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

POULTRY OPERATIONS: Use only in empty poultry houses (when birds are not present). Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, Close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. For animal application, direct the spray over backs and spray 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN AND STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. In animal application, lightly mist over backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horse's face or eyes. Repeat application daily or as necessary.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. For animal use, thoroughly spray entire animal as it is being released into pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

NON-AGRICULTURAL USES

KENNELS AND CATTERIES: Remove animals before spraying. Ventilate after treatment is completed.

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

ADULT FLEAS in Buildings (Not on Animals). Close windows and doors. Hold aerosol 36" above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet, making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatments after 6 days.

FOOD STORAGE AREAS, MEAT AND FOOD PROCESSING PLANTS

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths, Fruit Flies. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, CARPET BEETLES, DRUG STORE BEETLES in exposed adult larvae stages: Contact as many insects as possible when treating infested areas such as cracks, crevices, stored products, machinery, and dark secluded areas. For Carpet Beetles, treat infested area of carpets, localized areas of floors, baseboards, and into cracks and under carpets where these pests may be found. Spray 10 to 20 seconds per 1000 square feet of floor space. Repeat treatment weekly or as insects appear.

ROACHES, SPIDERS, ANTS, SILVERFISH, CRICKETS: Hold aerosol close to cracks and mist directly into them, contacting as many insects as possible. Apply freely in all possible hiding places behind baseboards, sinks, cabinets, coffee makers, meter boxes, manholes. Repeat as necessary.

NOTICE

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Chem-Tech, Ltd, the manufacturer or seller. In no case shall Chem-Tech, Ltd, the manufacturer or seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer.

Except as expressly provided herein, Chem-Tech, Ltd, the manufacturer or seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. Buyer's or user's exclusive remedy, and Chem-Tech, Ltd's, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.

EPA Reg. No. 47000-72

EPA Est. No. 47000-IA-01

NET WEIGHT: _____

MFG. By:
Chem-Tech, Ltd.
Des Moines, IA 50321