

<p>U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460</p> <p>NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION</p> <p><i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i></p>	<p>EPA REGISTRATION 47000-73</p> <p>TERM OF ISSUANCE</p> <p>NAME OF PESTICIDE PRODUCT GT 511</p>	<p>DATE OF ISSUANCE FEB 11 1986</p>
<p>NAME AND ADDRESS OF REGISTRANT (Include ZIP code)</p> <p><input type="checkbox"/> <input type="checkbox"/></p> <p>Chen-Tech, Ltd. 4515 Elm Drive #303 Des Moines, IA 50321</p> <p><input type="checkbox"/> <input type="checkbox"/></p>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p>		
<p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p>		
<p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p>This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:</p>		
<ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data. 2. Revise the language of the side panel precautionary statement by deleting the current paragraph in the draft labeling beginning "when using this product . . ." and replace it with the following: "When using this product as a space spray in food processing plants, foods must be removed or covered during treatment. Apply only when the facility is not in operation. All food-processing surfaces must be thoroughly cleaned with an appropriate cleaning agent and rinsed with potable water before reuse. 3. Add the phrase "EPA Registration No. 47000-73" to your label before you release the product for shipment. 		
<p><input type="checkbox"/> ATTACHMENT IS APPLICABLE</p>		
<p>SIGNATURE OF APPROVING OFFICIAL</p>	<p>DATE</p>	

4. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure



511

FOR
USE IN

"Only for sale to, use, and
storage by service persons."

Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restuarants,
schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food Handling
Plants, Dairies, Milk House, Cattle, Horse, Poultry, and Swine quarters. For
use on cattle, horses and swine.

with
DUAL ACTIVATORS
Kills & Repels

House Flies, Horn Flies, Stable Flies, Horse Flies, Face Flies, Deer Flies,
Gnats, Mosquitos, Small Flying Moths (millers), Roaches, Waterbugs, Sowbugs,
Spiders, Centipedes, Ants, Crickets, Wasps and Hornets.

ACTIVE INGREDIENTS:

Pyrethrins-----0.5
Piperonyl Butoxide, Tech*-----1.0
N-Octyl Bycycloheptene
Dicarboximide-----1.0
Refined Petroleum Oil-----2.0

INERT INGREDIENTS-----95.5

*Equivalent to 0.8%
(Butlycarbityl)(6-Propylpiperonyl)
ether and 0.2% of Related Compounds.

KEEP OUT OF REACH OF CHILDREN (8 Pt. Type)
WARNING (12 Pt Type)

Contents Under Pressure

See Side Panels for additional Precautions

Net Wt: 2 lbs.

EPA REG. NO. 47000-

EPA EST. NO 47000-1A-01

MFG. BY

CHEM-TECH, LTD.
Des Moines, IA 50321

ACCEPTED
with COMMENTS
in EPA Letter Dated:

FEB 11 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

47000-83

SIDE PANEL

PRECAUTIONARY STATEMENTS (8 PT)
HAZARDS TO HUMANS & DOMESTIC ANIMALS (8 PT)

WARNING (12 Pt. Type)

Harmful if swallowed. Avoid contact with skin and eyes. Avoid inhalation of spray mist. Avoid contamination of feed and foodstuffs.

When using this product as a space spray in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation.

Do not remain in treated areas and ventilate the area after treatment is completed.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS:

Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL (10 Pt. Type)

STORAGE: Store in a cool, dry, secure area.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

Remove protective cap. Aim spray opening away from person. Push button to spray.

HOME AND FOOD PROCESSING AREA USE:

For flies, mosquitoes, small flying moths (millers), and gnats: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated areas. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

Roaches, Waterbugs, Sowbugs, Spiders and Centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary. ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

STORED PRODUCT INSECTS (accessible stages):

Flying Mediterranean flour moths, Indian meal (chocolate) moths, angoumois grain moths: Close doors and windows. Direct aerosol mist to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be resting. Use a minimum treatment of six to eight seconds per 1000 cubic feet of space. Keep area closed for 30 minutes after each treatment. Repeat treatment weekly or as flying moths appear. Infested stored food products should be fumigated or treated with an approved product.

DAIRY FARM USE: Milking parlor and milk room. Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed, double the dosage. Stand in middle of room. Direct spray in all directions. Keep room closed for 15 minutes after treatment. Do not remain in treated area. Ventilate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

SIDE PANEL

PRECAUTIONARY STATEMENTS (8 PT)
HAZARDS TO HUMANS & DOMESTIC ANIMALS (8 PT)

WARNING (12 Pt. Type)

Harmful if swallowed. Avoid contact with skin and eyes. Avoid inhalation of spray mist. Avoid contamination of feed and foodstuffs.

When using this product as a space spray in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation.

Do not remain in treated areas and ventilate the area after treatment is completed.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS:

Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL (10 Pt. Type)

STORAGE: Store in a cool, dry, secure area.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

Remove protective cap. Aim spray opening away from person. Push button to spray.

HOME AND FOOD PROCESSING AREA USE:

For flies, mosquitoes, small flying moths (millers), and gnats: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated areas. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

Roaches, Waterbugs, Sowbugs, Spiders and Centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary. ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

STORED PRODUCT INSECTS (accessible stages):

Flying Mediterranean flour moths, Indian meal (chocolate) moths, angoumois grain moths: Close doors and windows. Direct aerosol mist to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be resting. Use a minimum treatment of six to eight seconds per 1000 cubic feet of space. Keep area closed for 30 minutes after each treatment. Repeat treatment weekly or as flying moths appear. Infested stored food products should be fumigated or treated with an approved product.

DAIRY FARM USE: Milking parlor and milk room. Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed, double the dosage. Stand in middle of room. Direct spray in all directions. Keep room closed for 15 minutes after treatment. Do not remain in treated area. Ventilate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

SIDE PANEL

DIRECTIONS FOR USE. (CON'T)

DAIRY ANIMAL USE: From approximately 2 foot distance thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN AND STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. In animal application, lightly mist over backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horses face or eyes. Repeat application daily or as necessary.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

NOTICE TO USER: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

DOT 39 NRC 13C/180 M1089

Federal law forbids transportation if refilled - penalty up to \$25,000.00 and 5 years imprisonment (49 U.S.C. 1809)