

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid inhalation of spray mist. Avoid contamination of feed and foodstuffs.

When using this product as a space spray in food processing plants, foods must be removed or covered during treatment. Apply only when the facility is not in operation. At home and in food processing establishments, cover or remove food prior to treatment. Cover food contact surfaces, benches, utensils, equipment, etc., during treatment or thoroughly clean prior to use with an effective cleaning compound followed by a potable water rinse.

In hospitals or other medical facilities, remove patients prior to treatment. Ventilate treated area for two hours prior to returning patients.

Do not remain in treated areas and ventilate the area after treatment is completed.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS: Extremely flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or around electrical equipment due to the possibility of shock hazard.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, birds, and other wildlife. Keep out of lakes, ponds, or streams. Do not contaminate water when disposing of equipment washwater. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry, secure area.
DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

Remove protective cap. Aim spray opening away from person. Push button to spray.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: For poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. Treat daily or as necessary.

EPA Reg No. 47000-73 EPA Est No. 47000-1A-01



CT-511

Aerosol Insecticide

PRESSURIZED

DUAL SYNERGISTS -- KILLS & REPELS

House Flies, Horn Flies, Stable Flies, Horse Flies, Face Flies, Deer Flies, Gnats, Mosquitoes, Small Flying Moths (millers), Roaches, Waterbugs, Sowbugs, Spiders, Centipedes, Ants, Crickets, Wasps and Hornets.

For Use In Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food Handling Plants, Dairies, Milk House, Cattle, Horse, Poultry, and Swine Quarters. For use on cattle, horses and swine.

ACTIVE INGREDIENTS:

- Pyrethrins..... 0.5%
- Piperonyl Butoxide, Technical..... 1.0%
- N-Octyl Bicycloheptene Dicarboximide..... 1.0%
- **INERT INGREDIENTS..... 97.5%

*Equivalent to 0.8% (Butylcarbitolyl 16 Propylpiperonyl) ether and 0.2% related compounds.

**Contains Petroleum Distillates

Statement of Practical Treatment

If Swallowed: Call a physician or poison control center immediately. Do not induce vomiting. Vomiting may cause aspiration pneumonia.
If inhaled: Remove patient to fresh air. Apply artificial respiration if indicated. **If on Skin:** Remove contaminated clothing and wash affected skin areas with soap and water. Get medical attention if irritation persists. **If In Eyes:** Flush eyes with plenty of water. Get medical attention if irritation persists.

NET WEIGHT:

DIRECTIONS FOR USE Cont

HOME AND FOOD PROCESSING AREA USE: For flies, mosquitoes, small flying moths (millers), and gnats: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated areas. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed. **Roaches, Waterbugs, Sowbugs, Spiders and Centipedes:** spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary. **ANTS:** spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

STORED PRODUCT INSECTS (accessible stages): Flying Mediterranean flour moths, Indian meal (chocolate) moths, angoumois grain moths: Close doors and windows. Direct aerosol mist to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be resting. Use a minimum treatment of six to eight seconds per 1000 cubic feet of space. Keep area closed for 30 minutes after each treatment. Repeat treatment weekly or as flying moths appear. Infested stored food products should be fumigated or treated with an approved product.

DAIRY FARM USE: Milking parlor and milk room. Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed, double the dosage. Stand in middle of room. Direct spray in all directions. Keep room closed for 15 minutes after treatment. Do not remain in treated area. Ventilate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

DAIRY ANIMAL USE: From approximately 2 foot distance thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

HOG OPERATIONS: In hog houses, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. For animal application direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN AND STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. Lightly mist over backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horse's face or eyes. Repeat application daily or as necessary.

NOTICE TO USER: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions. 2-01



Mfg. By:
CHEM-TECH, LTD.
Des Moines, IA 50321



0 45446 00145 7

47000-73

ACCEPTED
APR 23 2001
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pseudo registration under 47000-73, Reg. No.

04/23/2001

1/1