

272

Check amount of liquid in insecticide tank or auxiliary tank before and after treatment to assure proper dosage has been applied. Return unused insecticide to original container and clean machine outdoors using an approved solvent before returning to truck or storage area. This will help preserve equipment and prevent cross-contamination.

Treatment should be made during the hours when buildings are unoccupied. Buildings should be closed and ventilation kept at a minimum during application of the insecticide. Lock all entrances and do not allow unprotected workers to enter buildings when being treated.

Place danger signs at all entrances stating:

"This building should be opened and aired before re-entry of unprotected workers, or remain unoccupied by unprotected workers for at least 6 hours after treatment. Do not enter prior to \_\_\_\_\_ o'clock \_\_\_\_\_ date without protection."

Repeat treatment at weekly intervals during season of peak insect activity for optimum results.

Exposed food should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before processing is resumed. When using the product in these areas, apply only when the facility is not in operation.

**SPACE SPRAY AND CONTACT SPRAY** — Dilute to 1-3% with an approved solvent and apply at the rate of 1 gallon of spray per 64,000 cu. ft. to kill roaches, silverfish, fleas, waterbugs, earwigs, ants, spiders, in sheltered situation. Adjust a ULV spraying device to a course spray for control of the above listed crawling insects. Contact as many of the insects with the spray as possible. Thoroughly spray all parts of the room suspected of harboring these insects. Special attention should be paid to cracks, hidden surfaces under equipment, and food storage areas. Then adjust the ULV device to deliver a fine spray for use as a space spray. Sweep up and destroy dead insects. Repeat treatment as necessary.

**FOR USE AS A CONTACT SPRAY** — Dilute to 1% and spray 1 pint per 10,000 cubic feet or 1,000 square feet of floor space.

Do not use in the edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

#### GENERAL OPERATING INSTRUCTIONS

(For specific operating instructions, refer to  
Manufacturer's Instruction Booklet furnished with machine)

1. Having calculated amount of time required for treatment and flow rate required, place machine at farthest point from intended exit.
2. Direct nozzle(s) upward and pointed down aisle, not directly toward any nearby object.
3. Start machine and adjust flow to desired rate. Calibrate if necessary. Refer to operating manual.
4. Operate for calculated time period. Begin by backing along precalculated path at predetermined speed.
5. When treatment time has been completed, stop insecticide flow by closing valve. Wait until spray stops, stop machine and remove from area.
6. Wait at least 6 hours after completion of treatment before ventilating. 12 or 16 hours exposure may be used, if convenient or preferred, in which case ventilation is not necessary.

#### WARRANTY

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other WARRANTY of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.



DICHLORVOS MIXTURE

NA2783 RQ