

PM 17

47000-84

17/1/2

DEC 21 1990

3833/2307  
17/1


Mr. James R. Melton  
Chem-Tech, Ltd.  
4515 Fleur Drive, #303  
Des Moines, IA 50321

Dear Mr. Melton:

Subject: CT-25  
EPA Registration No. 47000-84  
Your Submission Dated May 21, 1990

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,



Phil Hutton  
Product Manager (17)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

57656:1:A-4:Nelson:C.Disk:KENCO:10/5/90:EK:VO:EK

CONCURRENCES

SYMBOL	SURNAME	DATE						

ACCEPTED

DEC 21 1990

47000-54

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid inhalation of spray mist. Avoid contamination of feed and foodstuffs.

When using this product as a space spray in food processing plants, foods must be removed or covered during treatment. Apply only when the facility is not in operation. All food processing surfaces must be thoroughly cleaned with an appropriate cleaning agent and rinsed with potable water before reuse.

Do not remain in treated areas and ventilate the area after treatment is completed.

Remove pets, birds and cover fish aquariums before spraying.

**PHYSICAL OR CHEMICAL HAZARDS:** Extremely Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry, secure area.

**DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

Remove protective cap. Aim spray opening away from person. Push button to spray.

**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

**POULTRY OPERATIONS:** For poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.



**CT-25**

**PRESSURIZED**

For use in Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food Handling Plants, Dairies, Milk House, Cattle, Horse, Poultry, and Swine Quarters. For use on cattle, horses and swine.

**with DUAL ACTIVATORS  
KILLS & REPELS**

House Flies, Horn Flies, Stable Flies, Horse Flies, Face Flies, Deer Flies, Gnats, Mosquitos, Small Flying Moths (millers), Roaches, Waterbugs, Sowbugs, Spiders, Centipedes, Ants, Crickets, Wasps and Hornets.

**ACTIVE INGREDIENTS:**

Pyrethrins . . . . . 0.25%  
Piperonyl Buloxide, Tech\* . . . . . 1.0%  
N-Octyl Jycycloheptene Dicarboximide . . . . . 1.0%

**INERT INGREDIENTS . . . . . 97.75%**

\*Equivalent to 0.8% (Butylcarbyl) (6-Propylpiperonyl) ether and 0.2% of Related Compounds.


**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Contents Under Pressure

See Side Panels for Additional Precautions

**NET WEIGHT: 1.5 lbs.**

Mfg By:  
**CHEM-TECH, LTD.**  
4515 Fleur Dr., #303  
Des Moines, IA 50321



DI

**HOME AND FOOD PLANTS:** For small flying moths (millers), Point nozzle upward, especially windows and doors. Fill the room with spray for at least 15 minutes. Repeat as necessary. **Roaches and Centipedes:** spray thoroughly in moist areas, opening baseboards and storerooms where possible. Repeat as necessary. **Ants:** Spray on ant entry points.

**STORED PRODUCT:** For grain moths: Close doors and windows of the room, especially product containers with airtight treatment of six to eight weeks or as flying insects should be fumigated.

**DAIRY FARM USE:** In rooms cannot be closed. Direct spray in all directions. Do not re-enter until treatment is complete. Avoid contamination from spray.

**DAIRY ANIMAL USE:** Spray entire animal directly toward animal's face or eyes where saliva accumulates.

**STANCHION BARN:** Spray backs allowing 1 second each morning.

**HOG OPERATIONS:** Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Do not spray directly towards animal's face or eyes.

**HORSE BARN:** Spray entire animal directly toward animal's face or eyes. Repeat application daily or as necessary.

**NOTICE TO USER:** Read entire label, and see side panels for proper use of the products on the label.

**EPA Reg. No. 4700-54-0101**  
5/90

Only for sale to, use, and storage by service persons.