

MAY 31 1991

Mr. James R. Melton
Chem-Tech, LTD.
4515 Fleur Drive #303
Des Moines, IA 50321

Dear Mr. Melton:

Subject: Formula revision
CT-25
EPA Registration Number 47000-84
Submissions dated May 8, 1991 and November 30, 1990


The amendment cited above, adding the phrase "Extremely flammable" to the product label, is acceptable.

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comment listed below. A stamped copy is enclosed for your records.

Add the following statement to the label:

In hospitals or other medical facilities, remove patients prior to treatment. Ventilate treated area for two hours prior to returning patients.

Sincerely,


Richard F. Mountfort
Product Manager 10
Insecticide-Rodenticide Branch
Registration Division (H7505C)

292

ACCEPTED
With COMMENTS
In EPA Letter Dated
MAY 31 1991
under the Federal Insecticide,
Fungicide, and Rodenticide Act
registered for the pesticide
under EPA Reg. No.

F T E R

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid inhalation of spray mist. Avoid contamination of feed and foodstuffs.

When using this product as a space spray in food processing plants, foods must be removed or covered during treatment. Apply only when the facility is not in operation. All food processing surfaces must be thoroughly cleaned with an appropriate cleaning agent and rinsed with potable water before reuse.

Do not remain in treated areas and ventilate the area after treatment is completed.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS: Extremely Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry, secure area.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

Remove protective cap. Aim spray opening away from person. Push button to spray.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: For poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.



CT-25

PRESSURIZED

For use in Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food Handling Plants, Dairies, Milk House, Cattle, Horse, Poultry, and Swine Quarters. For use on cattle, horses and swine.

**with DUAL ACTIVATORS
KILLS & REPELS**

House Flies, Horn Flies, Stable Flies, Horse Flies, Face Flies, Deer Flies, Gnats, Mosquitos, Small Flying Moths (millers), Roaches, Waterbugs, Sowbugs, Spiders, Centipedes, Ants, Crickets, Wasps and Hornets.

ACTIVE INGREDIENTS:

Pyrethrins	0.25%
Piperonyl Butoxide, Tech*	1.0%
N-Octyl Bicycloheptene Dicarboximide	1.0%
INERT INGREDIENTS	97.75%

*Equivalent to 0.9% (Butylcarbitol) (6-Propylpiperonyl) ether and 0.2% of Related Compounds.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
Contents Under Pressure
See Side Panels for Additional Precautions

NET WEIGHT: 1.5 lbs.

Mfg. By:
CHEM-TECH, LTD.
4515 Fleur Dr., #303
Des Moines, IA 50321



HOME AND FOOD PLANTS: Small flying moths (millers). Point nozzle upward, especially windows and doors. Fill the room with product. Close room closed for at least 15 minutes. Repeat as necessary. Reach pedes: spray thoroughly moist areas, opening baseboards and skirting possible. Repeat as necessary. Spray on ants.

STORED PRODUCT: Mediterranean flour moth, grain moths: Close doors of the room, especially product containers with treatment of six to eight area closed for 30 minutes weekly or as flying insects should be fumigated.

DAIRY FARM USE: Milk and doors. Spray at 1000 cubic feet. Direct spray in all directions. Do not re-treatment is complete. Contamination from insects.

DAIRY ANIMAL USE: spray entire animal directly toward animal where saliva accumulates.

STANCHION BARN: backs allowing 1 second each morning.

HOG OPERATIONS: Apply at the rate of 1 to 2 seconds per 1000 cubic feet. double the dosage. and spray for 1 to 2 seconds animal's face or eyes.

HORSE BARN: ANIMALS: windows and doors. feet. Keep area closed. Do not spray directly daily or as necessary.

NOTICE TO USER: of the label, and persons or property the products on the EPA Reg. No. 470 5/90