

PM03

47000-93

2/25/99

1085



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

FEB 25 1999

47000-93

Term of Issuance:

Conditional

Name of Pesticide Product:

CT 55 Dairy Aerosol

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Chem-Tech Ltd.  
4515 Fleur Drive #303  
Des Moines, IA 50321

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 47000-93".
3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

*Marion Johnson* /PM/IB/KD

Date:

FEB 25 1999

(FRONT PANEL)

CT 55

DAIRY AEROSOL

PRESSURIZED

KILLS & REPELS

House Flies, Horn Flies, Stable Flies, Horse Flies, Face Flies, Deer Flies, Gnats, Mosquitoes, Small Flying Moths (millers), Roaches, Waterbugs, Sowbugs, Spiders, Centipedes, Ants, Crickets, Wasps and Hornets.

For use in Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food Handling Plants, Dairies, Milk Houses; Cattle, Horse, Poultry, and Swine Quarters. For use on cattle, horses and swine.

ACTIVE INGREDIENTS:

Pyrethrins .....	0.5%
Technical Piperonyl Butoxide* .....	5.0%

INERT INGREDIENTS: .....	94.5%
TOTAL .....	100.0%

\*Equivalent to 4.0% (butylcarbityl)(6-propylpiperonyl)ether and 1.00% related compounds.

KEEP OUT OF REACH OF CHILDREN


CAUTION

Contents Under Pressure  
See Side Panels for Additional Precautions

WARNING: Contains methyl chloroform a substance which harms public health and environment by destroying ozone in the upper atmosphere.

NET WEIGHT:

Distributed by:  
CHEM-TECH LTD.  
DES MOINES, IA 50321



Quality Seal

3 of 5

(LEFT SIDE PANEL)

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS &  
DOMESTIC ANIMALS  
CAUTION

Harmful if swallowed or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

When using this product as a space spray in food processing plants, foods must be removed or covered during treatment. Apply only when the facility is not in operation. At home and in food processing establishments, cover or remove food prior to treatment. Cover food contact surfaces, benches, utensils, equipment, etc., during treatment or thoroughly clean prior to use with an effective cleaning compound followed by a potable water rinse.

In hospitals or other medical facilities, remove patients prior to treatment. Ventilate treated area for two hours prior to returning patients.

PHYSICAL OR CHEMICAL HAZARDS

Extremely Flammable. Contents under pressure. Keep away from fire, spark and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TV's, computers, etc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry, secure area in original container.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: DO NOT REUSE. DO NOT INCINERATE. Wrap empty container and put in trash collection.

Hold container upright, nozzle away from you and press actuator button spraying as directed.

4 of 5

**TO TREAT ANIMALS:** To protect cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

**IN ANIMAL QUARTERS & MILK ROOMS:** For rapid control of house flies, mosquitoes, gnats, wasps, and flying moths close windows and doors and spray at the rate of one second per 1000 cubic feet of space. For best results keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

(RIGHT SIDE PANEL)

**AREA AND FOOD PROCESSING AREA USE:** For Flies, Mosquitoes, Small Flying Moths (millers), and Gnats: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

**Roaches, Waterbugs, Sowbugs, Spiders and Centipedes:** spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary.

**ANTS:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**STORED PRODUCT INSECTS (accessible stages):** Flying Mediterranean flour moths, Indian meal (chocolate) moths, Angoumois grain moths: Close doors and windows. Direct aerosol mist to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be resting. Use a minimum treatment of 6 to 8 seconds per 1000 cubic feet of space. Keep area closed for 30 minutes after each treatment. Repeat treatment weekly or as flying moths appear. Infested stored food products should be fumigated or treated with an approved product.

**BEEF CATTLE OPERATIONS:** In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

**POULTRY OPERATIONS:** For poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

**HOG OPERATIONS:** In hog houses, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

5 of 5

HORSE BARN AND STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. In animal application, lightly mist over backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horses face of eyes. Repeat application daily or as necessary.

NOTICE TO USER: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use of handling and accepts the products on these conditions.

EPA Reg. No. 47000-69  
EPA Est. No. 47000-1A-01

