

47000-94

1-23-2001

1/4

-- CT LOGO --

Premium Dairy Aerosol

Especially Formulated For:

Dairy Farms (including Milk House, Milking Parlor, Loafing Sheds and Holding Lot). Poultry and Hog Operations, Horse Barns and Stables, Beef Cattle Operations, Kennels, Catteries, Food Storage Areas, Meat Packing and Food Processing Plants, Homes, Hotels, Stores and Restaurants.

KILLS: House Flies, Horn Flies, Stable Flies, Gnats, Mosquitoes, Wasps, Small Flying Moths, Fruit Flies, Adult Fleas, Grain Weevils, Flour Beetles, Chocolate Moths, Carpet Beetles, Drug Store Beetles, Roaches (Waterbugs), Spiders, Ants, Silverfish, and Crickets.

ACTIVE INGREDIENTS:

Pyrethrins	0.7%
Technical Piperonyl Butoxide*	5.0%
INERT INGREDIENTS**:	94.3%
TOTAL	100.0%


*Equivalent to 4.00% (butylcarbityl) (6-Propylpiperonyl)ether and 1.0% related compounds

**Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

WARNING

Contents Under Pressure
See Side Panels for Additional Precautions



NET WEIGHT: 2 lbs.

Mfg. By:
CHEM-TECH, LTD.
Des Moines, IA 50321

Quality
Seal

(Left Side Panel)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. When using this product as a space spray in food areas, do not apply while food processing is under way. Except in federally inspected meat and poultry plants, careful crack and crevice application for inspection purposes is permissible at all

times. After space spraying in meat packing plants, bakeries and other food processing plants wash with an appropriate cleaning agent and rinse with potable water where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish and birds. Do not apply directly to water.

PHYSICAL OR CHEMICAL HAZARDS:

Extremely flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry, area away from heat or open flame.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Replace cap and discard empty container in trash, sanitary landfill, or by other approved state and local procedures. Do not incinerate or puncture.

AGRICULTURAL USES

For control of House Flies, Horn Flies, Stable Flies, Gnats, Mosquitoes, Small Flying Moths.

Dairy Farm Use: MILKING PARLOR AND MILK ROOM--Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed tight, double the dosage. Stand in middle of room. Direct fog upward and spray in all directions. Seconds may be counted by saying one-thousand, two-thousand, etc. Keep room closed for 15 minutes after treatment. Example: a room 20 feet long, 12 feet wide, and 8 feet high requires 2 seconds if closed tight; 4 seconds if not. Do not remain in treated areas. Ventilate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

STANCHION BARN USE--Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

(Continued)

3/4

(Right Side Panel)

DIRECTIONS FOR USE (Con't)

ANIMAL USE--From approximately 2 foot distance, thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

POULTRY OPERATIONS: Use only in empty poultry houses (when birds are not present). Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.

HOG OPERATIONS: In hog houses, Close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. For animal application, direct the spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN AND STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. In animal application, lightly mist over backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horse's face or eyes. Repeat application daily or as necessary.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

NON-AGRICULTURAL USES

KENNELS AND CATTERIES: Remove animals before spraying. Ventilate after treatment is completed.

FLYING INSECTS: Flies, Mosquitoes, Gnats, Wasps, Small Flying Moths. Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cubic feet. Disperse in all locations. Keep area closed for 15 minutes.

ADULT FLEAS in Buildings (Not on Animals). Close windows and doors. Hold aerosol 36" above floor and direct fog toward floor and lower walls at a rate of 10 seconds per 100 square feet, making sure that all floor, sofa, and chair surfaces are contacted. Repeat treatments after 6 days.

FOOD STORAGE AREAS, MEAT AND FOOD PROCESSING PLANTS:

HOME, MEAT, AND FOOD PROCESSING AREA USE: For flies, mosquitoes, small flying moths (millers), and gnats: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated areas. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.
Roaches, Waterbugs, Sowbugs, Spiders and Centipedes: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under

4/4

refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

FOR USE IN STORAGE SITES: This product can be used in warehouse bins and trucks, cargo ships, mills, bin hoppers, elevators, and conveying equipment as a clean up prior to using them for storage. Close doors and windows. Direct aerosol mist to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be resting. Use a minimum treatment of 6 to 8 seconds per 1000 cubic feet of space. Keep area closed for 30 minutes after each treatment. In mills and elevators, all grain infested accumulations should be removed from the bin hoppers. All storage areas and conveying equipment should be thoroughly cleaned by sweeping out the waste grain, cobwebs and other debris from the walls and rafters as well as on the floor and door frames with special attention to material lodged in the cracks and crevices. All of the debris should be removed and burned to kill eggs and insects that might be present. Repeat treatment weekly or as flying moths appear.

FOR USE ON STORED PRODUCTS: This product can be safely used on rice, grain nuts, dried fruit, beans, almond nutmeat and shells, pistachio, dried prunes, dried apricots, raisins, figs, wheat, oats, corn, barley, rye, tobacco and peanuts held in storage for control of the accessible stages of Almond Moths, Angoumois Grain Moths, Cadelle Beetles, Cigarette Beetle, Confused Flour Beetles, Flat Grain Beetles, Granary Weevils, Indian Meal Moths, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Sawtoothed Grain Beetles, Square Necked Grain Beetles, and Tobacco Moths.

ROACHES, SPIDERS, ANTS, SILVERFISH, CRICKETS: Hold aerosol close to cracks and mist directly into them, contacting as many insects as possible. Apply freely in all possible hiding places behind baseboards, sinks, cabinets, coffee makers, meter boxes, manholes. Repeat as necessary.

NOTICE TO USER: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts and accepts the product on these conditions. (RV 102999)