

Prozap® GARDEN DUST

An Insecticide Dist For Use Against Certain Insects On Vegetables.

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl methylcarbamate) 5%

OTHER INGREDIENTS:.....95%

TOTAL100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

- If Swallowed: Call a poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
- If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise.
- If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advise.
- If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advise.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For emergency information contact: National Pesticide Information Center at 1-800-858-7378 between 6:30 am and 4:30 Pacific Time. After 4:30 call your poison control center.

NOTE TO PHYSICIANS: Carbaryl is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

ACCEPTED
MAY 13 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 47000-109

EPA Reg. No. 47000-109

EPA Est. No. 47000-IA-1

Manufactured By:
Chem-Tech, Ltd.
Des Moines, IA 50321

NET WEIGHT:

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash hands thoroughly with soap and water after handling. Remove clothing and wash clothing before reuse.

USER SAFETY REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.

Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

Do not allow children or pets to contact treated areas until dust has settled.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer, or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

For control of Ants, Centipedes, Crickets, Cutworms, Earwigs, Grasshoppers, Millipedes, Sowbugs, Springtails and ticks, apply product evenly over the soil surface.

CROP	POUNDS OF DUST PER ACRE	POUNDS OF DSUT PER 1000 SQ. FT.	PREHARVEST INTERVAL	SPECIFIC DIRECTIONS
Asparagus	40	0.90	1 day	Repeat applications as necessary up to a total of 3 times prior to harvest or a total of 4 times per year but not more often than once every 7 days.
Beans, Peas, Cowpeas, Lenatis	30	0.70	3 Days (fresh beans or peas) 14 Days (forage) 21 Days (dried beans or peas, seed or hay)	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard greens, Kohlrabi, Chinese Cabbage	40	0.90	3 Days (Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kohlrabi) 14 Days (Chinese Cabbage, Collards, Kale, Mustard greens)	Repeat applications as necessary up to a total of 3 times per year but not more often than once every 7 days.
Carrots, Garden Beets (roots), Horseradish, Parsnips, Potatoes, Radishes, Rutabagas, Salsify, Sweet Potatoes, Turnips (roots)	40	0.90	7 Days	Repeat applications as necessary up to a total of 3 times per year but not more often than once every 7 days.
Celery, Dandelion, Endive, Garden Beets (tops), Hanover Salad, Lettuce (head and leaf), Parsley, Rhubarb, Spinach, Swiss chard, Turnips (tops)	40	0.90	14 Days	Repeat applications as necessary up to a total of 3 times per year but not more often than once every 7 days.
Corn (Field and Pop)	40	0.90	14 Days (forage, silage) 48 Days (grain, fodder)	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 14 days.
Sweet Corn	40	0.90	2 Days (ears) 14 Days (forage) 48 Days (fodder)	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.
Cucumber, Melon, Squash	20	0.45	3 Days	Repeat applications as necessary up to a total of 6 times per year but not more often than once every 7 days.
Eggplant, Peppers, Tomatoes	40	0.90	3 Days	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.
Grain Sorghum (not sweet)	40	0.90	14 Days (grazing, forage)	Repeat applications as necessary up to a total of 3

			21 Days (grain, fodder)	times per year but not more often than once every 7 days.
Strawberries	40	0.90	7 Days	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days. Carbaryl may injure Early Dawn and Sunrise varieties of strawberries.
Sugar Beets	30	0.70	28 Days (Roots, forage)	Repeat applications as necessary up to a total of 2 times per year but not more often than once every 14 days.
Wheat	30	0.70	21 Days (grain, straw) 7 Days (forage)	Repeat applications as necessary up to a total of 2 times per year but not more often than once every 14 days.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in original container, tightly closed. High temperatures may shorten shelf life of product.

Container Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call you local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE

It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Chem-Tech, Ltd, the manufacturer or seller. In no case shall Chem-Tech, Ltd, the manufacturer or seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Except as expressly provided herein, Chem-Tech, Ltd, the manufacturer or seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. Buyer's or user's exclusive remedy, and Chem-Tech, Ltd's, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.

Manufactured by:
Chem-Tech, LTD
Des Moines, IA 50321

09/04