

ORLIK

DURSBAN GRANULES

Controls Chinch Bugs, Sod Webworms (Lawn Moths), Ants,
Fire Ants, Brown Dog Ticks, Crickets, Cutworms, Earwigs,
Sowbugs (Pillbugs), Certain White Grubs*, Fleas and
Hyperodes (Turfgrass) Weevils** Infesting Home Lawns
and Other Ornamental Turf Areas

ACTIVE INGREDIENT:

Chlorpyrifos (O,O-diethyl O-(3,5,6-trichloro-2-pyridyl)
phosphorothioate)..... .5%

INERT INGREDIENTS:..... .99.5%

100.0%

Dursban^(R) is a Registered Trademark of DowElanco

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Additional Precautionary Statements

NET WEIGHT: 8 POUNDS

MANUFACTURED BY:

Phaeton Corporation
1000 118th Avenue North
St. Petersburg, Florida 33716

RECEIVED
JUL 21 1992

EPA REG. NO. 47006-5(E:5/24/90)

EPA EST. NO. 47006-FL-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water, and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

Statement of Practical Treatment

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply uniformly to home lawns and other ornamental and recreational turf grass areas using any hand or power operated gravity or rotary spreader designed for application of granular materials. Treat when insects first appear, using this product at a rate of 200 pounds per acre. To help obtain uniform distribution, apply one-half of required amount of product in one direction, then apply the balance at a right angle. Immediately after application, sprinkle the turf with 1,300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine amounts of this product and water needed for small sized areas. Retreat as necessary.

RATES TO USE

393

Area to be treated	Pounds of this Product	Gal. of Water to Sprinkle on Turf after Treatment
200 sq. ft.	1	6

FOR CONTROL OF FLEAS AND BROWN DOG TICKS: Thoroughly apply this product as a spot treatment to infested areas such as: PET BEDS: Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment of pet areas. Do not apply this product indoors. To control the source of your flea infestation, treat pets with an EPA approved product registered for direction application to pets. YARD AREAS: Apply this product uniformly to home lawns, under shrubbery, and other ornamental recreational turf grass areas. Treat when insects first appear using at the rate of one pound per 200 sq. ft. or 4 pounds per 800 sq. ft. To help obtain uniform distribution, apply one-half required amount of product in one direction, then apply the balance at a right angle. Immediately after application, sprinkle the turf with 40 gallons of water per 800 sq. ft. so as to wash the granules into the zone of insect infestation. Retreat as necessary.

DO NOT TREAT PETS WITH THIS PRODUCT

* TO CONTROL CERTAIN WHITE GRUBS (larvae of European chafer and Japanese beetle) use two to four times the amount shown on the label, i.e., 2 to 4 lbs. per 200 sq. ft. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually late July and August. Consult your local Agricultural Extension Service Specialist for proper time to treat in your area.

** TO CONTROL HYPERODES (TURFGRASS) WEEVILS use twice the rate of the product shown on the label. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service Specialist for best time to treat in your area.

FOR FIRE ANT MOUNDS: To control fire ants apply 4 ounces to each mound. After application of granules, sprinkle the mound with one quart of water per six inches of mound diameter. For best results apply in cool weather, 65-80°F, or in early morning or late evening hours. Do not disturb the mound before or during the application. Treat new mounds as they appear.

STORAGE AND DISPOSAL

STORAGE Store in original container only, preferably in a locked area. Keep container tightly closed when not in use. Store in a cool, dry place. Do not use food or drink containers for mixing or storage.

PESTICIDE Securely wrap original container in several layers of newspaper and **DISPOSAL:** discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.