

JUN 23 1988

PH 23
47056-48
P. 1

EPA\HRM\47056-AP.LA1 NO #49
EPA File Symbol: 47056-

Page 1 of 2

TURFIC CRABOUT 92 PLUS

BALAN® PRE-EMERGENT HERBICIDE PLUS 25-3-3 FERTILIZER

20 POUNDS COVERS 5,000 SQ.FT. FOR NORTHERN LAWNS
20 POUNDS COVERS 4,000 SQ.FT. FOR SOUTHERN LAWNS

For Use On Established Home Lawns and Ornamental Turf BLUEGRASS-RYEGRASS-FESCUE-BANHAGRASS-BERMUDA GRASS CENTIPEDE GRASS- ST. AUGUSTINE GRASS- ZOYSIA GRASS

To Control Crabgrass, Foxtail, Goosegrass, Watergrass and Annual Bluegrass (Poa Annua)

BALAN® is a Reg. Trademark of the Elanco Products Company

ACTIVE INGREDIENT:

N-butyl N-ethyl, alpha, alpha, alpha,
trifluoro-2,5-dinitro-p-toluidine 0.92%

INERT INGREDIENTS: 99.08%
TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED. Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water.


SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 47056-

EPA EST. NO. 47056-FL-1

NET WEIGHT 20 POUNDS

H. R. MCLANE, INC.
7210 S.W. 57TH AVENUE
MIAMI, FLORIDA 33143



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters. This product is toxic to fish. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original container in a cool, dry, locked place out of reach of children. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. HOUSEHOLD: Do not re-use empty bag. Wrap and put in trash.

DIRECTIONS FOR USE (Cont.)

When applied according to these Directions this product can be used safely on the following turfgrass varieties. PERENNIAL BLUEGRASS VARIETIES, PERENNIAL RYEGRASS VARIETIES, BERMUDAGRASS, BANHAGRASS, CENTIPEDEGRASS, FESCUE VARIETIES, ST. AUGUSTINE GRASS and ZOYSIAGRASS lawns and ornamental turf.

WEEDS CONTROLLED

This product contains a pre-emergence herbicide plus a complete lawn food. It should be applied to established turf for the control of CRABGRASS (Smooth and Hairy), GOOSEGRASS (Silver Crab or Crowfoot), BARNYARDGRASS (Watergrass), FOXTAIL (Green and Yellow), and ANNUAL BLUEGRASS (Poa Annua)

When properly applied it will control these grassy weeds and will supply southern grasses with 1.00 to 1.25 lbs. nitrogen fertilizer/1,000 Sq.Ft.

BEST AVAILABLE COPY

WHERE, WHEN AND HOW TO APPLY THIS PRODUCT: Apply to turf prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns.

DO NOT: 1. Apply in the spring to Bermuda Grass which has been overseeded with winter grasses (rye) since thinning will occur.

DO NOT: 2. Do not use on Dichondra, nor to newly sprigged areas of Bermudagrass, St. Augustine Grass or Centipede Grass, nor to Bentgrass.

APPROPRIATE TIMES TO APPLY THIS PRODUCT FOR CONTROL OF WEED SEEDS THAT GERMINATE DURING THE FOLLOWING PERIODS

NORTHERN LAWNS - FOR CRABGRASS, FOXTAIL, GOOSEGRASS, WATERGRASS

Germinates - Mid March up to mid May - -Apply Mar 1st to May 1st

- FOR POA ANNUA (Annual Bluegrass)

Germinates - Mid August to October 1st. - Apply Aug. 1st to Sept 15th.

SOUTHERN LAWNS - FOR CRABGRASS, FOXTAIL, GOOSEGRASS, WATERGRASS

Germinates - February First to Mid April - - Apply Jan 15th to April 1st

- FOR POA ANNUA (Annual Bluegrass)

Germinates - October: First thru December- Apply Sept. 15th to Dec. 1st.

APPLY THIS PRODUCT APPROXIMATELY 2 WEEKS AHEAD OF WEED SEED GERMINATION Complete all cultural practices, such as aeration before application. Best weed control will be obtained if area is lightly irrigated soon after treatment.

HOW MUCH TO USE

NORTHERN LAWNS (Cool Season Grasses) - For Crabgrass (etc.) apply 20 lbs. to cover 5,000 Sq. Ft. (equal to 1.62 lbs. Active /acre). For Poa Annuua control apply 20 lbs. to cover 4,000 /sq./ft.

SOUTHERN LAWNS (Warm Season Grasses) - For Crabgrass (etc.) apply 20 lbs. to cover 4,000 Sq. Ft. (equal to 2.02 lbs. Active /acre). For Poa Annuua control apply 20 lbs. to cover 2,700 /sq./ft. (2.99# Herbicide/A.)

RESEEDING TREATED AREAS

Delay reseeding 12 to 16 weeks after treatment. When reseeding increase amount of seed needed for area by 25%.

SPREADER SETTINGS

Spreader Make	Southern Lawns	
	Crabgrass	Poa Annuua
SCOTT #35	5 1/2	6
CYCLONE B-1	4	4 1/2

These are approximate settings. You must calibrate your own spreader since it may vary due to worn or damaged condition.

20 lbs. to treat 5,000 Sq.ft.= 176 lbs/acre or 1.62 lbs. Herbicide/acre.

GUARANTEED FERTILIZER ANALYSIS

- Total Nitrogen (N) 25%
- Ammoniacal 1.2
- Urea Nitrogen 17.3
- Sulfur Coated Urea 6.5
- Available Phosphoric Acid (P2O5) 3%
- Soluble Potash (K2O) 3%

Primary Plant Food derived from: Urea, Sulfur Coated Urea, Diammonium Phosphate, Muriate of Potash.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.