

HANDY READY TO USE DUST GUN

TO OPERATE — Remove tape from vent and use pumping stroke

Dust underneath and top of foliage thoroughly. Apply to both plants and insects. Begin application when larvae are well established or when new ones appear. Dust early in the morning or evening when plants are dry. Repeat every 7 to 10 days or as necessary, whether insects or diseases are present or not to help prevent infestation. Use freely on the soil around garden plants for added protection against certain insects which work on the surface.

CONTROL CHART

Apply before plants are infested and damaged by the following insects and diseases

- 7 Day Interval Between Last Application and Harvest: Corn, Sweet: Helminthosporium Leaf Blight, Flea Beetle, Japanese Beetle, Corn Earworm. Do not feed straw to dairy animals or animals being finished for slaughter.
- Tomato: Early and Late Blight, Anthracnose, Septoria Leaf Spot, Vinegar Flies, Drosophila spp.
- 5 Day Interval Between Last Application and Harvest: Santolugo, Mushmelo, Watermelon: Downy Mildew, Alternaria Blight, Anthracnose, Cucumber Beetle, Cabbage Looper, Melonworm, Serpentine Leaf Miner, Thrips, Aphids, Leaf Hopper, Mites.
- 5 day interval Between Last Application and Harvest: Peppers: Anthracnose, Ripe Rot, Flea Beetle, Aphids, Serpentine Leaf Miner.
- 7 Day Interval Between Last Application and Harvest: Beans: Rust, Anthracnose, Downy Mildew, Mexican Bean Beetle, Black Bean Aphid, Leaf Miner, Leafhopper.
- Cabbages, Cauliflower: Downy Mildew, Cercospora & Alternaria Leaf Spot, Imported Cabbageworm, Diamond-back Moth, Cabbage Looper, Flea Beetle, Aphid.
- Cucumber: Alternaria Blight, Cucumber Beetle, Serpentine Leaf Miner, Aphids, Thrip, Strawberry: Leaf Spot, Leaf Scorch, Strawberry Leaf Roller, Aphids.
- 10 Day Interval Between Last Application and Harvest: Onions: Onion Thrip, Peas: Pea Aphid, Leaf Miner.
- Carrots, Radishes, Turnips: Cercospora and Alternaria Leaf Spot, Downy Mildew, Flea Beetle, Aphids, Dipterous Leaf Miner, Lettuce, Spinach: Downy Mildew, White Rust, Flea Beetle, Dipterous Leaf Miner, Celery: Early and Late Blight, Flea Beetle, Aphids. Remove residues at harvest by washing, stripping and trimming.
- 14 Day Interval Between Last Application and Harvest: Beets: Downy Mildew, Cabbageworm, Aphids, Dipterous Leaf Miner.

FLOWERS:

- Carnations, Chrysanthemums, Spandromas: Thrip, Rust, Leaf Spot, Anthracnose. Gladiolus: Thrips, Botrytis, Curvularia, Sclerophyllum. Azaleas, Camellias: Petal Blight. Apply dust liberally to ground and litter under the plants and into the flowers of azalea and camellias, beginning when flower buds swell and repeating at 5 to 7 day intervals through blooming season. Roses: Aphid, Strawberry Rootworm Beetle, Leafhopper, Black Spot.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risks to persons or property arising out of use or handling and accepts the product on these conditions.

EPA Reg No. 3770-261


Manufactured by:

ECONOMY PRODUCTS COMPANY

OSWEGO, IOWA 51001

MULTI-PURPOSE BUG DUST

ACCEPTED
1/27/72



CAUTION: Keep Out Of Reach Of Children.

May cause irritation of nose and throat. Do not breathe dust. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling. Harmful if swallowed. May be absorbed through the skin. Wash all vegetables thoroughly before using. Keep container closed when not in use. Store in a dry place.

This product is toxic to fish and wildlife. Do not apply where runoff is likely to occur. Apply this product only as specified on this label. Birds feeding on treated areas may be killed. This product is highly toxic to bees exposed to direct treatment of residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

ACTIVE INGREDIENTS:

Zinc (Zinc Ethylene Dithiocarbamate)	8.25%
O,O-Dimethyl C-(2-isopropyl-4-methyl-4-pyrimidinyl) phosphorothioate*	1.50
Zetozene	8.75
Other Cane Resins	1.50
INERT INGREDIENTS:	98.00

*Diazinon®

®Trademark of Geigy Chemical Co. U. S. Patent No. 2,754,243 and other patents pending

NET CONTENTS 8 .