

47123-7
P.M. 12

P 1/2

PARROTT

FORM

BLACK MAGIC

FOR ROSES

FLOWERS • ORNAMENTALS

CONTROLS MANY INSECTS & DISEASES

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-carbamate)	5.00%
CAPTAN*	4.73%
Related Derivatives27%
Sulphur	25.00%
Pyrethrins05%
(Equivalent to ground Pyrethrum Flowers 5.00%)	
INERT INGREDIENTS	64.95%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN
DANGER
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

1 LB. NET WT. (0.453 KG.)
Mfg. by PARROTT AGRO INDUSTRIES, INC.
STAMFORD, CT 06902
Reg. No. 47123-7 EPA Reg. No. 47123-CT-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

A dust mask must be worn when mixing and loading this product.
Clothing worn while loading, mixing and applying this product must be laundered separately from other clothing before reuse. Clothing that may have been drenched or heavily contaminated must be disposed of in accordance with State and local regulations.

Wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to Captan when using indoors or outdoors. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

ACCEPTED
JUN 23 1987
Under FIFRA, this insecticide, Fungicide, or Rodenticide, is Act. as amended, for the pesticide registered under 47123-7
EPA Reg. No. 47123-7

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection

FOR DUSTING: apply thoroughly, early and regularly; make certain to apply dust to lower as well as upper surfaces of plant foliage; make certain to dust early in the season upon the first appearance of the insects and thereafter about every two weeks depending upon the severity of the infestation. It is always advisable to dust when the plants have moisture on them, preferably after a rain.

EFFECTIVE AGAINST

Red Spider, Mites, Aphids, Leaf Hopper, Japanese Beetle, certain caterpillars, Box Elder Bug, White Fly, Lace Bug, Rose Slug, Tarnished Plant Bug, Leaf Roller, Fullers Rose Weevil, Thrips, Rose Chaffer, Rose Midge, Spotted Cucumber Beetle, Harlequin Bug, Rose Scale, and certain other insects on Roses & other Flowers. . . . Also diseases such as Black Spot of Roses, Septoria Leaf Spot of Chrysanthemums, Rust of Snapdragons, and Powdery Mildew of Roses and other Flowers. Make first application when new growth starts in the Spring and repeat applications at 7 to 10 day intervals.

For use in Alabama, New Mexico, Tennessee, Texas & Virginia, the following endangered species restrictions must be followed:

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and general location of their habitats have been identified by county:

Slackwater Darter,	Alabama (Lauderdale, Limestone, and Madison) Tennessee (Lawrence and Wayne)
Pecos gambusia	New Mexico (Chaves and Eddy) Texas (Reeves, Jeff Davis, and Pecos)
Freshwater Mussels	Tennessee (Hancock, Claiborne, Hawkins, and Sullivan) Virginia (Smyth, Scott, Lee, Washington, and Grayson)
Commanch Springs Pupfish	Texas (Reeves and Jeff Davis)

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or State Wildlife Agency if there is any doubt as to the areas where the outdoor use of captan is prohibited.

FOR WET SPRAYING

LUCKY STRIKE ROSE DUST can be used for wet spraying as well as dry dusting. For wet spraying, we recommend the mixing of one to two pounds of dust to about ten gallons of water or approximately five to ten tablespoonsful to each gallon of water. Quantity to be used may vary according to infestation and hardness of insects. Always add the powder to the water and mix thoroughly, and keep material agitated while spraying.