PM 31 47230-4

ACCEPTED
For Acceptance Recommended
Date 47230-4

Sanitizing of Food Processing Equipment and Other Hard Surfaces in Food Contact Locations For sanitizing food processing equipment, dairy equipment, [illegible] refrigerated storage and display equipment and other hard surfaces listed on the

No Potable water rinse is required.

Wash and rinse all articles thoroughly then apply a solution of one **E C C FORMULA 3D4** in 4 gallons of water. To open surfaces. Surfaces should remain wet for at least one minute as indicated by adequate draining and air drying. Fresh solution should be prepared daily when use solution becomes visibly dirty. For manual application the solution may not be reused for sanitizing application.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush or mechanical spray device. **No Potable water rinse is needed.**

Dishes, silverware, glasses, cooking utensils and other similar size food or utensil equipment can be sanitized by immersion in a ½ oz. of gallon solution of **E C C FORMULA 3D4**. No Potable water rinse is required.

At ½ oz. of gallons **E C C FORMULA 3D4** furnishes the [illegible] of Appendix F of the Grade A Pasteurized Milk Ordinances 1978 Recommendations of the U.S. Public Health Services in excess of 800 ppm of hardness calculated as $CaCO_3$ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of ½ oz. **E C C FORMULA 3D4** in 4 gallons of warm water. **No Potable water rinse is required.** Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not let cows lie down in the sanitizing solution. When solution becomes visibly dirty, discard and prepare fresh solution.

Precautionary Statements
Hazards to Humans and domestic animals

Keep out of reach of children. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling mixing or applying. Treatment: Do not breathe spray mist. Avoid contamination of food.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.
OPEN DUMPING IS PROHIBITED.
DO NOT REUSE EMPTY CONTAINER.

PESTICIDE DISPOSAL
PESTICIDE OR RINSATE THAT CANNOT BE USED AS THEM, AUTHORIZED USE SHOULD BE DISPOSED OF IN A LANDFILL OR PESTICIDES IS PERMITTED IN A STATE OR AREA APPROVED BY STATE OR EPA.

CONTAINER DISPOSAL
TRIPLE RINSE (OR EQUIVALENT) AND PUNCTURE CAN TO PREVENT REUSE. THEN OFFER FOR RECYCLING OR REUSE IF AVAILABLE OR DISPOSE IN A SANITARY LANDFILL.
FOR ONE GALLON OR SMALLER CONTAINER: TRIPLE RINSE CONTAINER AND OFFER FOR RECYCLING OR REUSE IF AVAILABLE OR DISPOSE IN A SANITARY LANDFILL.
BULK: WHEN CONTAINER IS EMPTY, DISPOSE IN A SANITARY LANDFILL.

GENERAL
CONSULT FEDERAL, STATE AND LOCAL DISPOSAL AUTHORITIES FOR ANY ADDITIONAL OR ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

EPA Registration No. 47230-4
EPA Establishment No. 47230-CA-01

3940100 G-2

ECC
FORMULA 3D4

Disinfectant Sanitizer Fungicide Deodorizer with Organic Soil Tolerance for Hospital Institutional Industrial School Dairy and Other Farm Use

Active Ingredients

Alkyl (C 50% C 40% C 10%) dimethyl benzylammonium chlorides	10.00%
Octyl decyl dimethyl ammonium chloride	2.25%
Didecyl dimethyl ammonium chloride	1.25%
Dioctyle dimethyl ammonium chloride	1.25%
Inert Ingredients	85.25%

For Sale For Use And Storage By Maintenance Personnel Only

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. Call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, or gelatin solution if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

FOR INSTITUTIONAL USE ONLY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply **E C C FORMULA 3D4** with a cloth mop or mechanical spray device. When applied with a mechanical spray device surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty. **Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions**

For disinfecting floors, walls, countertops, bathroom areas, lavatories, washrooms, tables, chairs, garbage cans and other similar porous surfaces.

Add 1½ / **E C C FORMULA 3D4** to 4.5 gallons of water. Apply to previously cleaned hard surfaces. At this use level, **E C C FORMULA 3D4** is effective against Pseudomonas aeruginosa, Staphylococcus aureus and Salmonella choleraesuis in the presence of 5% blood serum when evaluated by the AOAC Use Dilution test.

Disinfection in Institutions, Industry and Schools Add 2 oz. of **E C C FORMULA 3D4** to 4.5 gallons of water. At 0.5 oz. to 2.25 gallons of water use level the fungicidal effectiveness of **E C C FORMULA 3D4** in the presence of 5% blood serum against [illegible] mentagrophytes has been shown utilizing the AOAC Fungicide Test.

Disinfection of Barber Tools. Pre-cleaned barber tools (such as combs, hair shears, razors and scissors) can be disinfected by immersing in a 0.5 oz. gallon solution of **E C C FORMULA 3D4**.

Disinfection of Poultry Equipment, Animal Quarters and Kennels [illegible] brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of **E C C FORMULA 3D4** to 4 gallons of water. Similar results should be obtained with this solution. Prior to disinfection all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other [illegible] equipment or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces [illegible] before poultry or other animals or poultry equipment with treatment has been absorbed. All treatment equipment should be rinsed [illegible] with potable water prior to [illegible] water for drinking use.

NET CONTENTS: 1 GALLON

MANUFACTURED BY
ENTERPRISE CHEMICAL COMPANY
Garden Grove, California 92641
South Elgin, Illinois 60177

BEST AVAILABLE COPY

1 of 1