


BUG-JUICE WHITE INSECTICIDE FLAT LATEX PAINT (R.T.U.)

Kills Crawling Insects When Exposed to Treated Surfaces. Kills Flying Insects Alighting on Treated Surfaces.

Net Contents 50 US Gallons (190 Liters)

CAUTION
KEEP OUT OF REACH OF CHILDREN



Active Ingredient:

Deltamethrin: (s)-alpha-cyano-3-phenoxybenzyl-(1R,3R)-3-(2-2dibromovinyl)-2,2-dimethyl-cyclopropanecarboxylate..... 0.075%
Other Ingredients:..... 99.925%

Total..... 100.000%

EPA Registration No. 47332 – 12 contains 0.42 lb. Deltamethrin per gallon
EPA Est. No. 47332-WA-01

Shake well before using.

READ ENTIRE LABEL BEFORE USING

PRECAUCION AL USUARIO: Si usted no lea ingles, no use este producto hasta que la etiqueta le haya sido explicada amplimente. (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

PRECAUTIONARY STATEMENT: Hazards to humans and domestic animals.

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes or clothing. Avoid breathing vapors or spray mist. Contact with product may result in transient tingling and reddening of the skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash before reuse.

First Aid Statement for Inhalation: If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after 15 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

not apply to exterior surfaces if rain is expected within 12 hours. Close container after each use. Do not allow children or pets to enter the treated area until surfaces are dry.

Equipment: Apply paint with brush, roller, or conventional airless equipment.

PEST CONTROL INDOOR:

Treatment Sites: This product, when used in accordance with label directions, Bug Juice paint may be applied in and on residential and non-residential buildings and structures including residential kitchens and all types of food/feed handling establishments. This product can be applied to non-food contact surfaces of walls, floors, ceilings, in and around cabinets, behind and beneath equipment and appliances, around floor drains, window and door frames and around plumbing, sinks and other possible pest harborage sites. The paint can be applied only as a crack and crevice or spot treatment indoors. Do not use for broadcast interior applications such as entire walls, ceilings, and floors. Permitted areas of use include, but are not limited to, the food/feed and non-food/non-feed areas of:

Aircraft (cargo area only)	Industrial buildings
Apartment buildings	Installations
Bakeries	Kitchens
Bottling facilities	Laboratories
Breweries	Mausoleums
Buses	Meat, poultry and egg processing
and	packaging plants
Cafeterias	
Candy plants	Mobile and motor homes
Canneries	Nursing homes
Dairy product processing plants	Offices
Food manufacturing plants	Railcars
Food processing plants	Restaurants
Food service establishments	Schools
Granaries and grain mills	Ships and vessels
Hospitals	Trailers
Hotels	Trucks
Houses	Warehouses
	Wineries

Do not use this application method in the food/feed handling areas of the facility when the facility is in operation or foods/feeds are exposed. Do not apply directly to food/feed products. Cover or remove all food/feed processing and/or handling equipment during application. After application in food/feed processing plants, cafeterias and similar facilities wash all equipment, benches, shelving and other surfaces which food/feed will

Do not apply to classrooms when in use. Do not use in aircraft cabins. Do not apply this product in occupied patient rooms or in any rooms/areas while occupied by patients, the elderly or infirm. Do not apply to institutions including libraries, sports facilities or in the immediate area when occupants are present.

Pantry Pest Control: To kill adult and accessible stages of insect pests: *Cockroaches and Ants*. Make application to cupboards, shelving and storage areas. Remove all foodstuffs, all utensils, shelf paper and other objects before painting with Bug Juice paint. Allow all treated surfaces to dry and cover shelves with clean paper prior to replacing foodstuffs, utensils or other items. Any foodstuff accidentally contaminated with any paint solution should be discarded.

Crawling Insect Pests (such as Cockroaches, Mosquitoes, Imported and Southern Fire Ants, Leaf Cutter Ants, Silverfish, Citrus and Pecan Weevil): Apply Bug Juice paint to localized areas of walls, ceilings and other resting areas as a crack and crevice or spot treatment. May be used inside residential buildings, as well as in and around carports, garages, storage sheds, unoccupied horse stables and pet kennels.

Pet Kennels: For residual control of crawling and flying insects in and around these structures, apply as cracks and crevices or spot treatment. Optimal control results if facilities to be painted are clean, and painted surfaces are cleaned regularly (every 21 days). Apply only to areas of facility where animals are not present. Allow paint to dry before allowing re-entry of animals into painted areas. Do not apply to animal feed or watering equipment.

Food/Feed Handling Establishments: For control of all labeled pests, applications are permitted in food/feed and non-food/non-feed areas of food/feed handling and processing establishments as spot or crack and crevice treatment.

Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food/feed. Serving areas where food/feed is exposed and the facility is in operation are also considered food/feed areas.

Non-Food/Non-Feed Areas: Examples of non-food/non-feed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling). All areas that insects inhabit or through which insects may enter should be painted.

Do not use this application method in the food/feed handling areas of the facility when the facility is in operation or foods/feeds are exposed. Do not apply directly to food/feed products. Cover or remove all food/feed processing and/or handling equipment during application. After application in food/feed processing plants, cafeterias and similar facilities wash all equipment, benches, shelving and other surfaces which food/feed will

contact. Clean food/feed handling or processing equipment and thoroughly rinse with clean, fresh water. After application in bakeries, clean all food contact surfaces thoroughly with appropriate equipment.

Spot or Crack and Crevice Applications: Except in federally inspected meat and poultry plants, spot or crack and crevice applications may be made while the facility is in operation, provided exposed food is covered or removed from area being painted prior to application. Do not apply directly to food or food-handling surfaces. In the home, do not allow paint to contact food/feed surfaces. If the paint does contact food/feed surfaces, clean surface with soap and water.

POLE and TREE BANDING:

USE: Use to control cockroaches, imported fire ants, Southern fire ants, leaf cutter ants, silverfish and weevils. Apply product to pole with a 2 ft. band between ground level and 6 ft. Apply product to fruit and nut trees with a 2 ft. band between ground level and 6 ft. during the dormant or quiescent period. Product can be applied to ornamental trees at any time.

Equipment: For tree trunk or pole banding, apply product via course spray, using conventional airless sprayer.

PALLETS:

USE: Use to control cockroaches, silverfish and fire ants. Apply 1 gallon of Bug-Juice paint to 3000 square feet of pallet. May be applied by conventional airless sprayer or dipped in appropriate container. Do not apply to pallet surfaces that will come in direct contact with food or feed.

PEST CONTROL OUTDOOR

For Commercial/Industrial Use Only

Kills cockroaches in sewer manholes and related sanitary sites.

USE: Use to control cockroaches in manholes, sewers, sewer clean-outs, floor drains, and underground utility vaults. Apply product on manhole walls. Product may be applied via course spray using conventional airless spray equipment. Persons applying product in manhole must comply with CLOSED ENTRY requirements of government agencies.

Equipment: Apply paint with remote access application equipment.

Training: For sewer manholes and related sanitary sites, Walla Walla Environmental Inc. or its representatives will train and authorize all applicators in the proper use of Bug-Juice White Insecticide Flat Latex Paint.

Residual Application: Apply Bug Juice paint to surfaces of buildings, porches, window frames, eaves, patios and garages, to control American and German Cockroaches, Silverfish, Imported and Southern Fire Ants, Leaf Cutter Ants, Citrus Weevil, and Pecan Weevil. Do not apply to furniture, swing sets, or other outdoor playground articles.

CORRUGATED and CARDBOARD.

Bug-Juice Coating provides residual protection to the outer surface of corrugated cardboard boxes against insect attack or infestation. This product is for Commercial/Industrial use only. To protect the box from damage from crawling and flying insects, such as silverfish, cockroaches, ants and weevil, apply Bug-Juice Coating to outer surface of corrugated cardboard at a rate of coverage of 6000 to 8000 square ft. of surface. This coating can only be applied to the outside of the corrugated/cardboard box or to corrugated box liner that is to be used for the exterior portion of the corrugated cardboard box. Such means of application include size press, rod, blade, air knife, offset press, gravure and flexography. **DO NOT APPLY TO INNER SURFACE OF BOXES.**

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of equipment wash-water. Apply this product only as specified on this label.

PHYSICAL or CHEMICAL HAZARDS:

Do not use or store near heat or open flames. Do not ship or store with food, feeds, drugs or clothing.

STORAGE AND DISPOSAL:

Avoid extremes in temperature. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Container Disposal: If empty, do not reuse this container. Place in trash or offer for recycling if available.

NOTICE: Seller warrants that the product conforms to its chemical descriptions and is reasonably fit for the purposes stated on the label.

Do not use open container except for immediate use.

Walla Walla Environmental, Inc.
P.O. Box 1298
Walla Walla, Washington 99362
(509) 522-0490