

PH 31 47371-6

10/2

Formulation HS-652Q

One-Step Germicidal Cleaner and Deodorant

DISINFECTANT PSEUDOMONACIDAL STAPHYLOCIDAL SALMONELLACIDAL
BACTERICIDAL FUNGICIDAL MILDEWSTATIC *VIRUCIDAL

DILUTION: 1:256 1/2 OUNCE PER GALLON OF WATER
(488 ppm Active Quat)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL USE DIRECTIONS

Recommended for use in hospitals, nursing homes, schools, colleges, medical clinics, dental offices, office buildings, hotels, motels, airports, veterinary clinics, farm premises, poultry houses, federally inspected meat and poultry plants, and households. Cleans, disinfects, and deodorizes floors, walls, metal surfaces, painted surfaces, exterior bowl surfaces, empty basins, showers, conductive flooring, lavatory fixtures, plastic surfaces (such as polypropylene, polystyrene, etc.), and other hard, non-porous surfaces.

APPLICATION: Remove heavy soil deposits from surface. Then thoroughly wet surface with a solution of 1/2 ounce of the concentrate per gallon of water. The solution can be applied with a cloth, mop, sponge, or coarse sprayer, or by soaking. Let solution remain on surface for a minimum of 10 minutes. Rinse or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Prepare a fresh solution daily or more often if the solution becomes diluted or visibly soiled.

USDA: For use in federally inspected meat and poultry plants on all hard, non-porous surfaces in inedible product processing areas, non-processing areas, and/or exterior areas.

USDA: For use in federally inspected meat and poultry plants as a floor and wall cleaner for use in all departments. Food products and packaging materials must be removed from the room or carefully protected. All surfaces must be thoroughly rinsed with potable water.

TOILET BOWLS: To clean, apply diluted solution around the bowl and up under the rim. Stubborn stains may require brushing. To disinfect, first remove or expel over the inner trap the residual bowl water. Pour in 3 ounces of the diluted solution. Swab the bowl completely using a scrub brush or mop, making sure to get under the rim. Let stand for 10 minutes or overnight, then flush.

MILDEWSTATIC INSTRUCTIONS: Will effectively control the growth of mold and mildew plus the odors caused by them when applied to hard, non-porous surfaces such as walls, floors, and table tops. Apply solution (1/2 ounce per gallon of water) with a cloth, mop, sponge, or coarse sprayer, making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application weekly or when growth reappears.

A multi-purpose, germicidal detergent effective in the presence of a moderate amount of organic soil (5% serum). Disinfects, cleans, and deodorizes in one labor saving step. Effective against the following pathogens:

- Pseudomonas aeruginosa
- Staphylococcus aureus
- Streptococcus pyogenes
- Enterobacter cloacae
- Enterobacter aerogenes
- Klebsiella pneumoniae
- Proteus mirabilis
- Proteus vulgaris
- Escherichia coli
- Salmonella choleraesuis
- Staphylococcus epidemidis
- Staphylococcus aureus phage 80 & 81
- Shigella flexneri
- Shigella sonnei
- Serratia marcescens
- Mycoplasma gallisepticum
- Aspergillus flavus
- Aspergillus glaucus var. tonophilus
- Trichophyton mentagrophytes
- Candida albicans

- *HIV-1 (AIDS virus)
- *Adenovirus Type 3
- *Vaccinia
- *Influenza A/Texas
- *Herpes simplex Type 1
- *Canine parvovirus
- *Newcastle disease virus
- *Rubella (German Measles)
- *Infectious bronchitis (Avian IBV)
- *Bovine rhinotracheitis virus
- *Avian influenza

ACTIVE INGREDIENTS:

- Didecyl dimethyl ammonium chloride 7.5%
- n-Alkyl (C₁₁, 50%, C₁₂, 40%, C₁₄, 10%)
- dimethyl benzyl ammonium chloride 5.0%
- INERT INGREDIENTS: 87.5%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician immediately. If swallowed, immediately drink a large quantity of water. Get medical attention.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

CORROSIVE: Causes eye and skin damage. Wash contaminated clothing before reuse. Avoid contamination of food, water, or feed. When handling product, protect eyes by wearing goggles or a face shield and protect skin by wearing rubber gloves. Wash thoroughly with soap and water after handling.

EPA Reg. No. 47371-6

EPA Est. No. 47371-IN-01

NET CONTENTS _____
H&S CHEMICALS DIVISION

970 East Tipton, Huntington, IN 46750 (219) 356-7073

ACCEPTED
JUL 22 1994
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 47371-6

*Kills HIV-1 (AIDS virus) on pre-cleaned, environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other setting, in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1)(associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of blood/body fluids, and during decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces and objects before application.

Contact Time: Leave surface wet for 5 minutes at 25°C (room temperature). Use a 10-minute contact time for other viruses, fungi, and bacteria listed.

Disposal of Infectious Materials: Blood/body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL

Keep product under locked storage, inaccessible to children. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

PLASTIC CONTAINERS

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

(FOR 1 GALLON OR LESS)

Do not reuse empty container. Rinse thoroughly, securely wrap in several layers of newspaper, and discard empty container in trash.

FIBER DRUMS WITH LINERS

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

METAL CONTAINERS:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or dispose of by other procedures approved by state and local authorities.

040893

BEST AVAILABLE COPY

Formulation HS-052Q

FARM PREMISE AND POULTRY HOUSE DISINFECTANT

1. Remove all animals and feeds from premises, trucks, coops, crates, and enclosures.
2. Remove all litter and manure from floors, walls, and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent, and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, coops, cars, boats, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set, or dried.
8. After treatment with disinfectant, thoroughly scrub feed racks, troughs, automated feeders, fountains, and waterers with soap and detergent, and rinse with potable water before reuse.

040893

BEST AVAILABLE COPY

[For use as package insert or hang tag or place on label.]

