

Designed for use as a disinfectant for use in hospitals, nursing homes, medical and dental offices.

DISINFECTANT ½ ounce per gal. of water
AOAC USE-DILUTION TEST: Effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella choleraesuis* 1:256

Designed for use as a Sanitizer/Cleaner for use in restaurants & bars, food processing plants and federally inspected meat & poultry plants.

SANITIZER 1 ounce per 5½ gallons of water
AOAC AVAILABLE CHLORINE GERMICIDAL EQUIVALENT CONCENTRATION: Effectiveness against *Salmonella typhi* & *Staphylococcus aureus* 1:704

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FOR DISINFECTING: For use on hard non-porous pre-cleaned surfaces such as floors, walls, metal surfaces, porcelain and plastic surfaces. Remove gross filth and heavy soil deposits. Pre-clean all surfaces prior to disinfecting. Use 1 oz. per 2 gallons of water for a minimum contact time of 10 minutes in a single application. Can be applied by spraying or soaking. The recommended use solution is prepared fresh for each use then discarded.

FOR SANITIZING:

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected. A potable water rinse is not required following use of this compound on previously cleaned hard surfaces provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

FOR USE IN FOOD PROCESSING PLANTS:

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard surfaces. Do not rinse. Wash and rinse all articles thoroughly, then apply a solution of 1 ounce in 5½ gallons of water.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush. Do not rinse.

RESTAURANT AND BAR USE: Dishes, glassware, silverware, cooking utensils, wash with detergent, rinse thoroughly, and immerse in a solution (1 oz. per 5½ gallons) to provide 25 ppm of titratable iodine. Do not rinse.

FOOD AND BEVERAGE SANITIZATION PROCEDURES

BEER & Soft Drink Plants: Thoroughly rinse bottling and pre-mix dispensers with water. Drain thoroughly and fill equipment with 25 ppm solution (refer to use solution chart). Just before resuming operation, flush the 25 ppm solution from equipment with potable water. When changing over between soft drink flavors, rinse system well with water, flush with 25 ppm solution and rinse with potable water. Dilution is 1 oz. per 5½ gallons of water.

BEST AVAILABLE COPY

M-2205

Disinfectant/ Sanitizer/Cleaner

ACTIVE INGREDIENTS:

Nonylphenoxyethyl (ethyleneoxy) ethanol iodine complex	23.12%
Polyethoxy polypropoxy polyethoxy ethanol iodine complex (titratable iodine - minimum 1.75%)	3.62%
INERT INGREDIENTS:	73.26%
Total	100.00%

DANGER: KEEP OUT OF REACH OF CHILDREN. ONLY FOR SALE TO, USE AND STORAGE BY SERVICE PERSONS.

(See right panel for additional precautionary and practical treatment statements)

NET CONTENTS

EPA REG. NO. 47371-26 EPA EST. NO. 47371-IN-01

H&S CHEMICALS DIVISION
 970 EAST TIPTON, HUNTINGTON, IN 46750
 219/356-8100

Sanitizing bottles and cans: Proportion into the final rinse of the container washer or rinses in concentrations of 8-12 ppm titratable iodine. Dilution is 1 oz. per 11-12 gallons of water.

Sanitizing Storage Tanks: After cleaning use a 25 ppm solution for sanitizing beer fermentation and holding tanks. Use solution can be applied by spraying or can be used in conjunction with automated clean-in-place systems. No water rinse is required after sanitizing. Dilution is 1 oz. per 5½ gallons of water.

Milk, Citrus and Ice Cream Processing Plants: Applied routinely as a sanitizing spray (12½ ppm to 25 ppm titratable iodine) on walls, conveyors, equipment and floors will reduce or eliminate odors in the processing area. Use on filling equipment to reduce or eliminate bacteria associated with condensation on equipment and pipes. Regular use as a sanitizer at 25 ppm helps prevent milk stone formation on equipment. Dilution is 1 oz. per 5½ gallons of water.

DANGER

Precautionary Statements-Hazards to Humans and Domestic Animals
 May cause severe eye irritation or eye damage. For eyes, wear goggles or face shield when handling the concentrate. Causes skin irritation. Wear rubber gloves when handling the concentrate. Harmful if swallowed. Avoid contamination of food, water or animal feed. Wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician immediately. If swallowed, drink promptly a large quantity of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be needed.

STORAGE AND DISPOSAL

This product must be kept under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

PLASTIC CONTAINERS

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FIBER DRUMS WITH LINERS

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

METAL CONTAINERS

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

ACCEPTED

47371-26

JUN 12 1988

U.S. DEPARTMENT OF AGRICULTURE
 NATIONAL ANIMAL HEALTH SERVICE
 VETERINARY MEDICAL DIVISION

U.S. Reg. No.