

PH 31

47371-89

1071

FORMULATION AE-90

One-step Disinfectant, Cleaner, and Deodorant

BACTERICIDAL FUNGICIDAL
MILDEWSTATIC *VIRUCIDAL

- Broad-spectrum activity.
- Hospital-use disinfectant effective against Staph, Strep, and Pseudomonas.
- Non-acid and non-abrasive; will not etch or scratch delicate surfaces.
- Effective against *HIV-1 (AIDS virus) and *Herpes simplex type 2.
- Controls mold and mildew.
- Dissolves soap scum and foams away dirt.
- Disinfects, cleans, and deodorizes in one labor-saving step.

A ready-to-use, multi-purpose formula effective in the presence of a moderate amount of soil (5% organic serum) against Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, Trichophyton mentagrophytes, *Herpes simplex type 2, and *HIV-1 (AIDS virus).

ACTIVE INGREDIENTS:

n-Alkyl (C₁₄ 60%, C₁₂ 30%, C₁₀ 5%, C₈ 5%)
 dimethyl benzyl ammonium chloride 0.09%
 n-Alkyl (C₁₂ 50%, C₁₄ 30%, C₁₆ 17%, C₁₈ 3%)
 dimethyl ethylbenzyl ammonium chloride 0.09%
 INERT INGREDIENTS: 99.82%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
PHYSICAL AND CHEMICAL HAZARDS
CAUTION

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

EPA Reg. No. 47371-89
EPA Est. No. 47371-IN-01

Net Contents: _____

H&S Chemicals Division
970 East Tipton Street
Huntington, IN 46750
(219) 356-7073

ACCEPTED
JUN 17 1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 47371-89

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE BEFORE USING

GENERAL USE DIRECTIONS: Recommended for use in households, hospitals, nursing homes, schools, colleges, airports, hotels, motels, garages, and kennels. Disinfects, cleans, and deodorizes non-porous, inanimate surfaces such as floors, walls, metal surfaces, stainless steel surfaces, plastic surfaces (such as polypropylene, polystyrene, etc.), plastic/vinyl chairs, porcelain, garbage cans, laundry chutes, bathroom fixtures, shower stalls, exterior bowl surfaces, empty basins, knobs, handles, railings, vehicles (ships, trains, and airplanes), and many other similar public and residential surfaces which could harbor and transmit dangerous microorganisms.

TO CLEAN: Shake can. Spray evenly over surface, allowing sufficient time for the foam to penetrate heavy soil deposits. Wipe off. Chrome fixtures may be rinsed for high shine.

TO DISINFECT: Clean surface thoroughly as instructed above. Then apply a thin coat of foam, making sure to cover entire surface to be disinfected. Allow foam to remain for at least 10 minutes, then wipe off or allow to air dry.

MILDEWSTATIC INSTRUCTIONS: Follow the disinfection instructions above. Repeat application weekly or when growth reappears.

*Kills HIV-1 (AIDS virus) on pre-cleaned, environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of human immunodeficiency virus type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and during decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of disinfectant.

Contact Time: HIV-1 (AIDS virus) is inactivated after a contact time of 5 minutes at 25°C (room temperature). Use a 10-minute contact time for disinfection against all other viruses, fungi, and bacteria listed.

Disposal of Infectious Materials: Blood/body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL

Keep product under locked storage, inaccessible to children. Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Replace cap and discard container in trash. Do not incinerate or puncture.

BEST AVAILABLE COPY

1994