

CUIDADO: SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO.

DILUTION: 1:64
(600 ppm quat)

2 OUNCES PER GALLON OF WATER

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical devices prior to sterilization or high-level disinfection.

GENERAL USE DIRECTIONS

Recommended for use in hospitals, nursing homes, schools, colleges, commercial and industrial institutions, office buildings, veterinary clinics, animal life science laboratories, federally inspected meat and poultry establishments, equine farms, tack shops, pet shops, airports, kennels, hotels, motels, poultry farms, turkey farms, dairy farms, hog farms, breeding establishments, grooming establishments, funeral homes, and households. Disinfects, cleans, and deodorizes floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, plastic surfaces (such as polypropylene, polystyrene, etc.), and other hard, non-porous surfaces.

APPLICATION: Remove heavy soil deposits from surface. Then thoroughly wet surface with a solution of 2 ounces of the concentrate per gallon of water. The solution can be applied with a cloth, mop, sponge, or coarse spray, or soaking. Let solution remain on surface for a minimum of 10 minutes. Rinse or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Prepare a fresh solution daily or more often if the solution becomes visibly dirty or diluted.

USDA: For use in federally inspected meat and poultry plants on all hard, non-porous surfaces in inedible product processing areas, nonprocessing areas, and/or exterior areas.

USDA: For use in federally inspected meat and poultry plants as a floor and wall cleaner for use in all departments. Food products and packaging material must be removed from the room or carefully protected. All surfaces must be thoroughly rinsed with potable water.

USDA: For use in federally inspected meat and poultry plants as a disinfectant agent for use in all departments. Food products and packaging materials must be removed from the room or carefully protected. Use product in accordance with its label. All surfaces must be thoroughly rinsed with potable water.

TOILET BOWLS: Swab bowl with brush to remove heavy soil prior to cleaning or disinfecting. Clean by applying diluted solution around the bowl and up under the rim. Stubborn stains may require brushing. To disinfect, first remove or expel over the inner trap the residual bowl water. Pour in three ounces of the diluted solution. Swab the bowl completely using a scrub brush or mop, making sure to get under rim. Let stand for 10 minutes or overnight, then flush.

MILDEWSTATIC INSTRUCTIONS: Will effectively control the growth of mold and mildew plus the odors caused by them when applied to hard, non-porous surfaces such as walls, floors, and table tops. Apply solution (2 ounces per gallon of water) with a cloth, mop, sponge, or coarse spray. Make sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

FORMULATION HWS-64/ Formulación HWS-64

ONE-STEP,

GERMICIDAL DETERGENT AND DEODORANT

Disinfectant	Pseudomonacidal	Staphylocidal	Salmonellacidal
Bactericidal	Fungicidal	Mildewstatic	*Virucidal

A multi-purpose, neutral pH, germicidal detergent and deodorant effective in hard water up to 400 ppm (calculated as CaCO_3) in the presence of a moderate amount of soil (5% organic serum) according to the AOAC Use-dilution Test. Disinfects, cleans, and deodorizes in one labor-saving step. Effective against the following pathogens:

<i>Pseudomonas aeruginosa</i> ¹	*Adenovirus type 4
<i>Staphylococcus aureus</i> ¹	*Canine distemper
<i>Salmonella choleraesuis</i>	*Canine parvovirus
<i>Acinetobacter calcoaceticus</i>	*Feline leukemia
<i>Bordetella bronchiseptica</i>	*Feline panleukopenia
<i>Chlamydia psittaci</i>	*Feline picornavirus
<i>Enterobacter aerogenes</i>	*Herpes simplex type 1
<i>Enterobacter cloacae</i>	*Herpes simplex type 2
<i>Escherichia coli</i> ¹	*HIV-1 (AIDS virus)
<i>Fusobacterium necrophorum</i>	*Infectious bovine rhinotracheitis
<i>Klebsiella pneumoniae</i> ¹	*Infectious bronchitis (Avian IBV)
<i>Listeria monocytogenes</i>	*Influenza A/Hong Kong
<i>Pasteurella multocida</i>	*Porcine parvovirus
<i>Proteus mirabilis</i>	*Pseudorabies
<i>Proteus vulgaris</i>	*Rabies
<i>Salmonella enteritidis</i>	*Respiratory syncytial virus (RSV)
<i>Salmonella typhi</i>	*Rubella (German Measles)
<i>Salmonella typhimurium</i>	*Transmissible gastroenteritis virus (TGE)
<i>Serratia marcescens</i>	*Vaccinia
<i>Shigella flexneri</i>	
<i>Shigella sonnei</i>	<i>Aspergillus niger</i>
<i>Staphylococcus aureus</i> , (methicillin resistant)	<i>Candida albicans</i>
<i>Staphylococcus epidermidis</i> ²	<i>Trichophyton mentagrophytes</i>
<i>Streptococcus faecalis</i> ¹	
<i>Streptococcus pyogenes</i>	

¹ ATCC & antibiotic-resistant strain
² antibiotic-resistant strain only

ACTIVE INGREDIENTS:

Didecyl dimethyl ammonium chloride	2.31%
n-Alkyl (C_{14} 50%, C_{12} 40%, C_{10} 10%) dimethyl benzyl ammonium chloride	1.54%
INERT INGREDIENTS:	96.15%

KEEP OUT OF REACH OF CHILDREN CAUTION/CUIDADO

(See side panel for additional precautionary and practical treatment statements)

EPA Reg. No. 47371-131

EPA Est. No. 47371-IN-01

NET CONTENTS _____

H&S CHEMICALS DIVISION

970 East Tipton Street, Huntington, IN 46750
(219) 356-7073

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water. If irritation persists, seek medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye and skin irritation. Avoid contact with eyes, skin, and clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARD

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

Keep product under locked storage, inaccessible to children. Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL PLASTIC CONTAINERS

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or if allowed by state and local authorities, burn. If burned, stay out of smoke (FOR 1 GALLON OR LESS)

Do not reuse empty container. Rinse thoroughly, securely wrap in several layers of newspaper, and discard empty container in trash.

FIBER DRUMS WITH LINERS

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

METAL CONTAINERS

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or dispose of by other procedures approved by state and local authorities.

ACCEPTED

FEB 27 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 47371-131

BEST AVAILABLE COPY

FORMULATION HWS-64

*Kills HIV-1 (AIDS virus) on precleaned, environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human immunodeficiency virus Type I (HIV-1)(associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate must be worn during all cleaning of blood/body fluids and during decontamination procedures.


Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of disinfectant.

Contact Time: HIV-1 (AIDS virus) is inactivated after a contact time of 4 minutes at 25°C (room temperature). Use a 10-minute contact time for other viruses, fungi, and bacteria listed.

Disposal of Infectious Materials: Blood/body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

[For use as package insert or hang tag or place on label.]

011395



Page 3 of 4
(label directions for use continued)

FORMULATION HWS-64

GERMICIDAL DETERGENT


FARM PREMISE, LIVESTOCK, POULTRY & TURKEY HOUSE DISINFECTANT

DILUTION: 1:64 2 OUNCES PER GALLON OF WATER

DIRECTIONS FOR USE

1. Remove all animals and feeds from premises, trucks, coops, crates, and enclosures.
2. Remove all litter and manure from floors, walls, and surfaces of barns, pens, stalls, chutes, vehicles, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent, and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, coops, cars, boats, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set, or dried.
8. After treatment with disinfectant, thoroughly scrub feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

(For use as package insert or hang tag or place on label.)



011395

4 7 4



H&S Chemicals Division

Page 4 of 4
(label directions for use continued)

FORMULATION HWS-64

CLEANSING OF BODY SURFACES AND BODY ORIFICES OF HUMAN REMAINS:

This product removes body secretions, soil, and blood, along with their accompanying malodor, from the surfaces and orifices of human remains. Using a sponge or washcloth, bathe the entire body with a solution of 2 ounces per gallon of water. A soft handbrush may be employed on surfaces other than the face. Prepare a fresh solution for each remains or when solution becomes visibly dirty or diluted.

[For use as package insert or hang tag or place on label.]

011395

