

PM 31 47371-147

10/4

FORMULATION HS-1210 DISINFECTANT/SANITIZER (3.85%)**germicide and deodorant**

DISINFECTANT 1:64 2 ounces per gallon of water (600ppm)
 SANITIZER 1:256 1 ounce per 2 gallons of water (150ppm)
 SANITIZER 1:192 1 ounce per 1 1/2 gallons of water (200ppm)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL USE DIRECTIONS FOR DISINFECTING: For use on precleaned, hard, non-porous surfaces such as floors, walls, metal surfaces, stainless steel surfaces, porcelain, and plastic surfaces (such as polypropylene, polystyrene, etc.). Thoroughly wet all surfaces with a solution of 2 ounces or 1/4 cup of product per gallon of water. Let solution remain on the surface for a minimum contact time of 10 minutes. Allow to air dry. Can be applied with a mop, sponge, cloth, coarse sprayer, or by soaking. The use solution is prepared fresh for each use then discarded. Rinsing is not necessary unless floors are to be waxed or polished.

TO CONTROL MOLD AND MILDEW: To control mold and mildew on hard, non-porous surfaces preclean surfaces. Then apply 2 fluid ounces per gallon solution with mop, sponge, cloth, or coarse sprayer so as to wet all surfaces thoroughly. Let air dry. Repeat application every 7 days to control recurrence of growth.

FISHERY, MILK, CITRUS, ICE CREAM & POTATO PROCESSING PLANTS: Applied routinely as a disinfectant on walls and floors will reduce or eliminate odors in the processing area. Use on filling equipment to reduce or eliminate odors in the processing area and bacteria associated with condensation on equipment and pipes.

FOR SANITIZING FOOD CONTACT SURFACES: For use on precleaned, hard, non-porous surfaces such as counter tops, eating utensils, and drinking glasses. Remove gross food particles and soil prior to washing and rinsing the surfaces. Rinse with potable water prior to application of sanitizing solution. Then use 1 ounce per 1 1/2 - 2 gallons of water (150-200 ppm active quat) for a minimum contact time of 1 minute in a single application. Can be applied with a mop, sponge, cloth, or coarse sprayer, or by soaking. Thoroughly wet all surfaces. The recommended use solution is prepared fresh for each use then discarded. Drain thoroughly. Do not rinse. For mechanical applications, use solution may not be reused for sanitizing applications.

USDA: Before use as a sanitizer in federally inspected meat and poultry plants and dairies, food products and packaging materials must be removed from the room or carefully protected. A potable water rinse is not required following use of this compound on previously cleaned hard surfaces provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

FOR WISCONSIN ONLY**SANITIZING EATING AND DRINKING UTENSILS IN EATING ESTABLISHMENTS**

1) Scrape and pre-wash utensils and glasses whenever possible. 2) Wash with a good detergent or compatible cleaner. 3) Rinse with clean water. 4) Sanitize in a solution of 1 fluid ounce to 1 1/2 - 2 gallons of water (200 - 150 ppm quat). Immerse all utensils for at least two minutes or for a contact time specified by governing sanitary codes. 5) Place sanitized utensils on a rack or drain-board to air dry. 6) Fresh sanitizing solution should be prepared at least daily or when visibly soiled or diluted.

NOTE: A clean potable water rinse following sanitization is not permitted under Section HSS 196.13 of the Wisconsin Administrative Code.

Pseudomonacidal Staphylocidal Fungicidal *Virucidal

A concentrated, multi-purpose germicide and deodorant effective in 5% organic serum. Disinfects and deodorizes in one labor saving step. Recommended for use in hospitals, nursing homes, schools, colleges, veterinary clinics and animal life science laboratories, equine farms, tack shops, pet shops, airports, kennels, hotels and motels, poultry and turkey farms, dairy farms and hog farms, breeding and grooming establishments, and federally inspected meat and poultry plants.

Bactericidal at 1:64 dilution (2 ounces per gallon of water) against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Salmonella choleraesuis* according to AOAC Use-Dilution Test method, modified in the presence of 5% organic serum. (See reference sheet for complete list of pathogenic organisms).

Fungicidal against *Trichophyton interdigitale* and *Candida albicans* according to the AOAC fungicidal test, modified in the presence of 5% organic serum at a 1:64 dilution.

*Virucidal against *Bovine rhinotracheitis, *Avian Infectious Bronchitis virus, and *Newcastle virus according to the virucidal qualification, modified in the presence of 5% organic serum at a 1:64 dilution.

ACTIVE INGREDIENTS:

Didecyl dimethyl ammonium chloride 2.31 %
 n-Alkyl (C₁₄ 50%, C₁₂ 40%, C₁₄ 10%) dimethyl benzyl

ammonium chloride 1.54 %
INERT INGREDIENTS: 96.15 %

KEEP OUT OF REACH OF CHILDREN**CAUTION**

(See side panel for additional precautionary and practical treatment statements.)

EPA Reg. No. 47371-147
 EPA Est. No. 47371-IN-01

Net Contents _____

H&S Chemicals Division

970 East Tipton Street, Huntington, Indiana 46750

(219) 356-2073

ACCEPTED

APR 2 1986

EPA Reg. No. 47371-147

RESTAURANT AND BAR USE: For use as a sanitizer on dishes, glassware, silverware, cooking utensils, wash with detergent, rinse thoroughly and immerse in a solution of 150-200 ppm active quat (1 ounce per 1 1/2 - 2 gallons of water) for at least 1 minute. Drain thoroughly. Do not rinse.

BEVERAGE PLANTS: For sanitizing of bottling or pre-mix dispensing equipment, after cleaning thoroughly rinse equipment with cold or lukewarm water. Fill equipment with solution of 150-200 ppm quat (1 ounce per 1 1/2 - 2 gallons of water). Allow the solution to remain in contact with equipment for at least 1 minute or until operations resume at which time the sanitizing solution should be drained from the system. To insure the removal of flavors, it is suggested that during changeover between products the system should be cleaned, rinsed and flushed with the sanitizing solution for at least 1 minute. Allow to drain completely before using.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION**

Causes eye irritation. Avoid contact with eyes. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL**PROHIBITIONS**

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL**PLASTIC CONTAINERS**

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

FIBER DRUMS WITH LINERS

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

METAL CONTAINERS

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

051295

Formulation HS-1210 Disinfectant/Sanitizer (3.85%) germicide and deodorant

SANITARY FILLING: Recommended for use in sanitizing bottles or cans in the final rinse application. The product is to be proportioned into the final rinse water line of the container washer or rinser in concentration of 150-200 ppm active quat. A sanitizing solution of 150-200 ppm active quat should be used in an external application for the filler and closing machine.


STORAGE TANK SANITIZING: Recommended for sanitizing beer fermentation and holding tanks. A solution of 150-200 ppm should be applied manually or used in combination with automated cleaning systems. Drain thoroughly. Do not rinse. For mechanical operations or automated systems, the used sanitizing solution may not be reused for sanitizing but may be reused for other purposes such as cleaning.


FOR SANITIZING IN FISHERIES, MILK, CITRUS, POTATO & ICE CREAM PROCESSING PLANTS: For use as a sanitizer on conveyors and equipment to reduce or eliminate odors in the processing area. Also, for use on filling equipment to reduce bacteria associated with condensation on equipment and pipes.

EPA Reg. No. 47371-147

[May be used as box insert and/or hang tag.]

051295





Formulation HS-1210 Disinfectant/Sanitizer (3.85%) germicide and deodorant

POULTRY & TURKEY HOUSE DISINFECTANT

Dilution: 1:64

2 ounces per gallon

Directions for Use

1. Remove all poultry and feeds from premises, trucks, coops, and crates.
2. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks, and other feeding and water appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
7. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

FARM PREMISE DISINFECTANT

Dilution: 1:64

2 ounces per gallon

Directions for Use

1. Remove all animals and feed from premises, vehicles, and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

CITRUS CANCER CONTROL

For prevention of Citrus Canker Disease through treatment of equipment. Effective against *Xanthomonas campestris* pv. *vesicatoria* (Citrus Canker Disease) at 2000 ppm active quat. Treat all equipment thoroughly at a dilution ratio of 1:25 (2000 ppm active quat). Treatments can be applied to field equipment, outer clothing, shoes, vehicles, tires, and any other equipment taken into infested areas. To prevent the spread of Citrus Canker by this artificial means of transportation, treatments are made by trigger spraying, dipping or brushing, using a maximum rate of 2000 ppm quat. Clothing should be either thoroughly rinsed or laundered before reuse. Footwear should be rinsed before reuse.

Formulation HS-1210 Disinfectant/Sanitizer (3.85%)

(Reference Sheet)

BACTERICIDAL AGAINST:

Effective against the following according to the AOAC Use-Dilution Test, modified in the presence of 5% organic serum.

Pseudomonas aeruginosa
Staphylococcus aureus
Salmonella choleraesuis
Escherichia coli
Salmonella typhi
Serratia marcescens
Klebsiella pneumoniae
Streptococcus pyogenes
Shigella flexneri
Proteus mirabilis
Staphylococcus aureus phage 80
Staphylococcus aureus phage 81
Enterobacter aerogenes
Mycoplasma gallisepticum
Listeria monocytogenes
Pseudomonas putida
Aeromonas salmonicida
Vibrio anguillarum

Effective against the following according to the AOAC Germicidal & Detergent Sanitizer Test, modified in the presence of 800 ppm water hardness (calculated as CaCO_3)

Staphylococcus aureus
Escherichia coli

FUNGICIDAL & MILDEWICIDAL AGAINST:

Effective against the following according to the AOAC Use-Dilution Test, modified in the presence of 5% organic serum.

Aspergillus flavus
Aspergillus fumigatus
Aspergillus glaucus
Candida albicans
Trichophyton interdigitale
Alternaria tenuis
Geotrichum candidum
Penicillium digitatum
Phytophthora infestans
Trichoderma harzianum
Fusarium oxysporum
Rhizopus nigricans (stolonifer)


VIRUCIDAL AGAINST:

Effective against the following according to the virucidal qualifications modified in the presence of 5% organic serum.

Bovine rhinotracheitis virus
Avian Infectious Bronchitis virus
Newcastle virus

EPA Reg. No. 47371-147

[May be used as a box insert and/or hang tag.]



051295