

MTP PHENOLIC GERMICIDAL DETERGENT

Cleans, Disinfects, Deodorizes

DILUTION: 1 to 256 ONE-HALF OUNCE PER GALLON
DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL USE DIRECTIONS

USES: For use on hard, non-porous surfaces such as floors, walls, metal surfaces, painted surfaces, exterior bowl surfaces, empty basins, showers, conductive flooring and lavatory fixtures. For use in hospitals, nursing homes, schools, colleges, medical and dental offices, farms, hog farms, equine farms, poultry and turkey farms, federally inspected meat and poultry plants, lark shops, kennels, pet shops, veterinary clinics, and animal life science laboratories. This includes barns, calf pens, laying operations, breeder operations, hatcheries, agricultural buildings and premises, live haul equipment, trucks, foot baths, and meat and poultry processing plants.

APPLICATION: Use 1/2 ounce MTP PHENOLIC GERMICIDAL DETERGENT per gallon of water for a minimum contact time of 10 minutes in a single application. For disinfecting, remove gross filth and heavy soil deposits, then thoroughly wet surfaces. MTP PHENOLIC GERMICIDAL DETERGENT can be applied with a mop, sponge, or cloth as well as soaking. The recommended use solution is used once and discarded. Rinsing is not necessary on floor surfaces unless floors are to be waxed or polished. Tuberculocidal effectiveness is in 10 minutes at 20 °C when used at a dilution of 1/2 ounce per gallon of water.

HATCHERY: Can be used for cleaning, disinfecting and deodorizing floors, walls and equipment throughout the hatchery. This includes incubators, setters, and racks. Add 1/2 ounce per gallon of water. Hot or cold water may be used. Remove loose waste, gross filth, and heavy soil deposits prior to disinfecting. Repeat with fresh solution to disinfect.

For use in federally inspected meat & poultry plants as a floor and wall cleaner. Before use of this compound, food products and packaging materials must be removed from the room and carefully protected. After use of this compound, all surfaces in the area must be thoroughly rinsed with potable water. Use 1/2 ounce per gallon of water.

*Kills HIV-1 (AIDS virus) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, and blood and during decontamination procedures.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application. This cleaning may be accomplished with any cleaning solution, including this product.

Contact Time: HIV-1 (AIDS virus) is inactivated after (5) minutes of contact at 25°C. Use a 10 minute contact time at 20°C for other common types of viruses and bacteria claimed.

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

PSEUDOMONACIDAL STAPHYLOCIDAL
*VIRUCIDAL FUNGICIDAL TUBERCULOCIDAL

MTP PHENOLIC GERMICIDAL DETERGENT is a multi-purpose germicidal detergent proven effective by the AOAC Use-Dilution Method in 400 ppm hard water (calculated as CaCO₃) in the presence of 5% organic bioload (Fetal Bovine Serum). MTP PHENOLIC GERMICIDAL DETERGENT disinfects, cleans, and deodorizes in one labor-saving step.

MTP PHENOLIC GERMICIDAL DETERGENT is effective against the following pathogenic organisms:

Pseudomonas aeruginosa	†Staphylococcus aureus
Enterobacter aerogenes	Streptococcus pyogenes
Escherichia coli	
†Proteus mirabilis	*Rubella Virus
Shigella flexneri	*HIV-1 (AIDS Virus)
Serratia marcescens	*Adenovirus Type 2
Salmonella choleraesuis	*Herpes Simplex Type 1
†Enterobacter cloacae	*Influenza Type A/Mich
Klebsiella pneumoniae	*Vaccinia Virus
Proteus vulgaris	*Newcastle Disease Virus
Salmonella typhimurium	*Canine parvovirus
Salmonella typhi	*Human rhinovirus Type 38
Pseudomonas cepacia	*Polio Type 1 (Chat strain)
**Enterococcus sp.	
Providencia rettgeri	Candida albicans
**Pseudomonas maltophilia	Trichophyton mentagrophytes
Streptococcus faecalis	
Mycobacterium tuberculosis var. bovis	

†Tested against standard ATCC strains and antibiotic strains.

**Tested against antibiotic resistant strain only.

ACTIVE INGREDIENTS:

o-phenylphenol	5.88%
o-benzyl-p-chlorophenol	9.12%
p-tert-amyphenol	2.97%
INERT INGREDIENTS:	82.03%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN DANGER

(See side panel for additional precautionary and practical treatment statements.)

EPA Reg. No. 47371-163
EPA Est. No. 47371-IN-01

Net Contents _____

H&S CHEMICALS DIVISION
970 East Tipton Street
Huntington, WV 25701

BEST COPY AVAILABLE

STATEMENT OF PRACTICAL TREATMENT

In case of skin contact, wash thoroughly with soap and water. In case of eye contact, immediately flush eyes with water for 15 minutes and get prompt medical attention. If swallowed, promptly drink a large quantity of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive: Concentrate causes eye and skin damage. May be absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse. Harmful if swallowed. Avoid contamination of food, water, or feed.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

PLASTIC CONTAINERS

Triple rinse (or equivalent). Then offer to recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or, if allowed by state and local authorities, burn. If burned, stay out of smoke.

(For 1 gallon or less)

Do not reuse empty container. Rinse thoroughly before discarding in trash. Securely wrap original container in several layers of newspaper and discard in trash.


FIBER DRUMS WITH LINERS

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

METAL CONTAINERS

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

121494



DIRECTIONS FOR USE ON FARM PREMISES

1. Do not use in milking stalls, milking parlors, or milk houses.
2. Remove all animals and feeds from premises, trucks, coops, crates, and enclosures.
3. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
4. Empty all troughs, racks, and other feeding and watering appliances.
5. Thoroughly clean all surfaces with soap or detergent and rinse with water.
6. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
7. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
8. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried.
9. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

(Refer to label for precautionary and practical treatment statements.)

[To be used as an insert or hang tag or inserted on label.]

121494

.....

2