

LABEL NO. 1

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles, face shield or safety glasses and handling. Avoid prolonged or repeated contact with skin.

DO NOT DILUTE OR PUT CONCENTRATE IN FOOD OR DRINK CONTAINERS.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

CHEMICAL AND PHYSICAL HAZARDS: EXTREMELY FLAMMABLE! Keep away from fire, sparks, and heated surfaces. Use with adequate ventilation.

STATEMENT OF PRACTICAL TREATMENT:

If in eyes: Flush with plenty of water. Get medical attention. If on skin: Wash with plenty of soap and water. If inhaled: Remove victim to fresh air.

STORAGE AND DISPOSAL: TRIACON should be stored at a temperature in the original container. Unusual conditions under these conditions have an indefinite shelf life, and small quantities of waste may be disposed of by flushing down the drain with a large excess of water. DO NOT CONTAMINATE DRINKING WATER OR WATER USED OR INHABITED BY FISH OR WILDLIFE. KEEP CONTAINER TIGHTLY CLOSED.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. NOT INTENDED FOR HOME USE.

TRIACON is recommended for use with TRIACON ADDITIVE. Four additives are available for use on 6 different crops. THE CORRECT ADDITIVE MUST BE USED FOR A PARTICULAR CROP AND DILUTED STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS.

Corn, barley, and wheat are treated at a stage when 3 to 5 leaves appear. Seedlings, beans and soybeans are treated between the second and third trifoliate stage, and other crops are treated when 2 to 5 sets of two leaves each appear. Application at earlier or later stages of development may not be effective. Only a single application is recommended per growing season.

(continued other side page 1)

Triacon

CROP YIELD ENHANCER

RECOMMENDED FOR USE WITH TRIACON ADDITIVE. See directions for use.

ACTIVE INGREDIENTS

1-Triacetonol...
Inert Ingredients...
Total.....

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for precautionary statements

NET CONTENTS _____ GALLONS.

BIOCHEMICAL RESEARCH CORPORATION
FALLS CHURCH, VIRGINIA 22404

MADE IN U.S.A.

EPA REG. NO. _____

EPA EST. NO. _____

Use the final TRIACON solution in water as soon as possible after dilution - DO NOT STORE OVERNIGHT. Apply the solution as a fine mist to plants in the field at the recommended application rates in the table below, and at the recommended plant height. Application should be made at a temperature of at least 60° F. Application is easily made using any conventional equipment designed to apply foliar fertilizers, pesticides, etc.

CROP:	TRIACON ADDITIVE:	GAL. TRIACON PER GAL. WATER:	APPLN. RATE:	PLANT HEIGHT:
LEANS, CUCUMBERS, WEST LIME	No. 1	1 gal./20 gal.	20 gal/a	6-10"
RADISHES, CHINESE RADISHES, LETTUCE, PEPPERS**	No. 1	1 gal./20 gal.	20 gal/a	4-6"
SOYBEANS	No. 2	1 gal./20 gal.	20 gal/a	10-12"
MATEMEDICINS, CARROTS, TOMATO**	No. 3	1 gal./20 gal.	20 gal/a	4-6"
TOMATOES**	No. 4	1 gal./20 gal.	20 gal/a	6-10"
SUGAR BEETS, ALFALFA	No. 2	1 gal./20 gal.	20 gal/a	4-6"
PEAS	No. 1	1 gal./20 gal.	20 gal/a	5-8"
BARLEY	No. 3	1 gal./20 gal.	20 gal/a	6-8"
WHEAT, RYE	No. 4	1 gal./20 gal.	20 gal/a	6-8"
OTHER VEGETABLE CROPS	No. 1	1 gal./20 gal.	20 gal/a	4-6"

*Use U.S. gallon of TRIACON concentrate is sufficient for one acre. **Plants should be sprayed after they are well established in the field, one to two weeks after transplanting.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE MUST IN ACCORDANCE WITH DIRECTIONS.

U.S. Patent No. 4,333,758,
Other U.S. and Foreign Patents Pending.

ACCEPTED
JAN 20 1984
EPA REG. NO. 47655-1

BEST AVAILABLE COPY

47655-1
Pm 25
1062

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Avoid contact. Do not get in eyes. Wash thoroughly with plenty of water. If on skin, wash with plenty of soap and water. If on clothing, remove clothing and wash separately.

DO NOT FEED OR PUT CONCENTRATE IN FOOD OR DRINK CONTAINERS.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

CHEMICAL AND PHYSICAL HAZARDS: EXTREMELY FLAMMABLE. Keep away from fire, heat, and hot surfaces. Use with adequate ventilation.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Get medical attention.
If on skin: Wash with plenty of soap and water.
If inhaled: Remove victim to fresh air.

STORAGE AND DISPOSAL: TRIACON should be stored at room temperature in the original container. Unused portions under these conditions have an indefinite shelf life, and small quantities of waste may be disposed of by flushing down the drain with a large excess of water. DO NOT CONTAMINATE DRINKING WATER OR WATER USED OR INHABITED BY FISH OR WILDLIFE. KEEP CONTAINERS TIGHTLY CLOSED.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. NOT INTENDED FOR HOME USE.

TRIACON is recommended for use with TRIACON ADDITIVE. The additive is available for use in different crops. THE TRIACON ADDITIVE MUST BE USED FOR A PARTICULAR CROP AND APPLIED STRICTLY IN ACCORDANCE WITH LABEL DIRECTIONS.

Corn, barley, and wheat are treated at a stage when 3 to 5 leaves appear in seedlings; beans and soybeans are treated between the second and third trifoliate stage, and other crops are treated when 2 to 3 sets of two leaves each appear. Applications at earlier or later stages of development may not be effective. A multiple application is recommended per growing season. (continued other side page)

Triacon

CROP YIELD ENHANCER

RECOMMENDED FOR USE WITH TRIACON ADDITIVE.
See directions for use.

ACTIVE INGREDIENTS

1-Triacontanol.....0.0033 g
Inert Ingredients.....89.9967 g
Total.....100.0000 g

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panel for precautionary statements.

NET CONTENTS GALLONS.

BIOCHEMICAL RESEARCH CORPORATION
FALLS CHURCH, VIRGINIA 22046

MADE IN U.S.A.

EPA REG. NO.

EPA EST. NO.

ACCEPTED
JAN 20 1984
U.S. Dept. of Agriculture
Federal Agricultural Marketing Administration
Washington, D.C. 20250
Registered under 716-55-1
EPA Reg. No. 716-55-1

BEST AVAILABLE COPY

Use the final TRIACON solution in water as soon as possible after dilution - DO NOT STORE OVERNIGHT. Apply the solution as a fine mist to plants in the field at the recommended application rates in the table below, and at the recommended plant height. Application should be made at a temperature of at least 60° F. Application is easily made using any conventional equipment designed to apply foliar fertilizers, pesticides, etc.

CROP:	TRIACON ADDITIVE:	GAL. TRIACON* PER GAL. WATER	APPLN. RATE:	PLANT HEIGHT:
BEANS, CUCUMBERS, SWEET CORN	No. 1	1 gal/200 gal	20 gal/a	8-10"
RADISHES, CHINESE RADISHES, LETTUCE, PEPPERS**	No. 1	1 gal/200 gal	20 gal/a	6-8"
TOMATOES	No. 1	1 gal/200 gal	20 gal/a	18-12"
WATERMELONS, CARROTS, TOBACCO**	No. 2	1 gal/200 gal	20 gal/a	4-6"
TOMATOES**	No. 2	1 gal/200 gal	20 gal/a	6-10"
SUGAR BEETS, ALFALFA	No. 2	1 gal/200 gal	20 gal/a	5-8"
PEAS	No. 3	1 gal/200 gal	20 gal/a	5-8"
BARLEY	No. 3	1 gal/200 gal	20 gal/a	6-8"
WHEAT, RYE	No. 4	1 gal/200 gal	20 gal/a	6-8"
OTHER VEGETABLE CROPS	No. 1	1 gal/200 gal	20 gal/a	4-6"

* 1 U.S. gallon of TRIACON concentrate is sufficient for ten acres.
** Ants should be sprayed after they are well established in the field, one to two weeks after transplanting.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS.

U.S. Patent No. 4,333,758,
Other U.S. and Foreign Patents Pending.

2082