

GENERAL INFORMATION

COPPER CONTROL GRANULAR provides effective control of various bluegreen, filamentous, planktonic and bottom algae which can occur in potable water reservoirs, recreation lakes, ornamental, golf courses, fish ponds and hatcheries. COPPER CONTROL GRANULAR treated water may be used to irrigate turf, fairways, putting greens, and ornamental plants immediately after treatment.

ENVIRONMENTAL HAZARDS ✓

Do not apply to water except as specified on label. This pesticide is toxic to fish and aquatic organisms. Direct application of this product to water may cause a significant reduction in populations of aquatic invertebrates, plants, and fish. Consult your State Fish and Game Agency before applying this product to public waters. Permits may be required.

FISH CAUTION: COPPER CONTROL GRANULAR may be toxic to trout and other species of fish. Fish toxicity is dependent upon the hardness of water. Do not use when hardness is less than 50 ppm. Treatment of algae can result in oxygen loss and cause fish suffocation. Read and follow the directions carefully.

PRECAUTIONARY STATEMENTS ✓

CAUTION, HAZARD TO HUMANS: Harmful if swallowed. May cause skin damage. Avoid contact with skin or eyes. In case of contact, wash thoroughly. For eyes; wash with copious amounts of water and get immediate medical attention. Seek medical attention if swallowed.

# COPPER CONTROL GRANULAR

EPA EST. NO. 47677-WA-01 EPA REG. NO.

ACTIVE INGREDIENT:

Copper\* as elemental.....3.7%

INERT INGREDIENTS.....96.3%

TOTAL.....100.0%

Copper Control Granular contains 2.2 lbs. of elemental copper (Cu) per 60 lbs.

✓\*From mixed copper-Mono and Triethanolamine complexes.

CAUTION

KEEP OUT OF REACH OF CHILDREN  
See Side Panel for Additional  
Precautionary Statements

ACCEPTED  
APR 24 1987

Do not register and use  
BPA Reg. No. 47677-8



47677-8

PM 23

1061

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Useful in the control of free floating and bottom algae. For best results apply COPPER CONTROL GRANULAR early in the day. Apply at the rate of 60 pounds per acre (67.2 kgs. per ha) for large area treatment. For small areas treatment apply at 1 pound per 720 square feet. Distribute evenly over the water surface over vegetation to be killed. Use a dry fertilizer spreader to insure even distribution over the area. Begin treatment along the shore and proceed outwards in bands to allow fish to move into untreated waters.

A single application early in the season is usually all that is necessary. For heavily infested areas, retreatment may be required. Retreatment should be after 10-14 days to avoid fish suffocation.

Treatment of algae can result in oxygen loss from decomposition of dead vegetation. This oxygen loss can cause fish suffocation, therefore, if treating an entire area, treat 1/3 to 1/2 of the water area in a single operation and wait 10-14 days before retreatment.

NOTICE: Neither the manufacturer or the seller makes any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use of this material when such use is contrary to label instructions. Read and follow the label directions carefully.

✓ STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal  
PESTICIDE DISPOSAL: Wastes resulting from use of this product must be disposed of according to applicable Federal, State or Local procedures.  
CONTAINER DISPOSAL: Triple rinse or equivalent. Then offer for recycling a reconditioning or dispose of in a sanitary landfill, or by incineration if allowed by State and Local authorities. If burned, stay out of smoke.

NET CONTENTS \_\_\_\_\_ lbs.

LOT# \_\_\_\_\_

**ARGENT** Chemical Laboratories

8702 152nd Ave. N.E. • Redmond, Washington 98053  
(206) 485-3777  
Customer Service 1-800-428-6258

BEST AVAILABLE COPY