



2 of 7 # 47916-47 1/11/81

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

*HJ*  
Henry M. Jacoby  
Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (TS-767)

Enclosure

JOB-10685; Jacoby; Kendrick; 898-1270; tar; RD: 85/12: 3/18/85; del. 3/25/85

1047 # 47916-47 PM21

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (FTR-947) WASHINGTON, D.C. 20460	EPA REGISTRATION NO. 47916-47	DATE OF ISSUANCE APR 4 1985
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	

NOTICE OF PESTICIDE:  REGISTRATION  
 REREГИSTRATION  
 (Under the Federal Insecticide, Fungicide,  
 and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Venley Industries Inc.  
 P.O. Box 494  
 Monticello, N.Y. 12559

161 / 136878  
 18 / 1

**BEST AVAILABLE COPY**

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below to the label for this product for shipment:

- Add the phrase "EPA Registration No. 47916-47."
- Insert the words "EPA Environmental Protection Agency,"
- Insert the heading "General Instructions" and insert the following instructions: "See the label for the complete instructions for use."
- Transfer the warning statement to the "DANGER" section of the label.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

*HJ*  
Henry N. Jacoby  
Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (TS-767)

Enclosure

**BEST AVAILABLE COPY**

JOB-10685; Jacoby; Kendrick; 898-1270; tax; RD: 85/12; 3/18/85; del. 3/25/85



# AGRICULTURAL FUNGICIDE

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

FUNGICIDE/BACTERICIDE

APR 4 1985

ACTIVE INGREDIENT  
CUPRIC HYDROXIDE  
INERT INGREDIENTS

**BEST AVAILABLE COPY**

37.5%  
62.5%  
100.0%

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

47916-47

(METALLIC COPPER  
EQUIVALENT ..... 24.1%)

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
HARMFUL IF SWALLOWED.  
IF SWALLOWED, INDUCE VOMITING. GET MEDICAL ATTENTION.  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

EPA Reg. No. 47916-


EPA Est. No.

NET CONTENTS:



**wesley industries, inc.**

P.O. BOX 490 MONTROSE, ALABAMA 36559 U.S.A.



# WESLEY AGRICULTURAL FUNGICIDE

## STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

## PRECAUTIONARY STATEMENTS

### HAZARD TO HUMANS AND DOMESTIC ANIMALS

**BEST AVAILABLE COPY**

**CAUTION:** Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

### ENVIRONMENTAL HAZARDS

**CAUTION:** This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

## GENERAL INSTRUCTIONS

Use as directed below. Wesley Agricultural Fungicide is adaptable to spraying from all types of spray equipment. Depending on the equipment used and the specific crop, the volume applied per acre will differ. For dilute foliage sprays: use from 25 to 100 gallons of water per acre (GPA) for most vegetable crops, 100 to 800 GPA for fruit orchards and up to 1500 GPA as may be required for large citrus groves. For concentrate ground sprays, apply from 5 to 20 GPA for most vegetable crops and 25 to 100 GPA for fruit and nut crops. For aerial spraying, 3 to 15 GPA are commonly used. No additional surfactants are needed. Add slowly to a spray tank partially filled with water. Spreader-stickers, insecticides, nutrients, etc. should be added last. Observe cautions and limitations on label of all products used in mixtures.

The following specific instructions are based on general applications. The recommendations of the State Agricultural Extension Services should be closely followed as to timing, frequency and number of sprays per season.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Residual spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures. Triple rinse for equipment. Then offer for recycling or reconditioning, or dispose of in a sanitary landfill or by incineration if allowed by State or Local authorities.

*See Attached Replacement*

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**HOMEOWNER USE:** 1 level teaspoon per gallon of water is equivalent to 1 pint per 100 gallons. Make application according to specific instructions given below for those crops where rates are given in pints per hundred gallons. (For example: 2 pints per 100 gallons are equivalent to 2 level teaspoons per gallon.)

**NOTICE:** Wesley Industries, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to the directions under normal use conditions on the plants and crops specified. **THESE ARE NO OTHER WARRANTIES EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE.** This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer. Buyer assumes all risk of any such use.

WESLEY AGRICULTURAL FUNGICIDE

EPA REG. NO. 47916-

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

STORAGE: Store unused product in original container only in cool, dry area out of reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

BEST AVAILABLE COPY





BEST AVAILABLE COPY

## DIRECTIONS FOR USE (CONTINUED)

**MANGO**, *Anthracnose*. Apply monthly after fruit set until harvest. 2 pints per 100 gallons or 10 $\frac{1}{2}$  to 13 pints per acre depending on equipment. Consult Extension Service for date and conditions.

**OLIVES**, (California) *Powdery Mildew*. Make first application at 2 $\frac{1}{2}$  to 4 pints per 100 gallons or at 10 $\frac{1}{2}$  to 16 pints per acre depending on equipment before winter rainfall. A second application in early spring could be made if disease is severe.

**ORION**, *Purple Blotch & Downy Mildew*. Apply 2 $\frac{1}{2}$  pints per acre when plants are 4 to 6 inches high and repeat at 7 to 10 day intervals.

**PLACES & NECTARINES**, *Leaf Curl & Coryneum Blight*. Dormant application. Apply at leaf fall. Dilute. Apply 2 $\frac{1}{2}$  to 5 pints per 100 gallons (minimum of 10 pints per acre). Use 2 $\frac{1}{2}$  pints per acre when rainfall is very heavy and disease pressure is high. Concentrate. Apply at equivalent rates in approximately 100 gallons of water. May be used with agricultural spray oil. *Brown Rot* (*Botrytis*) *Blight*. Apply at 2 $\frac{1}{2}$  to 4 pints per 100 gallons as a full cover spray at pink bud. (Application at this time also affords some control of leaf curl (*Coryneum Blight*)). *Bacterial Spot*. Dormant application. Apply at 2 $\frac{1}{2}$  pints per 100 gallons. Postbloom, apply  $\frac{1}{2}$  pint per 100 gallons at first and second cover sprays. Do not spray later than three weeks prior to harvest. Do not use at rates above those recommended. Caution: Slight defoliation and spotting of leaves may occur from use in cover sprays.

**PEANUTS**, *Cercospora Leaf Spot*. Begin spraying 40-45 days after planting or when disease symptoms appear. Make ground or aerial application at 2 to 4 pints per acre. For aerial application use 3-10 gallons of water. Continue applications at 10-14 day intervals. Use sufficient water to get adequate coverage. May be tank-mixed with Flowable Sulfur Products.

**PEARS**, *Fire Blight*. Apply at  $\frac{1}{2}$  pint per 100 gallons or 1 $\frac{1}{2}$  pints per acre at 5 day intervals throughout bloom period. *Pseudomonas Blight*. Apply before fall rains at a rate of 4-5 $\frac{1}{2}$  pints per 100 gallons (30-400 GPA) and again at dormant before spring growth starts. Excessive dosages may cause fruit russet.

**PEAS**, *Powdery Mildew*. Begin spray treatment when disease symptoms first appear. Use at 2-4 pints per acre depending on disease severity. Repeat applications at weekly intervals.

**PEPPERS**, *Bacterial Spot*. When disease threatens, apply 2 $\frac{1}{2}$  to 4 pints per acre in sufficient water for adequate coverage at 7 to 14 day intervals depending on disease severity.

**PENDON**, *Bacterial Leaf Spot*. Apply weekly before disease appears at 2 pints per acre plus mancozeb at recommended amount of mancozeb per 100 gallons of water.

**POTATOES**, *Early & Late Blight*. Apply at 7-10 day intervals starting when plants are six inches high until two weeks before harvest. Use 1 $\frac{1}{2}$  to 2 pints per acre in those locations where disease is light and up to 4 to 5 $\frac{1}{2}$  pints per acre where disease is more severe.

**PUMPKIN & SQUASH**, *Powdery Mildew*. Begin applications when plants are 3 weeks old or when first disease symptoms appear. Use at weekly intervals at 2-4 pints per acre depending on disease severity.

**STRAWBERRIES**, *Leaf Spot & Leaf Blight*. Apply at 2 $\frac{1}{2}$  to 4 pints in 100 gallons per acre. Begin application when plants are established and continue on a weekly schedule throughout season. Discontinue applications if signs of phytotoxicity appear.

**SUGAR BEETS**, *Cercospora Leaf Spot*. Start spray when disease threatens and continue for 4 to 5 applications. Spray at 10-14 day intervals depending on weather conditions at 2 $\frac{1}{2}$  to 6 $\frac{1}{2}$  pints per acre depending on disease severity. Addition of suitable agricultural spray oil is recommended at 2 quarts per acre.

**SYCAMORE**, *Anthracnose*. Make two applications using 2 $\frac{1}{2}$  to 4 pints per 100 gallons as a full cover spray. Make first application at bud crack and second application 7-14 days later at 10% leaf expansion.

**TOMATOES**, *Early Blight*. When disease threatens, apply 2 $\frac{1}{2}$  to 4 pints per acre at 7-10 day intervals. *Bacterial Speck*. Apply at 2 $\frac{1}{2}$  pints per acre at 10-30 day intervals beginning when the disease threatens. Use more frequent applications when disease pressure is high. *Bacterial Spot*. When disease threatens, apply 2 $\frac{1}{2}$  to 5 $\frac{1}{2}$  pints per acre at 7-10 day intervals, more frequently when disease is severe.

May be combined with mancozeb. Do not apply within five days of harvest.

**WALNUT**, *Walnut Blight*. Apply first spray at early pre bloom when catkins are partially expanded. Make three additional applications during bloom and early leaf out stages at 7-10 day intervals. Additional applications may be necessary when frequent rainfall occurs. Dilute. Apply 2 $\frac{1}{2}$  pints per 100 gallons water (minimum of 10 $\frac{1}{2}$  pints per acre). Concentrate. Apply at equivalent rates in 50-100 gallons water per acre. One pint of summer oil emulsion may be added per 100 gallons of spray. Do not apply more than 10 $\frac{1}{2}$  pints per acre per application.

**WATERMELON**, *Anthracnose & Downy Mildew*. Apply as soon as plants become established and at weekly intervals thereafter. *Anthracnose*. Use at 2 $\frac{1}{2}$  pints per acre. *Downy Mildew*. Use at 2-4 pints per acre, according to disease severity.

**WHEAT & BARLEY**, *Septoria Leaf Blotch & Helminthosporium Spot Blotch*. Apply 2 to 2 $\frac{1}{2}$  pints per acre. Make first application at early boot stage and follow with second application at early head stage.