

**CAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**DANGER:** Corrosive, may cause severe skin irritations, chemical burns to broken skin. Causes eye damage. Do not get on skin or on clothing. Wear goggles or face shield and gloves when handling this product. Wash after use. Avoid breathing vapors. Vacate poorly ventilated areas if possible. Do not return until odors have dissipated.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent should not be discharged into lakes, streams, ponds or rivers without a valid discharge permit. For guidance, contact your regional office of the Environmental Protection Agency.

**PHYSICAL AND CHEMICAL HAZARDS:**

**STRONG OXIDIZING AGENT:** Mix only with water according to label directions. Mixing this product with such as feces, urine, etc., or with ammonia, acids, or other chemicals may release hazardous gases. Irritates eyes, lungs and mucous membranes.

Manufactured by  
**S & S POOLS**  
Route 9, Box 1220  
Plant City, FL 33566

EPA EST. NO.: 48105-1

NET CONTENTS:

ACCEPTED  
48105-1  
SEP 14 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide use of this product.

**S & S POOL SHOCK**

**"FOR SWIMMING POOL CHLORINATION AND SANITIZING."**

ACTIVE INGREDIENT: SODIUM HYPOCHLORITE 9.2%  
INERT INGREDIENTS 90.8%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**FIRST AID:** If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

**DIRECTIONS FOR SWIMMING POOL CHLORINATION**

**Initial Chlorination:** Start up superchlorinate a three 1/2 pint (16 fl. oz.) of this product in a pool of 2500 gallons of water. This dosage is equivalent to 1.0 ppm available chlorine by weight.  
**Pool Maintenance:**  
A. Adjust pool water pH to 7.6-7.8 range and maintain.  
B. Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.6 to 1.0 ppm. One-half pint (8 fl. oz.) of this sodium hypochlorite solution for each 6000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend upon temperature and number of swimmers. Use test kit to make certain pH and chlorine residual are in the proper range.

**DIRECTIONS FOR DISINFECTION OF POTABLE WATER FOR HOME WELL WATER SYSTEMS**

Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 9 parts softened water. Mix sodium hypochlorite solution and water thoroughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm throughout the distribution system as determined by a DPD Chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the local health department. Primary Drinking Water Regulations. Check water frequently with DPD chlorine test kit.  
\*Contact your local Health Department for further details.

**DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT**

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Rinse with water and contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak over. Addition of one fluid oz. (2 lbsps) of this sodium hypochlorite solution per 4 gallons of water will provide approximately 200 ppm available chlorine by weight.  
**NOTE:** This product degrades with age. Use a chlorine test kit and increase dosage as necessary to obtain the required level of available chlorine.

**STORAGE AND DISPOSAL:**

Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Rinse empty container thoroughly with water and either return to manufacturer or discard by placing in trash collection or burying in an approved landfill. Product or residue that cannot be used, should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment.  
**STORE IN AN UPRIGHT POSITION**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**DANGER:** Corrosive. May cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

**PHYSICAL AND CHEMICAL HAZARDS:  
STRONG OXIDIZING AGENT:** Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc. or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

**S & S POOLS  
Route 9, Box 1220  
Plant City, FL 33566**

EPA REG. NO. 48105-1 EPA EST. NO.:

NET CONTENTS:

ACCEPTED

48105-1  
SEP 14 1982

Under the Federal Insecticide,  
Fungicide and Rodenticide Act,  
as amended, for the pesticide

**S & S POOL SHOCK**

**"FOR SWIMMING POOL CHLORINATION  
AND SANITIZING."**

ACTIVE INGREDIENT  
SODIUM HYPOCHLORITE 9.2%  
INERT INGREDIENTS 90.8%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**FIRST AID:** If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

**DIRECTIONS FOR SWIMMING POOL CHLORINATION**

1. Add 1 lb. of this product to 10,000 gallons of water. For example, if you have 20,000 gallons of water, add 2 lbs. of this product. The amount of product to add will depend upon the size of the pool and the amount of water. Adjust pH of water to 7.2 to 7.6 and maintain. If you are using a feeder device, add this sodium hypochlorite solution to the feeder device. For each 1000 gallons of water, add 1 lb. of this product. Frequency of additions to maintain a residual chlorine will depend upon temperature and number of bather. To make certain pH and chlorine residual are in the proper range, test water frequently.

**DIRECTIONS FOR DISINFECTION OF POTABLE WATER FOR HOME WELL WATER SYSTEMS**

1. Add 1/2 cup of sodium hypochlorite solution in the ratio of one part sodium hypochlorite to 10 parts softened water. Mix sodium hypochlorite solution thoroughly and begin feeding of solution with a hypochlorinator. Maintain a free available chlorine residual of at least 0.2 ppm continuously throughout the distribution system as determined by a DPD chlorine test kit. Great sampling must be conducted at a frequency no less than that of Federal Interim Primary Drinking Water Regulations. Check water for DPD chlorine test kit.

**DIRECTIONS FOR SANITIZING FOOD PROCESSING EQUIPMENT**

1. Add 1/2 cup of sodium hypochlorite solution in the ratio of one part sodium hypochlorite to 10 parts softened water. Mix sodium hypochlorite solution thoroughly and begin feeding of solution with a hypochlorinator. Maintain a free available chlorine residual of at least 0.2 ppm continuously throughout the distribution system as determined by a DPD chlorine test kit. Great sampling must be conducted at a frequency no less than that of Federal Interim Primary Drinking Water Regulations. Check water for DPD chlorine test kit.

**NOTE:** This product degrades with age. Use a chlorine test kit and, if necessary, to obtain the required level of available chlorine.

**STORAGE AND DISPOSAL:**

Store away from direct sunlight. In case of spill, flood area with large quantity of water. Empty container thoroughly with water and either return to manufacturer, place in trash collection or burying in an approved landfill. Product cannot be used, should be diluted with water and disposed of in a sanitary manner. Do not use for storage, disposal or cleaning of equipment. **STORE IN AN UPRIGHT POSITION**